

21-11-6


74
227 Plate

6 Vols 15/-

Plates all in

53
47

85



FLORA LONDINENSIS

PLANTAE DESCRPTIONES

ENIRONS of LONDON

With a Description of the Trees and Shrubs of the City and Environs of London

By J. G. Smith, Esq. F.R.S. &c.

London: Printed by J. G. Smith, Esq. F.R.S. &c.

1794

Price 1s. 6d.

By J. G. Smith, Esq. F.R.S. &c.

London: Printed by J. G. Smith, Esq. F.R.S. &c.

1794

Price 1s. 6d.

By J. G. Smith, Esq. F.R.S. &c.

London: Printed by J. G. Smith, Esq. F.R.S. &c.

1794

Price 1s. 6d.

By J. G. Smith, Esq. F.R.S. &c.

London: Printed by J. G. Smith, Esq. F.R.S. &c.

1794

Price 1s. 6d.

By J. G. Smith, Esq. F.R.S. &c.

London: Printed by J. G. Smith, Esq. F.R.S. &c.

1794

Price 1s. 6d.

By J. G. Smith, Esq. F.R.S. &c.

London: Printed by J. G. Smith, Esq. F.R.S. &c.

1794

Price 1s. 6d.

By J. G. Smith, Esq. F.R.S. &c.

London: Printed by J. G. Smith, Esq. F.R.S. &c.

1794

Price 1s. 6d.

By J. G. Smith, Esq. F.R.S. &c.

FLORA LONDINENSIS:

OR

PLATES AND DESCRIPTIONS OF SUCH PLANTS

AS GROW WILD IN THE

ENVIRONS of LONDON:

WITH

Their Places of *Growth*, and Times of *Flowering*; their several
Names according to LINNÆUS and other Authors:

WITH

A particular DESCRIPTION of each PLANT in LATIN and ENGLISH.

To which are Added,

Their several Uses in *Medicine*, *Agriculture*, *Rural Oeconomy*, and other *Arts*.

By WILLIAM CURTIS,

DEMONSTRATOR of BOTANY to the COMPANY of APOTHECARIES.

VOL. I.



the hand of culture on popular minds
to plants in different beds, and to each
decides its province in the common soil
come by the hand
with more intent
what herding vases swell the leaves, veins
of herbs & flowers; or what the beams of sunny
dewy dew
she leads her vales & mountains to explore
through forth, distilling from the dewy road
in balmy tears

LONDON:

Printed for and Sold by the AUTHOR, No. 51, Gracechurch-Street; and B. WHITE, Bookfeller, in Fleet-Street.

MDCCLXXVII.

FLORA LONDINENSIS:

OR

PLANTS AND DESCRIPTIONS OF SUCH PLANTS

AS GROW WITHIN THE

ENVIRONS OF LONDON:

WITH

THE NAMES OF GROWERS, AND THE NAMES OF PLANTS, THEIR
VARIETIES, AND OTHER PARTICULARS.

WITH


A LIST OF THE NAMES OF THE PLANTS IN LATIN AND ENGLISH.

THE SECOND EDITION, CORRECTED, AND WITH ADDITIONS.

BY WILLIAM CURTIS,

PROFESSOR OF BOTANY IN THE UNIVERSITY OF LONDON.

VOL. I.



LONDON:

Printed for and sold by the Author, Mr. W. Curtis, in the Strand, near the Theatre Royal, and by Messrs. D. and J. Heath, in Pall Mall.

MDCCLXXXV.

THE

P R E F A C E.

ALTHOUGH the Author does not here mean to give a Preface at large, reserving that until the first volume, containing thirty-six numbers or two hundred and sixteen plants, shall be completed; yet he presumes it will be satisfactory to his subscribers, and the public, to be informed a little more fully of the nature and design of the work; as it will also give him an opportunity of answering some few objections that have been made to the plan of it.

The primary design of it then, is to facilitate a knowledge of the plants of our own country, and establish each species and variety on a firm basis: this the Author considers as the grand desideratum at present; this arduous task once accomplished, a way will be opened, and a foundation laid for numberless improvements in Medicine, Agriculture, &c.

To be enabled to do this, he means to take the greatest pains in the examination of those plants which he figures; to have them drawn from living specimens most expressive of the general habit or appearance of the plant as it grows wild; to place each plant, as much as is consistent, in the most pleasing point of view; and to be very particular in the delineation and description of the several parts of the flower and fruit, more especially where they characterize the plant.

And in order that he may obtain a more perfect knowledge of each plant; that he may see it in every stage of its growth, from the germination to the maturity of its seed; that he may compare and contrast the several species together; that he may make experiments to elucidate the nature of such as are obscure, or bring into more general use those which bid fair to be of advantage to the public; he is now cultivating each of them in a garden near the city, into which, by the kind assistance of his friends, he has already introduced, in the course of one year, about five hundred different species, including sixty of that most valuable tribe of plants the grasses.

Although the ascertaining and fixing of the plants will be his principal object, yet to make the work more useful to the public, as well as instructive and entertaining to the young botanist, his utmost endeavours will be used to lay before them whatever may be found useful in old botanic writers; and here they must not be surprized to find many of the numerous and imaginary virtues, which they attributed to almost every plant, purposely omitted: the discoveries made by modern authors, particularly relative to *Agriculture and Rural Economy*, will be carefully attended to; as here seems to be a field just opening to view, from whence the public is likely to draw great and lasting advantages: and as a knowledge of the plants themselves is first necessary, and for want of which, indeed, the experimental farmer cannot effectually communicate his improvements, he finds himself peculiarly happy in contributing his share to the public good.

He is nevertheless sensible how inadequate his abilities, or indeed the abilities of any one person are, to render a work of this kind any ways complete; he therefore respectfully solicits the assistance of those, who wish well to the improvement of *English Botany* and *English Agriculture*: any information they shall be pleased to communicate, shall with those favours he has already received from divers of his friends, be gratefully acknowledged; and to induce them the more readily to communicate, he has subjoined a catalogue of those plants which (with many others) are already drawn, and which he intends shall form the next Fasciculus.

He is sorry it has not been in his power to publish his numbers so fast as was originally proposed: the delay has chiefly been occasioned by the loss of one of his principal artists, whose place is now supplied by two others equally eminent; so that the drawing and engraving, which before fell to the share of one person, being now divided between two, he flatters himself he shall be able to publish a number once a month, or six weeks at farthest—he is however determined never to sacrifice the accuracy or utility of the work to hurry—on this principle he has been at the expence of having some of his plates engraven twice, and even three times over before he could venture to publish them. As the delay has originated from this source, he hopes none of his subscribers that have hitherto so generously contributed to the carrying on of the work, will withdraw that assistance, which alone can enable him to prosecute it with advantage to the public, credit to himself, and satisfaction to them.

It now remains to obviate some few objections which have been made to the plan of this work; and first, it has been suggested to the Author, that it would have been better received, if, instead of pursuing the present plan, he had published those plants only which were not figured in the *Flora Danica*, a work now carrying on in Denmark under the auspices of the King: but a few moments reflection, must he presumes be sufficient, to convince every unprejudiced person, how inadequate such a partial publication would have been to the making a knowledge of the plants of our country more general among ourselves—at best such a work could only answer the purpose of those few individuals who are in possession of that part of the *Flora Danica* already published; and as that is still going on, there is no doubt but the same plants would be published by both Authors; thus, the *Butomus umbellatus*, *Solanum Dulcamara*, and *Eryon hirsutum*, have been published in the *Flora Danica* since they were published in the *Flora Londinensis*, so that in the end even those persons would be obliged to purchase duplicates of the same plant.

Another reason why the Author could not adopt the plan proposed to him, was the limited scale of the *Flora Danica*, which contains the figures and names of the plants only, but gives us no account of their properties, nor teaches us how to distinguish the difficult plants from one another; the plates likewise being *small folio*, cannot admit many of the plants of their natural size, several of the grasses for instance, as the *Festuca sulcata* and *Aira aquatica* are obliged to be so cut and diminished as scarcely to be known. Many other objections might be urged without any view to depreciate a work which, though not so complete in some respects as could be wished, has exceeding great merit:—but these will probably be deemed sufficient.

The

THE PREFACE.

The engraving of one plant only on each plate has been another objection which some have strongly urged, while others have in as warm terms testified their approbation of it. It may be proper to mention, that whether one or more had been engraven on a plate, the difference in the expence would have been trifling, and chiefly in the paper: as they now are, each is distinct, and every one is at liberty to place them according to that system which he most approves of.

The want of figures of reference to the plates, or letter-press, has been perhaps a more solid objection; but the Author hopes, that by the use of the indexes described below, this also will be obviated.

Having now, so far as he can recollect, answered every thing deserving the name of an objection, he willingly submits his performance to the judgment of a candid and impartial public; conscious of having used his best endeavours to be serviceable in his department.

Uses of the Indexes, with Directions for Binding.

In the first Index the plants are placed according to the System of LINNÆUS, with which it is presumed, the greatest part of his subscribers are best acquainted. To find out any plant, even though the person be not acquainted with this mode of arrangement, look in the alphabetical English or Latin Index, and you will find the figures corresponding with them as placed in the book: if for example I want to find Ivy, I look for it in Index, No. 3, where the English names are alphabetically arranged, and find it to be the 16 plate, as there are 72 plates in each Fasciculus, I can readily guess within a few plates where it is placed: to those who have been accustomed to look out plants in LINNÆUS's works it will come easier; but if each subscriber will take the small pains of figuring the plates with a black lead pencil, any plant may then be immediately referred to. The Author could not hit on any mode more eligible, consistent with the irregular order in which he has been obliged to publish his plants.

With every third Fasciculus will be given a general and more copious Index, with a Glossary of the technical terms used in the work.

He would recommend to his subscribers, that each Fasciculus containing twelve numbers, be bound in boards, and not cut at the edges; the plates to be placed in the same order in which they occur in the first Index; taking care that each plate be put opposite to the letter-press belonging to it, with a leaf of thin paper betwixt them. If any should be at a loss to have them properly done, they will be pleased to send them to *Raham Reepe's*, Bookbinder, in Crooked Lane, near the Monument, who binds the Author's.

N. B. It may be necessary to caution the Bookbinder against beating the Numbers, as that operation would probably destroy the beauty of the plates.

A

Catalogue of those Plants which are *intended* to be Published in the next *Fasciculus*.

Anemone nemorosa
Adoxa moschatellina
Ajuga reptans
Aira præcox
Arabis thaliana
Arenaria tenuifolia
Achillæa Ptarmica
Briza media
Corylus avellana
Chærophyllum sylvestre
Convolvulus arvensis
Circæa lutea
Chenopodium Vulvaria
Dipsacus sylvestris
Ephedra angustifolium
Epilobium ramosum
Erica cinerea
Fumaria officinalis
Festuca duriuscula
Festuca myuros
Glechoma hederacea
Geranium molle
Geranium rotundifolium
Geranium perenne

Geranium Columbinum
Hyacinthus non scriptus
Hyoscyamus niger
Hypericum montanum
Hypericum quadrangulum
Hypericum hirsutum
Ilex Aquifolium
Iris Pseudacorus
Lamium amplexicaule
Lyfimachia nemorum
Lyfimachia nummularia
Lyfimachia tenella
Lyfimachia vulgaris
Ligustrum vulgare
Lotus corniculata
Myosurus minimus
Malva officinalis
Malva minor
Medicago lupulina
Osmunda spicant
Oxalis acetosella
Orchis Morio
Ornithopus perpusillus
Plantago lanceolata

Plantago major
Plantago Coronopus
Plantago media
Poa rigida
Poa compressa
Polygonum amphibium
Polytrichum commune
Ranunculus hirsutus
Ranunculus Ficaria
Sagina erecta
Saxifraga tridactylites
Spergula nodosa
Sedum dasyphyllum
Sedum reflexum
Symphytum officinale
Sparganium erectum
Tusillago farfara
Tormentilla erecta
Thymus serpyllum
Trifolium fragiferum
Valeriana dioica
Veronica officinalis
Veronica hederifolia
Veronica arvensis

I N D E X I.

In which the Plants contained in the first Fasciculus are arranged according to the
System of LINNÆUS.


<i>Latin Name.</i>	<i>Class and Order.</i>
1 Veronica agrestis	DIANDRIA Monogynia.
2 Veronica Chamædrys	
3 Veronica serpyllifolia	DIANDRIA Digynia.
4 Anthoxanthum odoratum	
5 Aira aquatica	TRIANDRIA Digynia.
6 Poa annua	
7 Festuca fluitans	
8 Bromus mollis	TETRANDRIA Monogynia.
9 Bromus sterilis	
10 Diplacus pilosus	
11 Hottonia palustris	PENTANDRIA Monogynia.
12 Anagallis arvensis	
13 Convolvulus sepium	
14 Solanum Dulcamara	
15 Lonicera Periclymenum	PENTANDRIA Digynia.
16 Hedera Helix	
17 Conium maculatum	PENTANDRIA Trigynia.
18 Æthusa Cynapium	
19 Scandix Anthriscus	OCTANDRIA Monogynia.
20 Alfine media	
21 Erica tetralix	OCTANDRIA Digynia aut Trigynia.
22 Polygonum Bistorta	
23 Polygonum Persicaria	
24 Polygonum Pensylvanicum	
25 Polygonum var. caule maculato	
26 Polygonum Hydropiper	
27 Polygonum aviculare	
28 Polygonum minus	ENNEANDRIA Hexagynia.
29 Butomus umbellatus	
30 Saxifraga granulata	DECANDRIA Digynia.
31 Sedum album	
32 Sedum acre	DECANDRIA Pentagynia.
33 Lychnis Flos Cuculi	
34 Cerastium aquaticum	DECANDRIA Pentagynia.
35 Euphorbia peplus	
36 Euphorbia Helioscopia	DODECANDRIA Trigynia.
37 Potentilla reptans	
38 Ranunculus bulbosus	ICOSANDRIA Polygynia.
39 Ranunculus acris	
40 Caltha palustris	POLYANDRIA Polygynia.
41 Verbena officinalis	
42 Lamium rubrum	DIDYNAMIA Gymnospermia.
43 Thymus acinos	
44 Euphrasia Odontites	
45 Antirrhinum Cymbalaria	DIDYNAMIA Angiospermia.
46 Antirrhinum Elatine	
47 Antirrhinum Linaria	
48 Digitalis purpurea	
49 Draba verna	TETRADYNAMIA Siliculosa.
50 Thlaspi Bursa-Pastoris	
51 Geranium Cicutarium	MONADELPHIA Decandria.
52 Geranium robertianum	
53 Orobancha tuberosus	DIADELPHIA Decandria.
54 Ervum hirsutum	
55 Ervum tetraspermum	POLYADELPHIA Polyandria.
56 Hypericum pulchrum	
57 Hypericum perforatum	SYNGENESIA Polygamia Æqualis.
58 Leontodon Taraxacum	
59 Lappula communis	SYNGENESIA Polygamia Superflua.
60 Erigeron acre	
61 Senecio vulgaris	SYNGENESIA Monogamia.
62 Bellis perennis	
63 Viola odorata	SYNGENESIA Monogamia.
64 Viola hirta	
65 Viola tricolor	GYNANDRIA Diandria.
66 Ophrys Apifera	
67 Asplenium Scolopendrium	CRYPTOGAMIA Filices.
68 Polypodium vulgare	
69 Bryum scoparium	CRYPTOGAMIA Musci.
70 Bryum undulatum	
71 Bryum hornum	
72 Hypnum proliferum	

I N D E X II. I N D E X III.

In which the Latin Names of the Plants
are arranged Alphabetically.

In which the English Names of the
Plants are arranged Alphabetically.

	Plate.		Plate.
Anthoxanthum odoratum	4	AIRA sweet tasted water	8
Aira aquatica	5	BROME-GRASS soft	8
Anagallis arvensis	12	BROME-GRASS barren	9
Alfane media	20	BRYUM broom	69
Antirrhinum Cymbalaria	45	BRYUM curled	71
Antirrhinum Elatine	46	BRYUM swan's-neck	22
Antirrhinum Linaria	47	BISTORT common	51
Asplenium Scolopendrium	67	CRANE'S-BILL hemlock-leaved	52
Aethusa Cynapium	18	CRANE'S-BILL stinking or herb Robert	51
Bromus mollis	8	CYMBALARIA Ivy-leaved	45
Bromus sterilis	9	CHICKWEED common	20
Butomus umbellatus	29	CONVOLVULUS large white	13
Bellis perennis	62	CINQUEFOIL common	37
Bryum scoparium	69	CROWFOOT round-rooted	38
Bryum undulatum	70	CROWFOOT upright meadow	39
Bryum hornum	71	DAISY, common,	62
Convolvulus Sepium	13	DRABA, vernal,	49
Conium maculatum	17	DEAD-NETTLE purple	42
Cerastium aquaticum	34	DANDELION common	58
C. ltha palustris	40	EYEBRIGHT red	44
Draba verna	49	ERIGERON purple	60
Digitalis purpurea	48	FLOWERING-RUSH	29
l. pilosus	10	FOOLS-PARSLEY	18
Erica tetralix	21	FLUELLIN sharp-pointed	46
Euphorbia Peplus	35	FOXGLOVE purple	48
Euphorbia Helioscopia	36	FESCUE-GRASS floc	7
Euphrasia Odontites	44	GROUNDSELL common	61
Eryum hirsutum	54	HARTS-TONGUE	67
Eryum tetraspermum	55	HEMLOCK	17
Erigeron acre	60	HEATH cross-leaved	21
Festuca nuntans	7	HOTTONIA water	11
Geranium cicutarium	51	HYPNUM proliferous	72
Geranium robertianum	52	HONEYSUCKLE common	15
Hottonia palustris	11	IVY	16
Hedera Helix	16	KNOT-GRASS common	27
Hypericum pulchrum	56	MOUSEEAR-CHICKWEED marsh	34
Hypericum perforatum	57	MARSH-MARIGOLD	40
Hypnum proliferum	72	NIPPLEWORT common	59
Lonicera Periclymenum	15	NIGHTSHADE woody	14
Lychnis Flos Cuculi	33	ORCHIS Bee	66
Lamium rubrum	42	PINK meadow	12
Leontodon Taraxacum	58	PEA wood	33
Lapfana communis	59	POA common dwarf	53
Orobis tuberosus	53	PERSICARIA spotted leaved	6
Ophrys apifera	66	PERSICARIA pale flowered	23
Poa annua	6	PERSICARIA spotted stalk'd	24
Polygonum Bistorta	22	PERSICARIA biting	25
Polygonum Persicaria	23	PERSICARIA small creeping	28
Polygonum Pensylvanicum	24	POLYTRIDY common	68
Polygonum var. caule maculato	25	PANSIE wild	65
Polygonum Hydropiper	26	SPURGE small garden	35
Polygonum aviculare	27	SPURGE sun	56
Polygonum minus	28	St. JOHN'S WORT small upright	36
Potentilla reptans	37	St. JOHN'S-WORT common	57
Polypodium vulgare	65	SCANDIX rough-seeded	19
Ranunculus bulbosus	38	SAXIFRAGE white	30
Ranunculus acris	39	STONECROP white flowered	31
Solanum Dulcamara	14	STONECROP common yellow	32
Scandix Anthriscus	19	SHEPHERDS-PURSE	50
Saxifraga granulata	30	SPEEDWELL procumbent garden	1
Sedum album	31	SPEEDWELL germander-leaved	2
Sedum acre	32	SPEEDWELL smooth-leaved	3
Senecio vulgaris	61	TINE-TARE rough, podded	54
Thymus acinos	43	TINE-TARE smooth podded	55
Thlaspi Bursa Pastoris	50	TEASEL small	10
Veronica agrestis	1	TOAD FLAX common yellow	47
Veronica Chamadrys	2	THYME basil	43
Veronica serpyllifolia	3	VERNAL-GRASS sweet-scented	4
Verbena officinalis	41	VERVAIN	41
Viola odorata	63	VIOLET sweet-scented	63
Viola hirta	64	VIOLET hairy	94
Viola tuedon	65		



VERONICA AGRESTIS. PROCUMBENT GARDEN-SPEEDWELL.

VERONICA *Linnaei. Gen. Pl. DIANDRIA MONOGYNIA.*

Raii. Syn. Gen. 18. HERBÆ FRUCTU SICCØ SINGULARI FLORE MONOPETALO.

VERONICA *agrestis*, floribus solitariis, pedunculatis; foliis cordatis incisus, petiolatis; caule procumbente.

VERONICA *agrestis*, floribus solitariis, foliis cordatis incisus pedunculo brevioribus. *Linnaei Syst. Vegetab. p. 56.*

VERONICA floribus solitariis, foliis cordatis incisus petiolatis. *Hudson Fl. Angl. p. 6.*

VERONICA caule procumbente; foliis petiolatis, ovatis, crenatis. *Haller. Hist. V. 1. n. 594.*

VERONICA *agrestis. Scopoli. Fl. Carn. p. 21* DIAGN. Primiflora; foliis ovato-cordatis, crenatis, pedunculo brevioribus.

VERONICA floribus singularibus, in oblongis pediculis, Chamædryfolia. *Raii. Syn. p. 279.* Germander-Speedwell or Chickweed.

ALSINE foliis Triflaginis: *Gerard. emac. 616. Parkinson. 764.*

ALSINE Chamædryfolia flosculis pediculis oblongis infidentibus. *Bauhin. Pin. 250. Oeder. Fl. Dan. Icon. 449.*

RADIX annua, fibrosa.

CAULES plures, primum erecti, tandem procumbentes, semipedales, subvillosi, teretes.

FOLIA alterna, ovato-cordata, serrata, petiolis brevibus infidentia, subhirsuta.

FLORES pedunculati, pedunculi axillares, longitudine fere foliorum, post florescentiam reflexi.

CALYX: PERIANTHIUM quadripartitum, laciniis lanceolatis, hirsutis, subtoruosis, fig. 1.

COROLLA monopetala, subrotata, calyce brevior, lævissimo fere tactu decidua; TUBUS brevissimus; LACINIÆ concavæ, subrotundæ, nunc penitus cœruleæ, nunc venis cœruleis striatæ, fig. 2.

STAMINA: FILAMENTA duo, alba, medio crassiora; ANTHERÆ cœrulescentes, fig. 3.

PISTILLUM: GERMEN subcompressum, hirsutulum, basi nectario cinctum; STYLUS viridis, apice incrassatus, filaminibus brevior; STIGMA album, capitatum, fig. 4.

PERICARPIUM CAPSULÆ *Veronica serpyllifolia* similis, at major rotundiorque, fig. 5.

SEMINA pallide fusca, plerumque 6 in singulo loculamento, rugosâ, hinc convexa, inde concava, fig. 6.

ROOT annual and fibrous.

STALKS several, first upright, then procumbent, about six inches in length, round and somewhat villous.

LEAVES alternate, of an oval-heart shape, serrated, placed on short foot-stalks and slightly hairy.

FLOWERS placed on foot-stalks, which proceed from the Axillæ of the leaves and are nearly of the same length; after the flowers are gone off turning back.

CALYX: a PERIANTHIUM divided into four lacinia, which are lanceolate, hairy, and somewhat twisted, fig. 1.

COROLLA monopetalous, somewhat wheel-shaped and shorter than the Calyx, falling off on the least touch; the TUBE very short; the LACINIÆ concave, and roundish, sometimes wholly blue, sometimes striped with blue, fig. 2.

STAMINA: two FILAMENTS of a white colour and thickest in the middle; ANTHERÆ bluish, fig. 3.

PISTILLUM: GERMEN flattish, a little hairy and surmounted at bottom by a Nectarium; the STYLE green, thickest at top, and shorter than the Stamina; STIGMA roundish and white, fig. 4.

SEED-VESSEL a CAPSULE like that of the *Veronica serpyllifolia*, but larger and rounder, fig. 5.

SEEDS of a pale brown colour, generally 6 in each cavity, wrinkled, convex on one side and hollow on the other, fig. 6.


THERE are few Botanists but what are apt to confound this species of Veronica with the *Veronica arvensis*, and this appears to arise in some degree from their similarity to each other, but more perhaps from the similitude of their Latin, and the ambiguity of their English names. To prevent in some degree this confusion, I have taken the liberty of altering the English name of *Germander-Speedwell* or *Chickweed* to that of *procumbent garden Speedwell*, in order that the young Botanist may thereby more readily distinguish it from the species above mentioned. The stalks of the *Agrestis* are usually procumbent, and it is found generally in Gardens; whereas the *Arvensis* has an upright stalk, and with us is found most commonly on Walls. Besides such obviously distinguishing characters, these two plants differ considerably in many other respects. In the *Arvensis* the leaves are sessile, in *this* they are placed on footstalks; in the *Arvensis* the flowers are sessile, in *this* species they likewise, are placed on foot-stalks: and a difference still more remarkable, or at least more curious, exists, which seems not to have been attended to, viz. the largeness and roundness of the seed-vessel, and the particular structure of the seed. In most of the Veronicas the seed-vessel is heart-shaped, and even in this species it retains somewhat of that form, although each of the Cavities is large and round; and if we examine the form of the seeds, we shall not wonder at this particular construction, for each seed instead of being small and flat as in other Veronicas, is large, convex on one side, hollow on the other, and wholly different in its appearance. This peculiarity of structure, shows what inconsistency there is in the parts of fructification, and how improper it would be to found a Genus on the particular form of any one of them, since those which are in general the most uniform, are sometimes subject to such uncommon variations. The number of seeds in each Capsule is generally about 12, LINNÆUS says 8, SCOPOLI from 16 to 20.

This species grows frequently in Gardens, and flowers through most of the summer months. No particular virtues or uses are attributed to it.




Veronica agrestis

A series of small, stylized botanical symbols arranged horizontally, likely representing different parts of the plant or its reproductive structures.





Veronica Chamuedrys. 

VERONICA CHAMÆDRYS. WILD GERMANDER.

VERONICA *Linnaei* Gen. Pl. DIANDRIA MONOGYNIA.

Raii Syn. Gen. 18. HERBÆ FRUCTU SICCO SINGULARI, FLORE MONOPETALO.

VERONICA *Chamædris* racemis lateralibus, foliis ovatis rugosis dentatis sessilibus, caule bifariam piloso. *Lin.*
Syll. Vegetab. p. 57. *Fl. Suecic.* p. 6.

VERONICA foliis cordatis subrotundis, hirsutis, nervosis, ex alis racemosa. *Haller. hist. n.* 536.

CHAMÆDRYS spuria minor rotundifolia. *Baubin. pin.* 249.

CHAMÆDRYS spuria sylvestris. *Parkinson,* 107.

CHAMÆDRYS sylvestris. *Gerard. emac.* 657. *Raii* Syn. 281. Wild Germander, *Hudson, Fl. Angl.* p. 5.
Scopoli. Fl. Carniol. p. 15. (a) *Oeder Fl. Dan. icon.* 448.

RADIX perennis, repens, fibrosa.	ROOT perennial, creeping, and fibrous.
CAULES numerosi, decumbentes, teretes, duri, bifariam dense hirsuti, ramosi.	STALKS numerous, spreading, round, hard, hairy on each side, hairs very thick together, branched.
FOLIA cordato-ovata, opposita, nunc sessilia nunc petiolis brevibus insidentia, serrata, venosa, hirsutula.	LEAVES of an heart shaped oval form, opposite, generally sessile, sometimes standing on short foot-stalks, serrated, veiny, and slightly hirsute.
FLORES numerosi, ad 20, cærulei, petiolati: Petioli BRACTÆA lanceolata suffulti; RACEMI longi, nunc oppositi nunc solitarii.	FLOWERS numerous, to 20, of a bright blue colour, forming long RACEMI (which are sometimes opposite, sometimes single), standing on foot-stalks, each of which is supported by a long-pointed BRACTÆA.
CALYX PERIANTHIUM quadripartitum, persistens, foliolis lanceolatis, hirsutulis, fig. 1.	CALYX: a PERIANTHIUM divided into four segments, and continuing, the segments lanceolate and slightly hairy, fig. 1.
COROLLA monopetala, rotata, tubus brevissimus internè ad inferiorem partem villosus, LIMBO quadripartito, plano, lacinis subcordatis ad basin venis saturationibus striatis, inferiore angustiore, fig. 2.	COROLLA monopetalous and wheel shaped, the TUBE very short, internally villous on the lowermost side, the LIMB flat, and divided into four segments, the segments somewhat heart-shaped, striated at bottom with veins of a purple colour, the lowermost segment narrower than the rest, fig. 2.
STAMINA: FILAMENTA duo apice incrassata, adscendentia, fig. 3. ANTHERÆ sagittata, fig. 4. POLLEN album, fig. 6.	STAMINA: two FILAMENTS, thickest at top, rising upward, fig. 3, the ANTHERÆ arrow-shaped, fig. 4. the POLLEN white.
PISTILLUM: GERMEN compressum glandula nectarifera cinctum: STYLUS declinatus, cærulefcens, STIGMA obtusum, purpureum, fig. 5.	PISTILLUM: the GERMEN flattish, surrounded at bottom by a nectariferous gland, fig. 6. the STYLE hanging downwards, bluish; the STIGMA blunt, and purple, fig. 5.
PERICARPIUM: CAPSULA cordata, subcompressa, pallide fusca, calyce paulo brevior, ad marginem hirsutulum, fig. 7.	SEED-VESSEL: a CAPSULE, heart-shaped, flattish, of a light brown colour, a little shorter than the calyx, and slightly hairy at the edge, fig. 7.
SEMINA: plura, compressa, flavescens, fig. 8.	SEEDS several, flat, of a yellowish brown colour, fig. 8.


The flowers of this Veronica are the largest and most specious of all the Plants of that Genus which grow wild in this Kingdom; many plants with less beauty are cultivated in our Gardens with the greatest care.

The leaves have been recommended by some writers as a substitute for Tea.

It bears a considerable resemblance to the *Veronica montana*, but differs essentially from that plant in the size of its Seed-vessels and the great number of flowers which it bears on its Racemi. See *Jacquin. Flor. Astricæ. Vol. 2.*

When growing wild the leaves are usually sessile or placed on very short foot-stalks, when cultivated they become larger and the foot-stalks moderately long; a kind of monstrosity, which *Linnaeus* has likewise observed, is very frequent on the leaves at the extremity of the stalk; which are collected into a very hairy white knob, on opening one of these I found two or three insects in their Pupa or Chrysalis state, which most probably would have produced some species of Fly. This appearance is very common at the latter end of Summer.

This is an early blowing plant, and grows very common on dry banks, under hedges, and in orchards; it flowers in May and June.



VERONICA SERPYLLIFOLIA. LITTLE SMOOTH SPEEDWELL, OR PAUL'S BETONY.

VERONICA *Linnei Gen. Pl.* DIANDRIA MONOGYNIA.

Raii Syn. Gen. 18. HERBÆ FRUCTU SICCO SINGULARI, FLORE MONOPETALO.

VERONICA *Serpyllifolia* racemo terminali subspicato, foliis ovatis glabris, crenatis. *Linnei Syst. Vegetab. p.* 56. *Fl. Suecic. p.* 6.

VERONICA caule recto, folijs ovatis, glabris, crenatis, petiolis ex alis unifloris, brevissimis, *Haller hist. n.* 546.

VERONICA pratensis serpyllifolia. *Baubin Pin.* 247.

VERONICA pratensis minor. *Parkinson.* 551.

VERONICA minor. *Gerard emac.* 627.

VERONICA foemina quibusdam, aliis Betonica Pauli Serpyllifolia. *I. Baubin. III.* 285.

VERONICA *Raii Syn. p.* 279. *n.* 3. *Hudson, Fl. Angl. p.* 4. *n.* 4. *Scopoli Fl. Carniol. V.1. p.* 12. *n.* 10. *Oeder Fl. Dan. icon.* 492.

RADIX perennis, fibrosum.

ROOT perennial, and very fibrous.

CAULES numerosi, ad basin repentes, dein erecti, simplices, palmares, teretes, laeves.

STALKS numerous, creeping at bottom, then growing upright, simple, three or four inches high, round and smooth.

FOLIA opposita, subconnata, subrotundo-ovata, *rariter et obsolete serrata, glabra, trinervia.*

LEAVES opposite, nearly uniting at bottom, of a roundish-oval form, here and there slightly serrated, smooth and trinervous.

FLORES albi, venis caeruleis picti, spicati, pedunculati, alterni, BRACTEÆ magnæ, ovatae.

FLOWERS white, coloured with blue veins or stripes, growing in spikes on foot-stalks alternately. FLORAL LEAVES large and oval.

CALYX: PERIANTHIUM quadripartitum, lacinii ovato-acutis, glabris, *fig. 1.*

CALYX: A PERIANTHIUM divided into four parts, the Segments of an oval pointed shape, and smooth, *fig. 1.*

COROLLA monopetala, rotata; tubus brevissimus; laciniae subcordatae, inferiore angustiore; superiore lacinia ftrius aut venis purpureis octo notata, lateralibus venis duabus, inferiore penitus alba, *fig. 2.*

COROLLA monopetalous, wheel-shaped, the tube very short, the segments somewhat heart-shaped, the lower one narrowest; the upper segment marked with eight purple veins or stripes, the side ones with two, and the lower one entirely white *fig. 2.*

STAMINA: FILAMENTA duo, alba, apice incrassata, *fig. 5, 6.* ANTHERÆ caeruleo-florescentes.

STAMINA: two FILAMENTS, white and thickish towards the extremity; the ANTHERÆ blueish *fig. 5, 6.*

PISTILLUM: GERMEN subcompressum, STYLUS albus, apice paululum incrassatus, persistens. STIGMA capitatum, rubens, *fig. 3.*

PISTILLUM: the GERMEN flattish, the STYLE white, a little thicker towards the extremity, and continuing. STIGMA roundish, and of a reddish colour, *fig. 3.*

NECTARIUM ad basin germinis, ut in Veronica Chamædryd.

NECTARY at the bottom of the Germen as in the Veronica Chamædryd.

PERICARPIUM: CAPSULA subcordata, fusca, pro magnitudine plantæ magna, *fig. 4.*

SEED-VESSEL: a CAPSULE somewhat heart-shaped, of a brown colour, and large in proportion to the plant, *fig. 4.*

SEMINA plurima, 60 numeravi, e luteo-fusca, sub-ovata, *fig. 8.*

SEEDS numerous, of a yellowish brown colour, and somewhat oval shape, *fig. 8.* We counted 60 in one capsule.

No particular virtues are attributed to this little plant by Writers.


It is one of the least of the Veronics, and occurs frequently in Meadows and Fields, and sometimes in Gardens, flowering in the Spring and Autumnal Months.

There is a great deal of delicacy in its blossoms, but they are too minute to make its beauty conspicuous enough for the Garden.

Its small, round, smooth and shining leaves readily distinguish it from the other Speedwells.



Thymus serpyllifolius



ANTHOXANTHUM ODORATUM. SWEET-SCENTED OR VERNAL GRASS.

ANTHOXANTHUM *Linnei Gen. Pl.* DIANDRIA DIGYNIA.

Calyx. Gluma bivalvis, uniflora. *Corolla.* Gluma bivalvis, acuminata. *Semen* unicum.

Raii Synop. Gen. 27. HERBÆ GRAMINIFOLIÆ FLORE IMPERFECTO CULMIFERÆ.

ANTHOXANTHUM *odoratum* spica oblonga, ovata, laxa.

ANTHOXANTHUM *odoratum* spica oblonga, ovata, flosculis subpedunculatis arista longioribus, *Linnei Syst.*

Vegetab. p. 67. Fl. Suecic. No. 33.

AVENA diantha, folliculo villoso, calycis glumis inæqualibus, altera de imo dorso, altera de summo aristata. *Haller. bist. belv. No. 1491.*

ANTHOXANTHUM *odoratum* *Scopoli Fl. Carniol. No. 38. Hudson Fl. Angl. p. 10. Stillingfleet miscel. t. 1. Schreber Gram. tab. 5. p. 49.*

GRAMEN *pratense* spica flavescente. *Bauhin. Pin. 3.*

GRAMEN *vernum* spica brevi laxa. *Raii Syn. 389. Scheuch. bist. 88.*

RADIX perennis, fibrosa.

CULMI primum obliqui, demum erecti, dodrantes aut pedales.

FOLIA inter digitos attrita odorem Asperulæ odoratæ spargunt, plerumque pubescentia, sæpe leniter tortuosa, membranâ ad basin instructa, Vagina striata, levis.

SPICÆ oblongo-ovate, laxæ.

CALYX: GLUMA bivalvis, Valvulis inæqualibus, inferiore dimidio brevior, membranacea, acuta, superiore acuminata, nervis tribus viridibus extantibus, *fig. 3, 2.*

COROLLA: GLUMA bivalvis, valvulæ subæquales, membranacæ, pilosæ aristatæ, fuscæ; altera Arista quæ demum geniculata fit, prope basin exurgit, altera prope apicem, *fig. 4.*

NECTARIUM: GLUMULÆ duæ, pellucidæ, nitidæ, ovate, inæquales, germen includentes, *fig. 5, 6.*

STAMINA: FILAMENTA duo prælonga; ANTHERÆ oblongæ, purpureæ, utrinque furcatæ, *fig. 5.*

PISTILLUM: GERMEN minimum oblongo-ovatum; STYLII duo filiformes glumi longiores, versus apicem plumulosi, *fig. 7.*

SEMEN unicum, Nectario fusco, nitido, inclusum, *fig. 8.*

ROOT perennial and fibrous.

STALKS at first growing obliquely, finally becoming upright, usually from 8 to 12 inches high.

LEAVES, if rubbed betwixt the fingers, smelling like Woodruff, generally pubescent and often curled, furnished with a membrane at bottom; the Sheath striated and smooth.

SPIKES of an oblong oval shape and smooth.

CALYX: a GLUME of two Valves, the Valves unequal, the lowermost shorter by one half, membranous and acute; the uppermost acuminate, with three strong nerves or ribs, *fig. 3, 2.*

COROLLA: a GLUME of two Valves, the Valves nearly equal, membranous, hairy, of a brown colour, and furnished with Arista, one of the Arista, which finally becomes bent, springs from the base of the Valve, the other almost at the top, *fig. 4.*

NECTARIUM: two small, pellucid, shining, oval, unequal Glumes or Valves inclosing the Germen, *fig. 5, 6.*

STAMINA: two FILAMENTS very long; ANTHERÆ long, purple, forked at each end, *fig. 5.*

PISTILLUM: GERMEN very small, of an oblong oval shape; STYLES two, slender, longer than the valves, and towards the top a little feathered, *fig. 7.*

SEED single, inclosed within its brown, shining Nectarium, *fig. 8.*

THE *Anthoxanthum* is distinguished from the other Grasses by a very singular circumstance, viz. that of having only two Stamina, *fig. 1.* hence it is placed by LINNÆUS among the *Diandria* plants, and separated from all the other Grasses; this peculiarity, although it occasions a separation which does violence as it were to Nature, yet it serves in a very striking manner to discriminate this Genus from a numerous and difficult tribe of plants: exclusive of this singularity, it differs also very essentially in the other parts of its fructification; each of the Spiculæ contains in common with many other grasses, only one flower, *fig. 1:* one of the Glumæ *Calyceinæ*, or valves of the Calyx, is small and membranous, *fig. 3;* the other is large, and incloses, or wraps up in it, as it were, the whole of the fructification, *fig. 2;* these glumes, so far as I have observed, do not open and expand themselves in the manner observable in the *Avena*, and many other grasses, were they separate quite wide, and expose their little feathery Styles; but the Stamina and Pistilla appear to push themselves out, the glumes remaining closed, *fig. 1.* The Glumæ *Cordulacæ* are very dissimilar to those of most other grasses, being remarkably hairy, and having each of them an Arista, the longest of which springs from near the base of the glume, is at first straight, but as the seed becomes ripe, the top of it is generally bent horizontally inward; the other Arista arises from near the top of the opposite Glume or Valve, *fig. 4.* The Glumule *Nectarii* or little Glumes of the Nectarium, differ no less in their structure, being composed of two little oval shining Valves, one of which is smaller than the other; these closely embrace the Germen, and cannot be seen but with great difficulty, until they are observed just at the time that the Antheræ are protruding from betwixt them, when they are very distinct, *fig. 6;* as soon as the Antheræ are excluded, they again close on the Germen, and continue to form a coat to the seed which does not separate, *fig. 5, 8.*

The Farmer or those who have not been accustomed to examine plants minutely, may readily distinguish this grass by its smell; if the leaves are rubbed betwixt the fingers, they impart a grateful odour like that of Woodruff,—hence I have called it sweet-scented.

Like the *Typhium repens* or *Irish Clover*, and many others of our most useful plants, this Grass grows on almost every kind of soil, from the poorest and driest, to the most fertile and boggy; it seems however in general to prefer a soil that is moderately dry. It is subject, like all other plants, to vary in its size, according to the good or bad of the ground it grows in: that is, we have a particular tendency to be curled if the soil be rich; and when it grows in woods, the spikes are often much broader and longer.

It has been called by some Authors *Percol* or *Spring Grass*, from its coming into ear earlier than most others; towards the middle or May it is in full bloom, and about the middle of June the seed is ripe—and may be easily separated on rubbing.

There is great reason to believe, that this is one of our Grasses which might be cultivated with considerable advantage: in the meadows about town it grows to a considerable height, and forms a thick turf of leaves at bottom; but the circumstance most in its favour, is its early appearance in the Spring: this seems to point it out as a proper grass to sow with others in laying down meadow land, and probably the *Poa trivialis* or common Meadow Grass, with the *Festuca elatior* or Meadow Fescue joined to it, would form a mixture, the produce of which, would for this purpose, be of great utility to most cities.



Anthoxanthum odoratum.





Tilia aquatica.

AIRA AQUATICA. SWEET-TASTED WATER AIRA.

AIRA *Linnei Gen. Pl.* TRIANDRIA DIGYNIA.

Cal. 2 valvis, 2 florus. *Flosculi* absque interjecto rudimento.

Raii Syn. Gen. 27. HERBÆ GRAMINIFOLIÆ FLORE IMPERFECTO CULMIFERÆ.

AIRA *aquatica* panicula patente, floribus muticis lævibus calyce longioribus, foliis planis. *Linnei Syst.*

Vegetab. p. 96. *Fl. Suecic. No.* 68.

POA locustis bifloris; glabris, florali gluma majori plicata, serrata. *Haller hist. No.* 1471.

AIRA *aquatica* Scopoli *Fl. Carniol.* 94. *Hudson Fl. Angl.* 29.

AIRA culmo inferiore repente, flosculis muticis calyce longioribus, altero pedunculato. *Ray. Lugdb.* 60.

GRAMEN caninum supinum paniculatum dulce. *Baubin Pin.* 2.

GRAMEN miliaceum aquaticum. *Raii Syn.* 402. *Scheuz. agr.* 218.

GRAMEN miliaceum fluitans suavis saporis. *Merret. Pin.*

RADIX perennis.

CULMUS basi repit, furculosque emittit more Festuæ fluitantis qui longe excurrunt et ad geniculos radículas plures albas dimitunt; culmus dumtaxat erigitur, pedalis circiter, teres, erectus, fistulosus, tener.

FOLIA latiuscula, tenera, lævia, carinata, vaginæ striatæ, ad basin rubræ præcipue in furculis.

PANICULA erecta, diffusæ, laxa, racemi plures ex uno puncto, sæpe flexuosi.

SPICULÆ plerumque biflores, flosculo uno sessili, altero pedunculato, purpurei, apicibus albidis, *fig. 1.*

CALYX: GLUMA bivalvis, valvulis inæqualibus, purpureis, lævibus, Corolla multo brevioribus, *fig. 2.*

COROLLA: GLUMA bivalvis, valvulis æqualibus, subtruncatis, plicatis five angulatis, *fig. 3.*

STAMINA: FILAMENTA tria capillaria, longitudine Corollæ; ANTHERÆ flavæ, *fig. 3.*

PISTILLUM: GERMEN ovatum; STYLUS duo plumosi, *fig. 4.*

NECTARIUM GLUMULÆ duæ minimæ ad basin Germinis, *fig. 5.*

SEMEN ovatum, intra Glumas arcte clausum, *fig. 7.*

ROOT perennial.

STALK creeps at bottom, and sends out young shoots like the Flote Fescue grass, which run out to a considerable distance, and send down small white roots at the joints; it then becomes erect, grows to about a foot in height, is round, hollow, and tender.

LEAVES broadish, tender, smooth, carinated, the sheaths striated, red at bottom, particularly in the young shoots.

PANICLE upright, spreading, loose; branches several, proceeding from one point, frequently crooked.

SPICULÆ generally contain two flowers, one of which is sterile, and the other stands on a foot-stalk, purple, the tips white, *fig. 1.*

CALYX: a GLUME of two valves the valves unequal, purple, smooth, and much shorter than the Corolla, *fig. 2.*

COROLLA: a GLUME of two valves, the valves equal, as if cut off at top, folded or angular, *fig. 3.*

STAMINA: three capillary FILAMENTS the length of the Corolla; ANTHERÆ yellow, *fig. 3.*

PISTILLUM: GERMEN oval; STYLES two and feathery, *fig. 4.*

NECTARY two very minute GLUMES at the bottom of the Germen, *fig. 5.*


SEED oval, closely contained within the Glumes, *fig. 7.*


The same soil and situation which produces the *Festuca fluitans*, is productive also of this grass; they both grow in gently flowing streams, or in wet boggy meadows; this circumstance may serve among others to distinguish the *Aira aquatica* from some of the *Poa's*, with which at first sight the young botanist might easily confound it: it has however besides this, many other characters which point it out more obviously. The bottom of the stalk usually creeps on the ground, and when it gets into the water, it runs out like the *Festuca fluitans* to a considerable distance, throwing off roots and young shoots as it passes along, very much in the manner of that grass: the stalk grows about a foot or more in height, is hollow, and remarkably tender; the leaves are broader than any of the *Poa's*, except the *Poa aquatica*, which is in every respect a much stronger plant: but what more especially characterizes this grass, is the purple or blueish colour of the Panicles, which is discernible even at a distance; and the sweet taste of the flowers if drawn through the mouth, whence this grass has acquired the name of *dulce*. Its parts of fructification likewise above described, distinguish it very strongly: when dried and placed between papers, the flowers and seeds are very apt to fall off.

It flowers in June and July, and may be found almost every where in the situations above-mentioned.

With respect to its uses in rural œconomy, it is in every respect inferior to the *Flote fescue grass*, consequently not worth cultivating for the use of cattle.

In a country like ours, where cultivation has made a considerable progress, the water plants are confined to a small space compared to what they occupied in a state of nature; the draining of bogs and lakes has rendered many large tracts in several parts of the kingdom, capable of producing corn and grass adapted to the use of cattle, which were formerly inaccessible to man or beast. We ought not however to look on this or any other plant as made in vain, because we do not immediately see the uses they are applied to: several sorts of water-fowl which abound in uninhabited countries, are expert gatherers of the seeds of the *aquatic grasses*; and no less than five different species of *Musci* or *Flies*, were produced from a few handfuls of the seeds of this grass, which when I gathered it, were doubtless in their Pupa or Chrysalis state: How little do we know of nature's productions!





POA ANNUA. COMMON DWARF POA.

POA *Linnaei* Gen. Plant. TRIANDRIA DIGYNIA.

Raii Synop. Gen. 27. HERBÆ GRAMINIFOLIÆ FLORE IMPERFECTO CULMIFERÆ.

POA *annua*, panicula diffusâ, angulis rectis, spiculis obtusis, culmo obliquo compresso. *Lin. Syst. Vegetab.*

p. 97. Spec. Plant. ed. 3. p. 99. Fl. Suecic. p. 228.

POA culmo infracto, panicula triangulari, locustis trifloris glabris. *Haller. hist. Vol. 2. p. 223.*

GRAMEN pratense paniculatum minus. *Bauhin. Pin. p. 2*

GRAMEN pratense minimum album et rubrum. *Gerard. emac. 3. Parkinson. 1156.*

GRAMEN pratense minus seu vulgatissimum. *Raii Synop. 408. Hudson. Fl. Angl. p. 34. Scopoli. Fl. Carniol. 71. Stillingfleet. tab. 7*

RADIX *annua*, fibrosissima.

CULMI plures, cespitosi, semiprocumbentes, in pratis vero inter alias plantas crescentes, suberecti, paululum infracti, semipedales.

VAGINÆ compressæ, acipites, læves.

FOLIA plurima, breviora, carinata, glabra, sæpe transversim rugosa, margine minutissime aculeata. *fig. 8.*

PANICULA triangularis, subcompressa, flores subsecundi.

PEDUNCULI *universales* ad basin paniculæ plerumque *bini*, altero breviora, in medio frequenter *terni*, apice vero *solitarii*; anguli nunc recti, nunc obliqui.

SPICULÆ ovato-acutæ, compressæ, utrinque acutæ trifloræ, quadrifloræ. *fig. 2.*

CALYX: GLUMÆ bivalvis, valvulis concavis, inæqualibus. *fig. 1.*

COROLLA bivalvis, valvulis villosis, margine membranaceis, albidis, una majore, concava, obtusifidula; alterâ minore, angustiore. *fig. 3.*

STAMINA: FILAMENTA tria capillaria; ANTHERÆ flavescentes, bifurcatæ. *fig. 4.*

PISTILLUM. GERMEN ovatum, STYLI duo ramossissimi, pellucidi. *fig. 5.*

SEMEN ovatum, corolla adnascente tectum, ad basin villosulum. *fig. 7.*

ROOT annual and very fibrous.

STALKS numerous, forming a turf, semiprocumbent, but in meadows when growing among other plants, nearly upright, a little crooked, and about half a foot high.

SHEATHS flat, two edged, and smooth.

LEAVES very numerous, short, keel-shaped, smooth, frequently wrinkled transversely, the edge very finely serrated. *fig. 8.*

PANICLE of a triangular shape and flatfish, the flowers growing mostly to one side.

PEDUNCLES: the *universal peduncles* generally proceed from the bottom of the panicle in *pairs*, one of which is shorter than the other, from the middle often by *threes*, and at top *singly*; forming angles sometimes straight, sometimes oblique.

SPICULÆ oval and pointed, flatfish and sharp on both sides, containing three and four flowers. *fig. 2.*

CALYX: a GLUME of two valves, the valves hollow and unequal. *fig. 1.*

COROLLA of two valves, the valves villous, membranous and whitish at the edges, the one larger, hollow and bluntish, the other smaller and narrower. *fig. 3.*

STAMINA: the FILAMENTS very minute, the ANTHERÆ yellowish and forked. *fig. 4.*

PISTILLUM: the GERMEN oval, two STYLES exceedingly ramified and pellucid. *fig. 5.*

SEED oval, covered by the Corolla which adheres to it, at bottom slightly villous. *fig. 7.*

THE laudable Society established in London for the encouragement of Manufactures, Arts, and Commerce, sensible of the improvements which might be made in Agriculture, from a more general introduction of the most useful English Grasses, have offered Premiums to such as shall give the best account of their cultivation, and the *Poa Annua* above described, is one of those they have selected, from its appearing to them to be one of the most useful.

Mr. Stillingfleet observes that it makes the finest turf, that he has seen in high Suffolk whole fields of it, without any mixture of other Grasses, and that as some of the best salt Butter we have in London comes from that County, he apprehends it to be the best Grass for the Dairy; from observing likewise, that this Grass flourished much more from being trodden on, he concludes that frequent rolling must be very serviceable to it.

There is no Grass better entitled to Ray's epithet of *Vulgatissimum* than this, as it occurs almost every where, in Meadows, Gardens, at the sides of Paths, and on Walls: when it grows in a very dry situation, it frequently doth not exceed three inches, but in rich meadows it often grows more than a foot in height. The panicle is frequently green, but in open fields it acquires a reddish tinge; it flowers all the Summer long, and even in Winter if the weather be mild.

It appears to be the first general covering which Nature has provided for a fruitful soil when it has been disturbed; for which reason, in Walks, Pavements, or Pitching, it may be considered as one of the most troublesome of Weeds; the most expeditious method of destroying it, would probably be by pouring boiling water on it.

All the Authors that have described this Grass call it an annual, it differs however very considerably from the other annual Grasses, they throw up their Spikes or Panicles, produce their flowers and seeds, and then die away; this on the contrary keeps continually throwing out new shoots, and producing new flowers, and seeds, and if the ground be moist, a single plant will remain growing in this manner throughout the year, so that we generally find on the same plant, young shoots and ripe seeds.

"Hic ver æfiduum atque alienis mensibus æstas."

Perhaps this is the only vegetable we have that in this Circumstance imitates the Tropical plants.

Although its seed may be gathered the whole summer long, yet about the latter end of May, it will be found in the greatest plenty: Experience must determine the best method, in which this Grass should be cultivated, whether by sowing its seed, or dividing and transplanting the Grass itself; as this seed would with more difficulty be procured in large quantities than that of many others, and as a single tuft of this Grass may be divided into a vast number of young plants, probably transplanting it in wet weather would be the most eligible mode of cultivation.

These observations are submitted to the consideration of the Farmer and Gentlemen of landed property, who reside in the Country, and who have both leisure and opportunity to try experiments of this kind. Although the Authors province more particularly is to describe and figure these plants in such a manner as to make them as obvious as possible, yet he would be exceedingly happy to communicate to the public, any improvements which may be made in this or any other branch of Agriculture, that he may be favoured with.

FESTUCA FLUITANS. FLOTE FESCUE GRASS.

FESTUCA *Linnei Gen. Pl. TRIANDRIA DIGYNIA:*

Raii Gen. 27. HERBE GRAMINIFOLIE FLORE IMPERFECTO CULMIFERE.

FESTUCA panicula ramosa erecta, spiculis subsessilibus, teretibus muticis. *Linnei Syst. Vegetab. p. 107.*

Fl. Suecic. p. 32.

POA locustis teretibus multifloris, glumis floralibus exterioribus truncatis, interioribus bifidis. *Haller. kist. p.*

219. n. 1453. v. 2.

POA fluitans. *Scopoli Fl. Carniol. p. 73.*

GRAMEN aquaticum fluitans, multiplici spica. *Baubin Pin. 2.*

GRAMEN aquaticum cum longissima panicula. *I. Baubin. II. 490. Raii Syn. p. 412. Flore-Grafs.*

GRAMEN fluviatile. *Gerard emac. 14. Parkinson. 1275. Hudson. Fl. Angl. p. 38. Oeder. Fl. Dan. i. 237.*

Schreber. Gram. tab. 3. Stillingfleet. mis. tab. 10.

RADIX perennis, in limbum profunde penetrans.
CULMUS pro ratione loci pedalis ad tripedalem, basi repens furculosque promens, dein suberectus, vaginis foliorum ad paniculam usque amictus.

VAGINÆ foliorum compressæ, subancipites, striatæ.

FOLIA latiuscula, lævia; furculorum erecta, carinata, breviuscula, caulina longiora, planiuscula, flaccida, aquis tempore hyberno prostrata.

PANICULA longa, inclinata, nonnunquam subspicata sæpius vero ramosa, ramis nunc cauli adpressis nunc distantibus, ut pinxit C1: Schreberus.

SPICULÆ tenues, teretes, unciales aut sesquicunciales 9 ad 12 floræ, rachi adpressæ.

CALYX: GLUMA bivalvis, valvulis inæqualibus, membranaceis. *fig. 2.*

COROLLA bivalvis, valvulæ longitudine æquales, calyce majores, inferiore majore, concava, lineata, nervis apice sæpe coloratis, apice membranacea, obtusiuscula, sæpius erosa; superiori lanceolata, compressa, bicaudata. *fig. 3. 4.*

STAMINA: FILAMENTA tria capillaria, ANTHERÆ flavæ aut purpurascens, oblongæ, *fig. 5.*

PISTILLUM: GERMEN ovatum, STYLII duo subulati, reflexi, STIGMATA ramossissima. *fig. 7. 6. 8.*

NECTARIUM Glandula squamiformis, cordata, horizontalis, ad basin germinis. *fig. 9.*

SEMEN oblongum, nitidum olivaceum, bicorniculatum, nudum. *fig. 10. 11.*

FIG 12 Spicula morbo *Ergot* affecta.

ROOT perennial, striking deep into the mud.

STALK according to its place of growth from one to three feet in length, creeping at bottom and sending forth young shoots, afterwards nearly upright; covered with the sheaths of the leaves as far as the panicle.

SHEATHS of the leaves, flattened, two edged, and striated.

LEAVES rather broad and smooth, those of the young shoots upright, keel-shaped, and shortish; those of the stalk longer, flattish, weak, and hanging down, in the winter season lying flat on the water.

PANICLE long, generally inclined or bending down a little, sometimes forming a kind of spike, but most commonly branched; the branches sometimes pressed to the stalk, sometimes diverging from it in the manner represented by Schreber.

SPICULÆ slender, round, an inch or an inch and a half long, producing from 9 to 12 flowers, pressed to the stalk.

CALYX: a GLUME of two valves, which are unequal and membranous. *fig. 2.*

COROLLA of two valves, which are of an equal length and bigger than the Calyx, the lower valve largest, concave and nervous, the nerves towards the top frequently coloured, at top membranous, rather blunt with uneven points, the upper valve more pointed, flat and bifid. *fig. 3. 4.*

STAMINA: three FILAMENTS very slender, ANTHERÆ oblong and yellow or purplish. *fig. 5.*

PISTILLUM: GERMEN oval, STYLES two, tapering and bending back, STIGMATA very much branched. *fig. 7. 6. 8.*

NECTARY a small heart-shaped squamiform gland, placed horizontally at the bottom of the Germen. *fig. 9.*

SEED oblong, shining, of an olive colour, with two little horns, and naked. *fig. 10. 11.*

FIG 12 a Spicula affected with the disease called *Ergot*.

IN speaking of the *Bromus mollis*, we had occasion to remark the very great variety of appearance to which the Grasses were subject from soil and situation, and this observation is equally applicable to the *Festuca fluitans*.

This Grass appears to thrive best in still waters, or gently running streams, where its numerous fibres penetrate easily into the mud; in such situations it becomes very luxuriant, the leaves are large, tender and sweet, and the Panicle becomes very much branched; but in Meadows where it is deprived of its natural quantity of water, it becomes in every respect less, and the Panicle is frequently changed to a simple spike; when it has nearly done flowering, the branches of the Panicle generally project from the main stalk so as to form an acute angle. In every situation whether the Panicle be large, or small, the Spiculæ are always pressed close to the stalk or branches of the Panicle, and this circumstance joined to the length, and roundness of the Spiculæ, sufficiently characterize this species; if it should not however, its parts of fructification afford at once a most pleasing and satisfactory distinction, *vid. fig. 6. 9. 10.*



Distachya

spicata.

We have often had the singular pleasure of observing this Grass soon after being gathered, expand its Glumes and expose its delicate yellow Stamina, and still more delicate Pistilla, and in this expanded state each Spicula puts on a very different face, and seems to invite the Student to its investigation, and would he wish to become acquainted with the structure of this useful tribe of plants, he cannot select one more proper for his purpose, as it may be found in almost every watery ditch, flowering from the beginning to the end of Summer, and has all the parts of fructification which are peculiar to the Grasses, large enough to be distinctly discerned even by the naked Eye, and so exposed as to be visible without the trouble of dissection.

Modern Botanists seem much divided whether they should consider this as a *Poa* or *Peplosia*, as it does not appear to us that we should in the least advance our favourite Science by altering its generic name, we have continued that of LINNÆUS, although we are by no means satisfied with his generic characters of the Grasses in general, and are persuaded that future observations and a more accurate attention to the minute parts of their fructification, will place those Genera in a much clearer point of view than has yet been done by any author.

Professor OEDER in his *FLORA DANICA*, and the celebrated SCHREBER in his *AGROSTOGRAPHIA*, have both given a figure of this grass. As we have not seen it growing either in Denmark or Germany we cannot say that their figures do not express its particular mode of growth in those countries, but they do not convey to us its habit or manner of growing here; in both their figures the Panicle is represented quite upright, whereas with us it is always more or less inclined; this however is a matter of no great moment, a deviation from nature in the representation of the minute parts of the fructification is a matter of much greater consequence, and we are sorry to find that Mr. SCHREBER whose knowledge and accuracy can seldom be called in question, has not been sufficiently attentive to all the parts which characterize this species. He has represented the Styles as branched or feathered quite down to the Germen, whereas they are evidently naked at bottom and much branched at top only; the singular Squamula or Scale at the base of the Germen he has properly noticed, but the two little Horns at the top of the seed, which are the remains of the Styles, and which in a peculiar manner distinguish this important seed, he does not remark. In the *Flora Danica* the Styles are likewise feathered down to the Germen and the Squamula at the base of the Germen wholly omitted.

This Grass is found to be of considerable importance in the economy of Nature.

The *Phalena Fefcuse* or *Gold Spot Moth*, to which LINNÆUS with great propriety adds the epithet of *puleherrima*, *Coel. Funna Suecica*. p. 311. *Albin. pl. 81. lit. K. r. o. n.* is said by him to feed on this particular species; with us however it is always found on a different grass, viz. the *Poa aquatica* or *Large water Poa*; its history, with the particular manner of finding it we shall give under that grass.

From the observations of late writers, it appears that several sorts of Cattle are remarkably fond of this grass, particularly Kine and Hogs, and that in the spring time they are frequently enticed into bogs by endeavouring to get at its sweet young shoots, which appear earlier than those of most other Grasses.

“Professor KATM is a journey through part of Sweden, observed the Swine to go a great way into the water after this grass, the leaves of which, they eat with great eagerness; on this he was tempted to try if they would eat the same grass dried; he accordingly had small bundles of it gathered, dried, and call before them; the consequence was they ate it seemingly with as much appetite as horses do hay, hence he concludes that by cultivating this grass, wet and swampy places might be rendered useful, and a great deal of corn, &c. saved”.

He who introduced the method of feeding hogs in summer time on Clover, deserved very well of his country; and if the hay of this grass would keep them in heart during the winter, it might prove a very valuable discovery.

Mr. Kent in his *hints to Gentlemen of landed property*, lately published, considers this as a most valuable grass, and assures us (p. 24) it is to be improved above all others, and at a less expence, merely by flooding; (p. 54,) he informs us that flooding destroys all weeds, and enriches the land to a very high degree; (p. 56,) he says as rolling and pressure bring the annual meadow-grass, so flooding immediately begets the *slote fescue*. These assertions of Mr. Kent bespeak neither the Philosopher nor the accurately practical Farmer, they contain an exaggerated account of improving pasture land by a particular process, but show a great want of that minute attention which so important a subject required.

From a long residence in Hampshire, we well know that the meadows in that county are considerably improved by flooding them, that is stopping the water when there happens to be an unusual quantity from violent or long continued rains, and by means of trenches or gries, conveying the surplus water so as to overflow them entirely if possible; but we deny, that by this process all weeds are destroyed, the use of manure superadded, or that *slote fescue* grass is immediately begotten. Although it is a constant practice with the farmers to flood their meadows in the winter, it is no less a constant practice with such as wish to have good crops of grass to manure them with dung or ashes. Flooding can no otherwise destroy weeds than by altering the soil in which they grow, and if it destroys one set of weeds, it must certainly favour the growth of another: if those plants which thrive best in a dry situation are destroyed by the alteration which now takes place in the soil, those which are fond of a moist situation will proportionably flourish. If the *slote fescue* grass was immediately produced by flooding, we should find all those meadows which have undergone this operation to contain nothing but this kind of grass, whereas the richest and best meadows in Hampshire contain scarce a single blade of it: the fact is, this grass will not flourish in meadow land, unless you convert it into a kind of bog or tump, and I believe few landed Gentlemen will think this an improvement, or thank Mr. Kent for giving them such a hint.

“Mr. Stillington informs us that Mr. Deane a very sensible Farmer at Ruscomb, in Berkshire, assured him, that a field always lying under water of about four acres, that was occupied by his father when he was a boy, was covered with a kind of grass that maintained five farm-horses in good heart from April to the end of harvest without giving them any other food, and that it yielded more than they could eat. He at my desire brought me some of the grass, which proved to be the *slote fescue* with a mixture of *maish bent*; whether this last contributes much towards furnishing good pasture for horses I cannot say, they both throw out roots at the joints of the stalks and therefore likely to grow to a great length. In the index of dubious plants at the end of Ray's *Synopsis*, there is mention made of a grass under the name of *Grassum caninum supinum longissimum* growing not far from Salisbury twenty-four feet long; this must by its length be a grass with a creeping stalk: add that there is a grass in Wiltshire, growing in watery meadows, so valuable that an acre of it lets from ten to twelve pounds, I have been informed by several persons. These circumstances incline me to think it must be the *slote fescue*; but whatsoever grass it be it certainly must deserve to be enquired after”.

It may not be improper to add, that the account of the extraordinary long grass above mentioned, was taken by Ray from the *Phytographia Britannica*, which mentions the particular spot where it grew, viz. at Mr. Tucker's, at Maddington, nine miles from Salisbury; it is also remarked that they fat Hogs with it.

As it is now above a century since this enquiry was first made, is it not surprising that no succeeding Botanic Writer should have acquired satisfactory information concerning it? I am promised specimens of the roots and seeds. Upon

Upon the whole, from the observations which we ourselves have made on this Grass and from what is to be collected from Authors, it appears that if it be cultivated to any advantage it must be in such meadows as are naturally very wet and never drained.

The quickest and perhaps the best method of propagating it would be by transplanting the roots at a proper season, and if the soil prove suitable, from the quickness of its growth, and its creeping Stalk, it would soon exclude most other plants, and produce a plentiful crop.

In foreign countries the seed of this Grass seems to be an object of more importance than the grass itself, the following is the substance of what Mr. SCHREBER has said concerning it, (vid. *Beschreibung der Gräser* p. 40.)
 "The seed has a sweet and pleasant taste particularly before it comes to its full growth, whence the plant has acquired the name of *Manna Grass*. Ducks and other water-fowl feed on it with much eagerness (LINNÆUS has remarked that the Water-fowl are very well acquainted with the method of collecting these seeds) it has been observed likewise that Fish are fond of it, and that Trout in particular thrive in those rivers where this grass grows in plenty and sheds its seeds; but it is not only for Birds and Fish but also for Man a palatable and nutritious food, and has for many years past been known at Gentlemen's tables under the name of *Manna-Grout*.

"The Manna Grass is of two kinds the one *Panicum sanguinale* or *Cocks-foot*, *Panic-Grass* the other *Festuca fluitans* which we have now described; the former is cultivated in several parts of Germany, and its seed somewhat resembles that of Millet, the latter is collected in great abundance from the plant as it grows wild in Poland, Lithuania, the new Marche and about Frankfort and other places in *Silesia* as also in Denmark and Sweden and hence exported to all parts.

"The common method they make use of to gather and prepare this seed in Poland, Prussia, and the Marche is as follows. At sun rise the seed is gathered or beat from the dewy grass into a horse-hair sieve, and when a tolerable quantity is collected, it is spread on a sheet and dried fourteen days in the sun; it is then thrown into a kind of wooden trough or mortar, straw or reeds laid between it, and beat gently with a wooden Pestle so as to take off the chaff and then winnowed. After this it is again put into the mortar, in rows, with dried Marygold-flowers, Apple, and Hazel leaves, and pounded until the Hulk is entirely separated and the seed appears bright, it is then winnowed again, and when it is by this last process made perfectly clean it is fit for use. The Mary-golds are added with a view to give the seeds a finer colour. The most proper time for collecting them is in July. A Bushel of the seed and chaff, yields about two quarts of clean seed.

"When boiled with milk or wine they form an extremely palatable food, and are most commonly made use of whole in the manner of *Sago* to which they are in general preferred.

In the month of October last, I discovered in a watery ditch, which runs through a meadow not far from Kent-Street Road an uncommon appearance in some of the seeds of this grass, and on a farther examination, I found whole Panicles the seeds of which were affected in a similar manner, instead of being of their natural size, and colour, they were enlarged to a very great degree, assumed externally a blackish colour, and were more or less incurved. Struck with the novelty as well as oddity of the appearance I conjectured at first that it was a disease occasioned by some Insect, I examined it more attentively, but could not find the least cause to suppose that an Insect had been concerned in it. The surface of some of these seeds was rough, and chopped, they were light as to weight, internally of a whitish colour, insipid in their taste, but not disagreeable. Having a little before this been favoured with a sight of some *burned Rie* it now occurred to me that this was the same disease which had been said to affect the *Rie* only, and farther enquiry confirmed my conjecture.

As this singular disease of the *Rie* has first been noticed by the French, and as some very uncommon circumstances have attended it, it cannot fail of proving acceptable to our readers to lay before them the substance of what they have said concerning it. In the *Histoire de L'Academie royale des Sciences* there is an account given of a particular species of Gangrene or Mortification which attacked many persons in some particular provinces of France. It began generally at the toes and sometimes spread as high as the thigh. Out of fifty people there was but one that was attacked with this disease in the hands and what was equally remarkable there were no females affected with it except some little Girls.

"It appears that this singular malady attacked only the lower sort of people, and that too in years of scarcity, that it proceeded from bad nourishment, and principally from eating bread made of a certain black and diseased corn called *Ergot*, from the grains assuming somewhat of the form of a Cocks Spur. vid. fig. 12.

"The manner in which this singular monstrosity of the Corn is produced is thus related by Monsieur FAGON.

"There are certain mists which prove injurious to the corn, and from which the greatest part of the Ears of the *Rie* defend themselves by their beards. In those however which this hurtful humidity can strike and penetrate, it rots the skin which covers the grain, blackens it, and alters the substance of the grain itself, the juices which form the seed being no longer kept within their ordinary bounds by the skin, are carried hither in two great an abundance and amassing themselves irregularly form this monstrous appearance.

"He observes that it is only in *Rie* that the *Ergot* is to be found, that the poor people do not separate this grain from that which is good, that it was only in such particular seasons as favoured the growth of the *Ergot* that this disease was prevalent, that the country people after eating bread made of this bad corn perceived themselves as if drunk, and after this the mortification generally took place, that in some provinces where there was but little of this *Ergot* this species of disease was not known.

"From the observations made by the Farmers of that country it appears that this bad species of grain is produced in the greatest abundance in such land as is wet and cold, and particularly in rainy seasons. The Poultry refused it when given them, nevertheless if by accident they had eaten it they did not appear to be hurt by it, When sown (as might be expected) it did not vegetate."

A kind of mortification very similar to the above described was observed in this Kingdom some years ago; it affected the same kind of people and on enquiry it was found that they had sared very hard, and that the bread which they had eaten was made of the *tailings* or *screenings* of Corn, but it was not ascertained whether it contained any of the *Ergot* or not.

From the insipid taste of this corn, as well as from its not proving fatal to Poultry, it seems exceedingly probable that it is not in itself noxious, any otherwise than as it affords no nourishment; and that those people who have eaten of this corn, have in fact been abridged of a proportionate quantity of food, hence from an impoverished state of the fluids and a weak action of the vessels this species of Mortification might easily be induced.

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO
LIBRARY
1100 EAST 58TH STREET
CHICAGO, ILL. 60637

1967-1968

BROMUS MOLLIS. SOFT BROME GRASS.

BROMUS *Linneæ Gen. Pl.* TRIANDRIA DIGYNIA.

Raii Syn. Gen. 27. HERBÆ GRAMINIFOLIÆ FLORE IMPERFECTO CULMIFERÆ.

BROMUS *mollis* panicula erectiuscula, spiculis ovatis pubescentibus, aristis rectis, foliis mollissime villosis

Linneæ Syst. Vegetab. p. 102. *Sp. Pl. p.* 112.

BROMUS hirsutus, locustis septifloris, ovato concis. *Haller hist. p.* 1504.

BROMUS Polymorphus. *Scopoli Fl. Carniol. p.* 80.

FESTUCA avenacea hirsuta, paniculis minus sparsis. *Raii Synop. p.* 413 *Hudson Fl. Angl. p.* 39. n. 1.

Secalinus. Schreber. Gram. pl. 6. fig. 1.

RADIX	biennis *	ROOT	biennial *
CULMUS	pedalis ad tripedalem, erectus; GENICULI tumidi, cylindracei.	STALK	from one to three feet high, upright; the JOINTS swelled and cylindrical.
FOLIA	cum VAGINIS pilis mollibus vestita.	LEAVES	together with their SHEATHS covered with soft hairs.
PANICULA	erectiuscula, nunc coarctata nunc diffusa.	PANICLE	nearly upright, sometimes close, sometimes spreading.
SPICULÆ	ovato-acutæ, turgidæ, subcompressæ, plerumque villosæ, octofloræ, circa oras glumarum albidæ. <i>fig. 1.</i>	SPICULÆ	oval and pointed, turgid, flattish, generally villous, containing eight flowers, whitish about the edges of the Glumes. <i>fig. 1.</i>
CALYX:	GLUMA bivalvis, valvulis inæqualibus, muticis. <i>fig. 2.</i>	CALYX:	a GLUME of two valves, the valves unequal without any beard, or arista, <i>fig. 2.</i>
COROLLA:	GLUMA bivalvis, valvulâ exteriorē latâ, concavâ, striatâ, aristatâ, <i>fig. 4.</i> interiorē planiusculâ, ciliatâ, lanceolatâ. <i>fig. 3.</i> ARISTA valvulis paulo longior, subrectâ, <i>fig. 4.</i>	COROLLA:	a GLUME of two valves, the outermost valve broad, hollow, striated, and bearded, <i>fig. 4</i> ; the innermost flattish, ciliated or hairy at the edges and pointed, <i>fig. 3</i> ; the ARISTA a little longer than the valves and nearly straight, <i>fig. 4.</i>
NECTARIUM:	Glumula bipartita, ad basin petali interioris, <i>fig. 5.</i> parum auct:	NECTARIUM:	a small kind of Glume deeply divided, placed at the base of the inner petal, <i>fig. 5.</i> a little magnified.
STAMINA:	FILAMENTA tria capillaria, ANTHERÆ primum flavæ, oblongæ, dein fuscæ et bifurcatæ. <i>fig. 7.</i> 6. auct:	STAMINA:	three FILAMENTS very small, ANTHERÆ first yellow and oblong, lastly brown and forked at each end, <i>fig. 7.</i> 6. magnified.
PISTILLUM:	GERMEN ovatum, apice subemarginatum, <i>fig. 8.</i> STYLI duo, ad basin usque plumosi, ex uno latere germinis enati. <i>fig. 9.</i>	PISTILLUM:	GERMEN oval, with a slight depression at top, <i>fig. 8.</i> two STYLES feathery quite down to the bottom, proceeding from one side of the Germen, <i>fig. 9.</i>
SEMEN	oblongum, concavum, calyci adnatum <i>fig. 10.</i> denudatum <i>fig. 11.</i>	SEED	oblong, concave, adhering to the Calyx <i>fig. 10.</i> the Calyx taken off, <i>fig. 11.</i>


OUR Farmers in general are not very warm in their recommendations of this Grass, nevertheless it abounds in most of our best meadows; it springs up early, and ripens its seed generally about the time of Hay-making. The seed is large, and each panicle contains nearly as much as that of a common Oat, indeed it seems to have more pretensions to the name of Corn than of Grass.

Although Cattle may not be so fond of the leaves, and panicle of this Grass while green as of some others, yet may it not (when cut down as it usually is when the seed is nearly ripe) contribute to render the hay more nutritive? and hence may it not be a proper Grass to sow with others? It seems at least to deserve the attention of the Farmer.

There is perhaps no class of plants more affected by difference of soil and situation than the Grasses, hence the same plant has often been divided into several species; and to such Varieties is the present Plant incident, as to occasion SCOPOLI to give it the name of *Polymorphus*.


When it grows on a Wall, or dry Bank, the Spiculæ are generally more upright, and closer together; when the soil is rich and moist, the Spiculæ spread out, and the whole plant becomes much larger; in Meadows the Spiculæ frequently lose their villous appearance and become perfectly smooth. To determine this species then with more certainty, recourse must be had to the parts of fructification.

* According to LINNÆUS



Bromus Hollii







Bromus sterilis

BROMUS STERILIS. BARREN BROME GRASS.

BROMUS *Linnaei*. *Gen. Pl.* TRIANDRIA DIGYNIA.

Raii Gen. 27. HERBÆ GRAMINIFOLIÆ, FLORE IMPERFECTØ CULMIFERÆ.

BROMUS *sterilis*, panicula patula, spiculis oblongis distichis, glumis fubulato-aristatis. *Lin. Syst. Vegetab.* p. 103.

BROMUS panicula nutante; locustis septifloris; glumis argute lanceolatis, lûneatis, subhirsutis. *Haller. hist. n.* 1505.

FESTUCA avenacea sterilis elatior. *Baubin. pin.* 9. 16.

BROMOS herba, sive avena sterilis. *Parkinson, 1147.* Bromos sterilis. *Gerard. emac. Raii Synop.* p. 412. Great wild Oat-Grass or Drank. *Hudson. Fl. Angl.* p. 40. *Scopoli. Fl. Carniol.* p. 76.

RADIX fibrosa.

CULMI pedales ad bipedales, subrecti, teretes, læves, ad basin infracti; Geniculi tumidi.

FOLIA longa, plana, unâ cum vaginis molliſſime villosa.

PANICULA magna, nutans; PEDUNCULI plerumque simplices, ad basin tumidi.

SPICULÆ binucleales, subcompressæ, apice divergentes, fig. 1.

CALYX: GLUMÆ bivalvis, Valvulis inæqualibus lineari-lanceolatis, fig. 2.

COROLLA: bivalvis, Valvulis inæqualibus, exteriore longiore, concava, striata, apice membranacea, bifida, ARISTA recta Corollâ duplo longiore terminata, fig. 3. Valvulâ interiore planiuscula, ciliata, fig. 4.

NECTARIUM: GLUMULÆ duæ acuminatæ, ad basin biglandulosæ, fig. 6.

STAMINA: FILAMENTA tria, capillaria, ANTHERÆ flavæ, fig. 5.

PISTILLUM: GERMEN oblongum, apice truncatum sive emarginatum, pars inferior ex quâ styli prodeunt et quod verum Germen esse videtur, nitida, fig. 7. pars superior albida, villosa, fig. 8. STYLI duo plumosi, patuli, fig. 9.

SEMEN ex purpureo-fuscum, oblongum, aristatum, calyce tectum, fig. 10, denudatum, fig. 11.

ROOT fibrous.

STALKS from one to two feet high, nearly upright, round and smooth, at bottom crooked or elbowed; the Joints swelled.

LEAVES long and flat, covered, together with their sheaths with soft short hairs.

PANICLE large, and drooping, the PEDUNCLES generally simple, and swelled at their base.

SPICULÆ about two inches long, flatfish and diverging toward the extremity, fig. 1.

CALYX: a GLUME of two Valves, the valves unequal, long and narrow, fig. 2.

COROLLA: composed of two Valves, which are unequal, the exterior Valve longest, concave, striated, at top membranous and bifid, terminated by a straight ARISTA twice the length of the Corolla, fig. 3. the interior Valve nearly flat, and ciliated, fig. 4.

NECTARY: two small long-pointed GLUMES with a small gland at the base of each, fig. 6.

STAMINA: three small FILAMENTS: the ANTHERÆ yellow, fig. 5.

PISTILLUM: the GERMEN oblong, at top flat or slightly emarginate, the bottom part from whence the Styles proceed, and which seems to be the true Germen, is smooth and shining, fig. 7. the upper part white and villous, fig. 8. two STYLES, feathery and spreading, fig. 9. SEED of a purplish brown colour, oblong, bearded, enclosed within the Calyx, fig. 10. the Calyx stripped off, fig. 11.

Much praise is due to the late ingenious Mr. STILLINGFLEET for his attempts to introduce, more generally among Farmers, a knowledge of our most useful English Grasses: his observations on this subject are so exceedingly pertinent that the insertion of them cannot fail to prove highly acceptable to such as have the promotion of Agriculture at heart.


"It is wonderful to see how long mankind has neglected to make a proper advantage of plants of such importance, and which in almost every country are the chief food of cattle. The farmer for want of distinguishing, and selecting grasses for feed, fills his pasture either with weeds, or bad, or improper grasses; when by making a right choice, after some trials he might be sure of the best grass, and in the greatest abundance that his land admits of. At present if a farmer wants to lay down his land to grass, what does he do? he either takes his seeds indiscriminately from his own foul hay-rick, or sends to his next neighbour for a supply. By this means, besides a certain mixture of all sorts of rubbish, which must necessarily happen; if he chances to have a large proportion of good seeds, it is not unlikely, but that what he intends for dry land may come from moist, where it grows naturally, and the contrary. This is such a slovenly method of proceeding, as one would think could not possibly prevail universally; yet this is the case as to all grasses except the darnel grass, and what is known in some few countries by the name of the Suffolk grass; and this latter instance is owing, I believe, more to the soil than any care of the husbandman. Now would the farmer be at the pains of separating once in his life half a pint, or a pint of the different kinds of grass seeds, and take care to sow them separately; in a very little time he would have wherewithal to stock his farm properly, according to the nature of each soil, and might at the same time spread these seeds separately over the nation by supplying the seed-shops. The number of grasses fit for the farmer is, I believe small; perhaps half a dozen, or half a score are all he need to cultivate; and how small the trouble would be of such a task, and how great the benefit, must be obvious to every one at first sight. Would not any one be looked on as wild who should sow wheat, barley, oats, rye, pease, beans, vetches, buck-wheat, turnips and weeds of all sorts together? yet how is it much less absurd to do what is equivalent in relation to grasses? does it not import the farmer to have good hay and grass in plenty? and will cattle thrive equally on all sorts of food? we know the contrary. Horses will scarcely eat hay, that will do well enough for oxen and cows. Sheep are particularly fond of one sort of grass, and fatten upon it faster, than on any other in Sweden, if we may give credit to Linnæus. And may they not do the same in England? How shall we know till we have tried? Nor can we say that what is valuable in Sweden may be inferior to many other grasses in England; since it appears by the Flora Suecica that they have all the good ones that we have. But however this may be I should rather chuse to make experiments, than conjectures."

The present Grass is not one of those which are worth the Farmer's cultivation, but so much the reverse, that most Authors have called it *sterilis*, not because it is really barren but from its inutility with respect to Cattle.

It grows exceeding common under hedges and flowers in May and June.

In order to have a clear idea of the structure of the parts of fructification in the Grasses, they should be examined just at the time, or rather before the *Antheræ* have discharged their *Pollen*, a small space of time makes a considerable alteration in their appearance.

In this species of *Bromus* as well as in the *Bromus mollis* the Styles proceed from the middle of the Germen and not from the top, this is a peculiarity which seems to have escaped the notice of *Schreber* who has written professedly on the Grasses, and examined them with more accuracy than any preceding Writer. In his figures the Styles proceed always from the Apex of the Germen.





Lysimachia pilosula.

DIPSACUS PILOSUS. SMALL WILD TEASEL OR SHEPHERD'S R O D.

DIPSACUS *Linnaei Gen. Pl.* TETRANDRIA MONOGYNIA.

Calyx communis polyphyllus; proprius superus. Receptaculum paleaceum.

Raii Syn. p. 191. HERBÆ CORYMBIFERIS AFFINES.

DIPSACUS *pilosus* foliis petiolatis appendiculatis. *Linn. Syst. Vegetab. p. 120. Spec. Plant. 145.*

DIPSACUS foliis biauribus, capitulis hemisphæricis. *Haller. hist. herb. No. 199.*

DIPSACUS sylvestris capitulo minore vel virga pastoris minor. *Bauhin Pin. p. 385.*

DIPSACUS minor seu Virga pastoris. *Ger. emac. 1168.*

VIRGA PASTORIS. *Parkinson 984. Raii Synop. p. 192. Hudfon. Fl. Angl. p. 49.*

RADIX biennis.

ROOT biennial.

CAULIS orgyalis, erectus, ramosissimus, pene teres, aculeatus, fuscatus.

STALK about six feet high, upright, very much branched, nearly round, prickly and grooved.

RAMI oppositi, patentes, cauli similes.

BRANCHES opposite, spreading, like the stalk.

FOLIA ad basin CAULIS, connata, ovato-lanceolata, ferrata, nervo medio fubtus aculeato, indivisa, suprema appendiculata; RAMORUM; ima appendiculata, ferrata, suprema margine integerrima, lanceolata.

LEAVES at the bottom of the STALK connate, ovato-lanceolate, ferrated, the midrib prickly underneath, undivided, those at the top dividing at the base into two smaller leaves; the leaves on the branches at bottom similar to those last described, at top lanceolate, with the edges entire.

PEDUNCULI erecti, longi, ex dichotomiâ caulis, fulcati, aculeati, apice spinosissimi, uniflori.

FOOT-STALKS of the flowers, upright, long, proceeding from the middle where the stalks separate, grooved, prickly, at top very full of slender spines, supporting one flower.

FLORES albi, in capitulum hemisphæricum collecti, dum florent nutantes, postea capitula eriguntur.

FLOWERS whitish, collected together in a small hemispherical head, which, while the plant is in flower, droops, and afterwards becomes upright.

CALYX: PERIANTHIUM commune multiflorum, hexaphyllum, foliolis longitudine florum, patentibus, lanceolatis, mucronatis, fig. 1: PERIANTHIUM proprium parvum, superum, concavum, ciliatum, fig. 5. lente auctum.

CALYX: the common PERIANTHIUM supporting many flowers, composed of six leaves, the length of the flowers, spreading, lanceolate and pointed, fig. 1. The PERIANTHIUM of each floret small, placed above the Germen, hollow, and ciliated, fig. 5, magnified.

COROLLA propria monopetala, tubulosa, limbo quadrifido, lacinia inferiore longiore, fig. 3.

COROLLA: each floret monopetalous, tubular, the limb quadrifid, the lowermost segment longest, fig. 3.

STAMINA: FILAMENTA quatuor Corollâ longiora; ANTHERÆ purpureæ, fig. 3.

STAMINA: four FILAMENTS, longer than the Corolla; ANTHERÆ purple, fig. 3.

PISTILLUM: GERMEN inferum, tetragonum; STYLUS filiformis, longitudine Corollæ; STIGMA simplex, fig. 6.

PISTILLUM: GERMEN placed below the Calyx, quadrangular; the STYLE filiform, the length of Corolla; the STIGMA simple, fig. 6.

PERICARPIUM nullum.

SEED-VESSEL wanting.


SEMINA fusca, subtetragona. fig. 4.

SEEDS brown, nearly quadrangular.

RECEPTACULUM commune hemisphæricum, paleaceum, pars inferior palearum concava, alba, carinata, superior lanceolata, acuminata, spinulis obtusa. fig. 2.

RECEPTACLE common to all the flowers paleaceous; the lower part of the paleæ hollow, white, and angular behind; the upper part lanceolate, tapering to a point, and beset with little spines or hairs, fig. 2.

THIS species of *Teasel* may be considered as one of our *Plantæ rariores*; hitherto I have found it only in one place near town, viz. on the right hand side of the Turnpike-road leading from *Deptford* to *Lewisham*, not far from the latter: as it grows to a considerable height, it is conspicuous at a distance: the flowers appear in July, and the seed is ripe in September: it continues to blow for a considerable time, and did not the plant take up so much room, there is beauty enough in its flowers to recommend it for the Garden. Moths seem very fond of its blossoms, being found on them in great numbers after sun-set.



HOTTONIA PALUSTRIS. WATER HOTTONIA, OR WATER VIOLET.

HOTTONIA *Lin. Gen. Pl.* PENTANDRIA MONOGYNIA.

Raii Syn. Gen. 18. HERBÆ FRUCTU SICCO SINGULARI, FLORE MONOPETALO.

HOTTONIA *palustris*, pedunculis verticillato-multifloris. *Lin. Syst. Vegetab.* 164.

HOTTONIA florum verticillis spicatis. *Haller. bist. n.* 632.

MILLEFOLIUM aquaticum seu Viola aquatica, caule nudo. *Baubin. pin.* 141. *Parkinson,* 1256.

VIOLA palustris. *Gerard. emac.* 826. *Raii Syn. p.* 285. *Hudson. Fl. Angl. p.* 72. *Scopoli Fl. Carniol. n.* 213. *Fl. Dan. icon.* 487.

RADIX e plurimis fibrillis capillaceis albis constat, quæ in limbum profunde dimittuntur.

ROOT consists of numerous white capillary fibres, which penetrate deep into the mud.

CAULIS sive SCAPUS floriferus, pedalis, simplex, erectus, multiflorus, versus apicem glandulis scabriusculus, ad basin foliis plurimis instructus, unde per aquam longe excurrunt caules plures qui fibrillas dimittunt.

STALK or flowering SCAPUS, a foot high, simple, upright, sustaining many flowers, towards the top roughish with little glands, furnished at bottom with numerous leaves, from whence several stalks proceed and run out to a considerable length through the water throwing out numerous white fibres.

FOLIA plurima, plerumque immersa, pinnata, in apicibus caulium juniorum densa, reflexa, Pinnis linearibus planis.

LEAVES numerous, generally under the water, pinnated, growing in tufts on the tops of the young stalks, bending downwards, the Pinnæ linear and flat.

FLORES pallide purpurei, verticillati, spicati, Pedunculi ad 10, Bractæa, ad basin instructi, post florescentiam reflexi.

FLOWERS of a pale purple colour, growing in whorls, and forming a spike. Peduncles to 10 in number, furnished at bottom with a Bractæa, when the flowers are gone off turning downwards.

CALYX: PERIANTHIUM monophyllum, quinquepartitum: LACINIIS linearibus, erecto-patulis, *fig. 1.*

CALYX: a PERIANTHIUM of one leaf, divided into five segments, which are linear, upright and somewhat spreading, *fig. 1.*

COROLLA: monopetala, hypocateriformis, TUBUS longitudine calycis, LIMBUS quinquefidus, planus: LACINIIS ovato-oblongis, emarginatis, *fig. 2.*

COROLLA: monopetalous and salver-shaped, the TUBE the length of the calyx; the LIMB divided into five segments and flat; the SEGMENTS of an oval oblong shape with a notch at the extremity, *fig. 2.*

STAMINA: FILAMENTA quinque, subulata, brevia, erecta. ANTHERÆ oblongæ, flavæ, *fig. 3.*

STAMINA: five FILAMENTS tapering, short, and upright, ANTHERÆ oblong and yellow, *fig. 3.*

PISTILLUM: GERMEN subglobosum. STYLUS filiformis, brevis. STIGMA globosum, *fig. 4.*

PISTILLUM: GERMEN roundish, STYLE thread-shaped and short, STIGMA spherical, *fig. 4.*

PERICARPIUM: CAPSULA globosa, unilocularis, subpellucida, *fig. 5.*

SEED-VESSEL: a round CAPSULE of one cavity, slightly transparent, *fig. 5.*

SEMINA plurima, ovata, pallide fusca, *fig. 7.* receptaculo globoso intra capsulam affixa, *fig. 6.*

SEEDS numerous, oval, of a pale brown colour, *fig. 7:* affixed to a round receptacle within the capsule, *fig. 6.*

This singular plant abounds in most of our watry Ditches, particularly in such as divide the Meadows, and flowers in May and June, continuing for a considerable time in blossom; among a variety of other places it may be found in a ditch on the right hand side of the Field Way leading from Kent-street Road to Peckham.

We do not find any author that mentions its possessing any properties to recommend it but its beauty and singularity, both of which it possesses in a degree sufficient to command our admiration.

The leaves generally grow beneath the surface of the water and afford a Nidus if not Nourishment to the fresh-water Periwinkle and some other small shell fish.

Antient Botanists have given it the names of *Millefolium aquaticum*, and *Viola aquatica*; the great number of its leaves induced them, with some propriety, to call it *Millefolium*, but why they should call it a *Viola* seems difficult to determine, as the blossom has nothing in its structure similar to the flowers of that Genus. Boerhave afterwards called it *Hottonia*, in honour of Dr. Hotton, which name Linnæus has continued.



Helianthus patens.

AGRICULTURE - LAND - CULTIVATION

1. AGRICULTURE - LAND - CULTIVATION

2. AGRICULTURE - LAND - CULTIVATION

3. AGRICULTURE - LAND - CULTIVATION

4. AGRICULTURE - LAND - CULTIVATION

5. AGRICULTURE - LAND - CULTIVATION

ANAGALLIS ARVENSIS. PIMPERNEL.

ANAGALLIS *Linnei Gen. Plant.* PENTANDRIA MONOGYNIA.

Raii Gen. 18. HERBÆ FRUCTU SICCO SINGULARI, FLORE MONOPETALO.

ANAGALLIS foliis indivisis caule procumbente. *Lin. Spec. Plant.* 211.

ANAGALLIS phœniceo flore. *Baubin. pin.* 252.

ANAGALLIS mas *Fuschii* 18. *Gerard emac.* 617. *Parkinson* 558. *Order. Flor. Dan. tab.* 88. *Raii Syn.* 282. *Hudson.* 73. *Haller. hist.* 621. 626. *Scopoli. Fl. Carniol.* 139.

RADIX simplex, fibrosa, annua.

CAULIS ramosus, prostratus, quadrangularis, laevis, subtoruosus, *fig. 1.*

FOLIA opposita, sessilia, cordata, glabra, *subtus punctis fuscis notata.*

PEDUNCULI oppositi, foliis fere duplo longiores, inflexi.

CALYX persistens, quinquepartitus, segmentis triangularibus, alatis, membranaceis, *fig. 2.*

COROLLA monopetala, quinquepartita, laciniiis rotundis, coccineis, ad basin purpureis, margine crenatis, subpilosis, *fig. 3-4.*

STAMINA. FILAMENTA quinque, erecta, *filosissima (pili articulati!)* superne purpurea. ANTHERÆ oblongæ, biloculares, flavæ, infidentes, *fig. 5-6.*

PISTILLUM. GERMEN rotundum, STYLUS filiformis, *obliquus*, longitudine filamentorum; STIGMA subrotundum, *extra circulum filamentum locatum,* *fig. 7.*

PERICARPIUM. CAPSULA rotunda, nitida, quinquecervia, subdiaphana, circumscissa, fusca, *fig. 8.*

SEMINA plurima, angulosa, fusca, *fig. 9.*

ROOT simple, fibrous, annual.

STALK branched, procumbent, quadrangular, smooth, and a little twisted, *fig. 1.*

LEAVES opposite, sessile, heart-shaped, smooth, underneath dotted with brown.

PEDUNCLES opposite, nearly twice the length of the leaves, bending downwards.

CALYX persisting, divided into five segments, the segments triangular and membranous at the edges, *fig. 2.*

COROLIA monopetalous, quinquepartite, the laciniae scarlet purplish at bottom, the edges slightly notched, and hairy, *fig. 3-4.*

STAMINA. five FILAMENTS upright and very hairy (the hairs when magnified joined!) at top purplish: The ANTHERÆ oblong, bilocular, yellow, and sitting on the filaments, *fig. 5-6.*

PISTILLUM. the GERMEN round, the STYLE filiform, the length of the filaments, the STIGMA roundish, placed without the circle of the Stamina, *fig. 7.*

SEED-VESEL. a CAPSULE, round, shining, brown, slightly transparent, having five nerves, dividing into two equal parts, *fig. 8.*

SEEDS numerous, brown, and angular, *fig. 9.*

NATURE seems to have taken uncommon pains in the formation of the Flowers of this little Plant; few possess more liveliness of colour or greater delicacy of structure, this must be sufficiently obvious to every common observer, but when its minute parts come to be viewed by the microscope, we are charmed with beauties altogether novel and unexpected; we then find that the edges of the flowers, which to the naked eye appear a little uneven or hairy, are furnished with a number of little glands placed on footstalks, and that the hairs of the Filaments which partly tend to distinguish this Genus, are regularly jointed: the Pistillum which generally arises upright betwixt the Stamina, is here inclined to one side, so that the Stigma is placed without the circle of the Stamina: The care which Nature has taken likewise in the preservation of these delicate parts from the injury of the weather, is not less remarkable. Every morning, if the weather be fair and warm, the blossoms fully expand, but if rain falls, or there be much moisture in the air, the flowers quickly close themselves up to secure the inclosed Antheræ and Stigma, from having their functions destroyed: from this property which it has in common with many plants of the same class, it has acquired the name of the *Shepherd's*, or *Poor Man's Weather-glass*,—they have remarked that if the flowers be open in a morning it will prove a fine day, if shut, the contrary.

The small birds (PASSERES LINNEI) are fond of the seeds of this Plant; and according to experiments made by some of LINNÆUS's pupils, it appears that kine and goats feed on it.

It is very common in corn fields and gardens, flowering all the Summer. A variety with four leaves at a joint sometimes occurs in rich soil; but as it differs in no other part, and is a mere variety, it scarcely deserves a distinct figure. It is also found with blue and sometimes with white flowers, but we have not observed either of these varieties near London.



Anagallis arvensis.

1870-1871

1871-1872

1872-1873

1873-1874

1874-1875

1875-1876

1876-1877

1877-1878

1878-1879

CONVOLVULUS SEPIUM. LARGE WHITE CONVULVULUS OR GREAT BINDWEED.

CONVOLVULUS *Linnæi. Gen. Pl. PENTANDRIA MONOGYNIA.*

Raii Syn. Gen. 18. HERBÆ FRUCTU SICCO SINGULARI FLORE MONOPETALO.

CONVOLVULUS (*sepium*) foliis sagittatis, postice truncatis, pedunculis tetragonis, unifloris. *Linn. Syst. Vegetab. p. 168. Fl. Suecic. p. 64.*

CONVOLVULUS foliis sagittatis, hamis emarginatis, angulosis, petiolis unifloris, stipulis cordatis maximis. *Haller. bist. V. 1. p. 295.*

CONVOLVULUS Major albus. *Baubin. pin. 294.*

SMILAX laevis major. *Gerard emac. 861. Parkinson. 163. Raii Syn. p. 275. Great Bindweed. Hudson. Fl. Angl. p. 74. Scopoli. Fl. Carniol. 141. Fl. Dan. icon. 458.*

RADIX perennis, crassitie pennæ aserinae, alba, sub terra reptans et late se propagans, vix eradicanda, Hortorum pestis.

CAULES numerosi, volubiles, tortuosi, striati, orgyales, subramosi.

RAMI pauci, alterni, cauli similes.

FOLIA alterna, sagittata, postice truncata, glabra, petiolata.

PEDUNCULI uniflori, alterni, tetragonis.

CALYX INVOLUCRUM biphylum, foliolis oblongo-cordatis, subcarinatis, venosis, purpurascensibus. *fig. 2.*

CALYX PERIANTHIUM pentaphyllum, tubulosum, foliolis ovato-lanceolatis, pallide virentibus. *fig. 1.*

COROLLA monopetala, infundibuliformis, lactea, limbo lato, obscure diviso, paululum reflexo.

STAMINA: FILAMENTA quinque, fundo corollæ inferta, hirsutula, alba, subulata; ANTHERÆ sagittatæ, albæ, insidentes. *fig. 3.*

PISTILLUM: GERMEN subovatum; STYLUS subulatus apice tortuosus; STIGMA bifidum. *fig. 4. 5.*

NECTARIUM: Glandula crocea annuliformis ad basin Germinis.

PERICARPIUM: CAPSULA subrotunda, fuliginosa, mucronata. *fig. 6. 7.*

SEMINA angulosa, fusca, Cotyledonibus mire convolutis. *fig. 8. 9.*

ROOT perennial, about the thickness of a goose quill, of a white colour, creeping under the ground and propagating itself exceedingly, rooted out with the greatest difficulty, and hence very troublesome in Gardens.

STALKS numerous, twining, twisted, striated, generally about six feet high and somewhat branched.

BRANCHES few, alternate, like the Stalk.

LEAVES alternate, arrow-shaped, apparently cut off behind, smooth, and placed on foot-stalks.

FOOT-STALKS of the flowers, alternate, supporting one flower only, and four square.

CALYX an INVOLUCRUM composed of two heart-shaped leaves, slightly keel-shaped, veiny, and purplish. *fig. 2.*

CALYX a PERIANTHIUM, composed of five leaves and tubular, the leaves of an oval pointed shape and pale green colour. *fig. 1.*

COROLLA monopetalous, funnel shaped, of a white colour, the limb broad, obscurely divided, and turned back a little.

STAMINA: five FILAMENTS inserted into the bottom of the corolla, slightly hairy, white and tapering, the ANTHERÆ arrow shaped, white, and sitting on the filaments. *fig. 3.*

PISTILLUM. GERMEN somewhat oval, STYLE tapering, twisted at top; the STIGMA bifid. *fig. 4. 5.*

NECTARY a yellow gland surrounding the base of the Germen.

SEED-VESSEL a roundish CAPSULE of a sooty colour and pointed. *fig. 6. 7.*

SEEDS angular and brown, the Cotyledons folded up in a very singular manner. *fig. 8. 9.*


The plant which produces the *Scammony* is a species of *Convolvulus*, very similar to that which we have now described, hence Dr. Cullen and some other Physicians have conjectured that our *Convolvulus* might possess similar properties, but if it should be found to contain such properties, the smallness of its roots would prevent its juice from being collected in the same manner with that which flows on incision from the large root of the *Scammony* plant, and which hardens and forms that purgative substance. Whether an extract made from the expressed juice of the roots, or any other preparation of them might possess a purgative property, or if it should, whether such a purgative would be so far superior to any now in general use as to introduce it deservedly into practice, is what we cannot pretend to decide on. Hogs are said to eat and even to be fond of the roots.

It grows exceedingly common in our hedges, and flowers in August and September. Where it has once gained ground it is with the greatest difficulty eradicated: was it not for this property and its being so common, it would doubtless be considered, as it really is, a very ornamental plant.

My ingenious Friend Mr. CHURCH, Surgeon, at Islington, (who has taken much pains to collect and acquire a knowledge of our English Insects) informs me that the Caterpillar of the *Phalena Vivicaria* or *Bloody vein Moth*, (*vid. Clerc. Phalen. pl. 3. fig. 2.*) feeds on this plant, and the *Sphinx Convolvuli* or *Unicorn Hawk Moth*, (*vid. Roefel. Cl. 1. pap. noct. 1. 7.*) is well known to take its name from feeding on this plant also.



Convolvulus sepium.





Solanum Dulcamara.

SOLANUM DULCAMARA. WOODY NIGHTSHADE.

SOLANUM *Linnei Gen. Pl.* PENTANDRIA MONOGYNIA.

Raii Gen. 16. HERBÆ BACCIFERÆ.

SOLANUM *Dulcamara* caule inermi frutescente flexuoso, foliis superioribus hastatis, racemis cymosis. *Lin. Sp. Pl.* 264.

SOLANUM Scandens seu *Dulcamara Bauhin. Pin.* 176. Amara Dulcis. *Gerard. emac.* 350. Solanum lignosum *Parkinson. 350. Raii Synopsis* 265. *Hudson. Flor. Angl.* p. 78. *Scopoli Flor. Carniol.* p. 161. *Haller. bist. Plant. Helv.* p. 248.

RADIX perennis.

CAULIS fruticosus, scandens, fistulosus, ramosus, tuberculis parvis subasper, leniter angulosus, oxygalis et ultra.

RAMI alterni, juniores purpurei.

FOLIA petiolata, mollia, venosa, in caulem subdecurentia, inferiora ovato-lanceolata, integerrima; superiora trilobohastata.

FLORES in CYMAS racemosas dispositi; pedunculi florales ad basim bulbosi, aut ex acetabulo quasi prodeuntes.

CALYX. PERIANTHIUM monophyllum, parvum, quinquefidum, purpureum, segmentis obtusiusculis, persistentibus; *fig. 1.*

COROLLA monopetalâ, rotata; TUBUS brevissimus; LIMBUS quinquepartitus, LACINIIS lanceolatis, purpureis, reflexis; FAUX nigra, nitida, ad basin singulâe lacinia maculâe duæ, virides, *fig. 3—2.*

STAMINA. FILAMENTA quinque, brevissima, tubo corollæ inserta, nigro-purpurea. ANTHERÆ quinque, flavæ, erectæ, in tubum subconicum coalescentes, apicibus biforaminosis, *fig. 4—5.*

PISTILLUM. GERMEN pyriforme; STYLUS subulatus, staminibus paulo longior; STIGMA simplex, obtusum, *fig. 6.*

PERICARPIMUM. BACCA ovata, coccinea, glabra, bilocularis, receptaculo utrinque convexo, cui semina adnæctuntur, *fig. 8.*

SEMINA plures, lutescentia, compressa, subreniformia, pulpo odoris ingrati oblecta, *fig. 9.*

ROOT perennial.

STALK woody, climbing, hollow, branched, thinly beset with small pointed tubercles, slightly angular, and growing to the height of six feet, or more.

BRANCHES alternate, the younger ones purple,

LEAVES standing on foot-stalks, of an oval pointed shape, soft, veiny, running slightly down the stalk, the lower ones entire, the upper ones halbert shaped.

FLOWERS growing in branched CYMÆ, the proper peduncles of the flowers bulbous at their base, or growing out of a kind of socket.

CALYX a PERIANTHIUM of one leaf, small, and purple, divided into five segments, the segments bluntish, persisting; *fig. 1.*

COROLLA monopetalous, wheel-shaped, the TUBE very short, the limb divided into five segments, the SEGMENTS lancet-shaped, purple, and turning back; the MOUTH black and shining, at the bottom of each segment are two roundish green spots, *fig. 3. 2.*

STAMINA. five FILAMENTS very short, of a black purple colour, and inserted into the tube of the Corolla. Five ANTHERÆ yellow, upright, and uniting into a tube, with two holes at the top of each, out of which the POLLEN is discharged, *fig. 4. 5.*

PISTILLUM. the GERMEN pear-shaped; the STYLE tapering, a little longer than the Stamina; the STIGMA simple and obtuse; *fig. 6.*

PERICARPIMUM. an oval, scarlet, smooth BERRY of two cavities, the receptacle to which the seeds are connected, is round, on both sides; *fig. 8.*

SEEDS several, flat, somewhat kidney-shaped; *fig. 9.* of a yellowish colour, inclosed in the pulp, which has a disagreeable smell.

THE Woody Nightshade has been commended as a medicine for many distempers by the old Botanists, in their usually lavish manner; but PARKINSON says, he found the juice of it prove a very churlish purge. LINNÆUS prefers an infusion of the stalk of this plant to any of the foreign woods, as a cleanser of the blood, and recommends it in inflammatory fevers, obstructions, the itch, and rheumatism; and to render the knowledge of plants as extensively useful as possible, he does not think it beneath him to remark, that the Swedish Peasants make hoops of the stalk of this plant to bind their wooden cans. RAY informs us, that the inhabitants of *Wegysphalia*, who are subject to the scurvy, make use of a decoction of the whole plant as their common drink with success against that distemper.

FLOYER says, that thirty berries of this plant killed a dog in less than three hours, and remained undigested in his stomach; as these berries from their resemblance, may happen by mistake to be eaten for currants by children, it may not be improper to remark, that in such a case, it is advisable to pour down instantly as much warm water as possible, to dilute the poisonous juice, and provoke vomiting, till farther assistance can be had.

Goats and sheep are said to feed on this plant, but our other cattle, *vis.* kine, horses, and swine, refuse it.

It grows plentifully in moist hedges, and blows from July to August. The berries are ripe in September and October. It is sometimes found with a white flower.

1871

1871

1871

1871

1871

1871

1871

1871

1871

1871

1871

1871

LIBRARY OF THE UNIVERSITY OF TORONTO

Proc. J. Inst. Chem. Ind. 1961, 1, 101

LONICERA PERICLYMENUM. HONEY SUCKLE or WOODBINE.

LONICERA *Linnaei* Gen. Pl. PENTANDRIA MONOGYNIA.

Raii Synops. ARBORES ET FRUTICES FRUCTU FLORI PFTALOIDI CONTIGUO.

LONICERA capitulis ovatis imbricatis terminalibus, foliis omnibus distinctis. *Lin. Sp. Pl.* 247.

PERICLYMENUS *Fuschii* Icon 646.

PERICLYMENUM non perfoliatum Germanicum. *Bauhin pin.* 302.

CAPRIFOLIUM Germanicum *Dodon. Gerard. emac.* 891. *Parkinson.* 1460. *Raii Synop.* 458. *Hudson Fl.* 80.

Haller. hist. 301. *Scopoli Fl. Carniol.* p. 153.

CAULIS lignosus, volubilis, orgyalis et ultra; cortice pallide fusco; RAMI oppositi, purpurei.

FOLIA opposita, ovata, glabra, subtus cærulescentia.

FLORES terminales, verticillatim dispositi, patentes, rubri, interne flavi, odoratissimi.

CALYX PERIANTHIUM superum, brevissimum, quinquepartitum; segmentis ovato-lanceolatis, erectis, duobus inferioribus remotioribus. *fig. 1.*

BRACTEÆ subcordatæ, *fig. 8.* germina imbricatum cingentes, ad marginem præcipue scabræ, ut sunt calyx, et tubi basis pilis glanduliferis.

COROLLA monopetala, tubulosa, tubus oblongus, subinfundibuliformis, LIMBUS bipartitus, lacinis revolutis, superiore quadrifida, segmentis fere aequalibus, obtusis, inferiore integra. *fig. 2.*

STAMINA: FILAMENTA quinque filiformia, corolla longiora, alba, tubo corollæ inserta; *fig. 3.* ANTHERÆ dum pollinem involvunt oblongæ, incumbentes, postea lunatæ. *fig. 4.*

PISTILLUM: GERMEN subrotundum, inferum, *fig. 5.* STYLUS filiformis, Staminibus paulo longior, *fig. 6.* STIGMA capitatum, subrotundum, trifidum, viride. *fig. 7.*

PERICARPIA: BACCÆ plures, subrotundæ, rubræ, umbilicatæ, biloculares, omnes distinctæ. *fig. 9.*

SEMINA Plura, lutescentia, hinc convexa inde plana. *fig. 10.*

STALK woody, twining, growing to the height of six feet or more, the Bark a pale brown, the BRANCHES opposite and purple.

LEAVES opposite, oval, smooth, underneath of a blueish colour.

FLOWERS terminal, growing in a whirl, and spreading; externally red, internally yellow, and fragrant.

CALYX a Perianthium placed above the Germen, very short, divided into five segments, which are of an oval pointed shape, and upright, the two inferior ones most remote from each other. *fig. 1.*

FLORAL LEAVES laying one over the other, and closely embracing the Germina, reddish at the edges and covered as well as the Calyx and base of the tube with glandular hairs. *fig. 8.*

COROLLA monopetalous, and tubular, the TUBE long, and somewhat funnel shaped, the LIMB bipartite, the lacinie rolling back, the upper one divided into four blunt and nearly equal segments, the lower one entire. *fig. 2.*

STAMINA: five white FILAMENTS, of an equal thickness throughout, longer than the Corolla and inserted into its tube, *fig. 3.* the ANTHERÆ while they contain the Pollen oblong, afterwards femilunar and of a yellow colour. *fig. 4.*

PISTILLUM: the GERMEN roundish and placed below the Calyx, *fig. 5.* the STYLE filiform, a little longer than the Stamina, *fig. 6.* the STIGMA roundish, trifid, and of a green colour. *fig. 7.*

SEED VESSELS several roundish red BERRIES, having the remains of the Calyx adhering to them, and all distinct. *fig. 9.*

SEEDS several, of a yellowish brown colour, round on one side and flattish on the other. *fig. 10.*

The early writers attributed virtues to this officinal plant which the latter have been inclined to give up, as a medicine we must not expect much from it, but the beauty, singularity, and exquisite fragrance of its flowers have long given it a place in our Gardens. It is a Climber and turns from East to West with most of our other English Climbers, and in common with them it bears clipping and pruning well; for in a state of nature those plants that cannot ascend without twining round others are often liable to lose large branches, they have therefore a proportional vigour of growth to restore accidental damages. This plant is subject when placed near Buildings to be disfigured and injured by small insects called Aphides or vulgarly blights, these Animalculæ were formerly supposed to be brought by the East Wind, and consequently the mischief was looked upon as inevitable, but observation has of late years corrected that Error, their history is well known*, but no effectual remedy against them is as yet discovered. These Insects are not very numerous in Spring, but as the Summer advances they encrease in a surprising degree; to preserve the plant therefore from injury it is necessary to watch their first attacks, cut off and destroy the branches they first appear on, for when they have once gained ground they are defended by their numbers. We have seen small plants cleared of them by sprinkling Spanish snuff on the infected branches, but for large trees this remedy is scarcely practicable. The leaves are likewise liable to be curled up by a small caterpillar (*Phalæna Tortrix Linnaei*) which produces a beautiful little Moth, see *Albini's history of english Insects* pl. 73. It is fed on by Kine, Goats, and Sheep, but Horses refuse it.

To shew the confusion of ancient names it may not be improper to mention that this plant and Woodrose (*Asperula odorata*) have been both called *Matrifylva* by the old botanic writers. Our Poets also have strangely confounded the names of this plant, *SHAKESPEAR* says

*So doth the WOODBINE the sweet HONEYSUCKLE
Gently entwine*

MILTON seems to call this plant *Eglantine* although that is an undoubted name for the *Sweet Briar*

Through the SWEET BRIAR or the Vine


Or the TWISTED EGLANTINE.

We find it plentifully in Woods and Hedges flowering from July to September, such plants as grow in shady places produce blossoms of a paler colour, and they universally smell sweetest in the Evening; at which time some particular species of Sphinges (*Linnaei*) or Hawk Moths are frequently observed in Gardens hovering over the blossoms, and with their long tongues which are peculiarly adapted to the purpose, extracting honey from the very bottom of the flowers.

* Vid *Reaumur* and *Geoffroy*.




Lonicera Periclymenum.







Hedera Helix.



HEDERA HELIX. IVY.

HEDERA *Limæi* Gen. Pl. PENTANDRIA MONOGYNIA. *Petala* quinque oblonga. *Bacca* quinquesperma calyce cincla.

Raii Syn. ARBORES ET FRUTICES FRUCTU FLORI PETALOIDI CONTIGUO.

HEDERA *Helix* foliis ovatis lobatisque. *Linn. Syst. Vegetab. p. 202. Sp. Pl. 292. Fl. Suecic. p. 75.*

HEDERA *Helix* foliis trilobatis, fructiferis ovato-lobatis. *Holler hist. helv. n. 826.*

HEDERA *Helix. Scopoli Fl. Carniol. n. 271. Hudson Fl. Angl. p. 85.*

HEDERA arborea. *Baubin. Pin. 305.*

HEDERA poetica. *Baubin. Pin. 305.*

HEDERA major sterilis. *Baubin. Pin. 305.*

HEDERA humi repens. *Baubin. Pin. 305.*

HEDERA arborea five scandens et corymbosa communis. *Parkinson 678.*

HEDERA *Helix* Ger. Em. 858. *Raii Syn. 459.* Climbing or Berried Ivy: also Barren or Creeping Ivy.

TRUNCUS in arboribus hujus speciei senescentibus cortice rimoso cinereo vestitur, in novellis ramis viridis aut purpureo cernitur, fibrillas e latere interiori exerit, quorum ope proximis arboribus aut parietibus innixus alta petit.

TRUNK; the trunk in trees of this species, which are old, is covered with an ash-coloured chopped bark, in the young branches it is of a green or purple colour; from the inside of the trunk a great number of small fibres are thrown out, by the assistance of which, it supports itself on the nearest walls and trees, and climbs aloft.

FOLIA quam maxime varia, dum planta repit plerumque trilobata, quinquelobata etiam occurrunt; adnunculis derelictis, ovata sunt; glabra, nitentia, nunc rubedine ornata, nunc venis albis picta, prefertim in ramulis junioribus.

LEAVES as various as possible, while the plant creeps they are in general trilobate, sometimes quinquelobate, leaving its supporters, they become oval; smooth, shining, sometimes figured with red, sometimes painted with white veins, particularly in the young branches.

FLORES lutescentes, in summitatibus caulium umbellatim dispositi, UMBELLÆ densæ, globosæ.

FLOWERS yellowish, growing on the top of the stalks in thick round UMBELS,

COROLLA: quinque, ovata, flavescentia, patentia.

COROLLA: PETALS five, oval, yellowish and spreading.

STAMINA: FILAMENTA quinque longitudine Corollæ; ANTHERÆ basi bifidæ, incumbentes, fig. 1.

STAMINA: five FILAMENTS the length of the Corolla; ANTHERÆ bifid at bottom, and incumbent, fig. 1.

PISTILLUM: GERMEN turbinatum; STYLUS simplex, brevissimus; STIGMA simplex, fig. 2.

PISTILLUM: GERMEN roundish; STYLE simple and very short; STIGMA simple, fig. 2.

PERICARPIUM: BACCA globosa, nigra, intus purpurea, quadrilocularis aut quinquelocularis, coronata receptaculo et stylo conico brevi, loculis monospermis, fig. 3, 4.

SEED-VESSEL: a round BERRY, externally black, internally purple, with four or five cavities each containing one seed, crowned with the receptacle and short conic Style, fig. 3, 4.

SEMINA quinque, hinc gibba, inde angulata, fig. 6.

SEEDS five, on one side gibbous, on the other angular, fig. 6.

The *Hedera Helix* begins to blow in sunny aspects towards the end of September, and according to situation blows on through October, and November. This plant is one of the last blowers, and is much resorted to by bees, and flies of various species, which swarm on its branches, and feed on its blossoms, making such a humming on sunny days as may be distinguished at a considerable distance.

The berries encrease in bulk gradually all through the winter months, and are full formed by February; in April they ripen and turn very black, and are eaten by several species of thrushes, and wild pigeons. Thus does fructification manifestly obtain in this instance all through the winter months, as well as in the mosses and lichens.

Sheep are very fond of Ivy, which in hard weather is a warm and wholesome food; and therefore shepherds in snowy seasons cut down branches for their flocks to browse on. CATO directs that in a scarcity of hay, cattle should be foddered with Ivy.


Professor KALM, in his travels through the greatest part of N. America, saw but one plant of Ivy, and that was running up the walls of a man's house: this specimen was probably carried thither by some European who perhaps was desirous of propagating in that new world a plant that might still recall to his mind the pleasing Idea of his native cottage, tufted with the foliage of this beautiful Evergreen.

The ancients held this plant in great esteem; their Heroes and Poets are described as wearing garlands composed of it. The supposition of its preventing intoxication is of very early date: Homer therefore mentions his Bacchus as Ivy-crowned and often describes his Heroes drinking out of a Cup made of the wood of Ivy. (κεκυβηται.) CATO tells us that with a cup of this kind we may distinguish wine that has been adulterated with water, for the wine will be discharged and the water remain: to such an extravagant assertion has this grave author been probably led by relying on the supposed antipathy between the wine and ivy: This cup is still used in some parts of the kingdom as a remedy for a trembling hand; but rational practice has not admitted any part of the *Hedera* into the *Materia Medica*, Ivy-leaves however are said to be successfully applied to painful Corns. When it trails on the ground it grows branches are small and weak; and its leaves are divided into three lobes; but when it climbs walls or trees it grows much stronger and the leaf changes to an oval form: these different appearances induced old Botanists to suppose there were two or three different species. In its variegated state it sometimes appears almost white, and may perhaps be the *Hedera alba*, and *pallentes Hederae* of VIRGIL.

Few people are acquainted with the beauty of Ivy when suffered to run up a stake, and at length to form itself into a standard, the singular complication of its branches, and the vivid hue of its leaves give it one of the first places among evergreens in a shrubbery; In woods when suffered to grow large and rampant, this plant by twining round their bodies does great damage to timber trees; and therefore should be carefully destroyed: but in ornamented Out-lets, where evergreens do not abound, a few trees covered with Ivy have a very pleasing effect, and moreover induce birds to build their Nests particularly the Black-bird.

In the Stump of Ivy many birds build their Nests particularly the Black-bird. When Ivy is prejudicial, it may easily be destroyed, tho' it has spread to a great height, by cutting through its Trunk, and this shows that the fibres which the Stalk throws out in so singular a manner serve more to support than nourish it.

The soft wood of Ivy is made use of by Shoemakers to give a smooth edge to their cutting knives.





(*onium maculatum*).

CONIUM MACULATUM. HEMLOCK.

CONIUM *Linnaei* Gen. Pl. PENTANDRIA DIGYNIA.

Raii. Syn. Gen. II. UMBELLIFERÆ HERBÆ.

CONIUM *maculatum* feminibus striatis. *Linn. Syst. Vegetab.* p. 229.

CICUTA *Haller. bist. helv. n. 766. v. 1. p. 337.*

CONIUM *maculatum*. *Scoepoli Fl. Carniol.* p. 207.

CICUTA *major Bauhin. Pin.* 166.

CICUTA *Gerard emac.* 1061.

CICUTA *vulgaris major Parkinson 933. Raii Syn. p. 215. Hudson Fl. Angl. p. 100. Störck. Cicut. Suppl. p. 7.*

t. 1.

RADIX biennis, crassitudine digiti, longa usque ad pedalem, in crura sæpe divisa, juniori Pastinacæ haud dissimilis, odoris gravis, et saporis subdulcis: secundo anno in caulescente planta succo fere caret, firma solidiorque evadit.

CAULIS orgyalis, teres, nitidus, lævis, fistulosus, ad basin crassitie pollicis, rore glauco tectus, et maculis fanguloneis pictus, versus summum ramosus, et striatus.

FOLIA inferiora magna, etiam bipedalia, atro-virentia, nitentia, multiplicato-pinnata, pinnulis oblongis inciso-serratis; SPATHA fulcata.

INFLORESCENTIA. *Umbella universalis* Radiis plurimis patentibus striatis; *partialis* confimilis.

CALYX: *Involucrum universale* e foliolis 5-7 constat, lanceolato acuminatis, reflexis, margine albidis, *fig. 1*; *partiale* 3 aut 4 dimidiatis, extrorsum patentibus, *fig. 2.*

COROLLA: PETALA quinque, alba, inæqualia, inflexo-cordata, *fig. 3.*

STAMINA: FILAMENTA quinque, alba, longitudine Corollæ; ANTHERÆ albæ, *fig. 3.*

PISTILLUM: STYLI duo, albi, filiformes, non admodum breves; STIGMATA subrotunda; GERMIN inferum, striatum, *fig. 3. 4.*

FRUCTUS subrotundus, e binis feminibus fuscescentibus compositus; hinc planiusculus, illic gibbus, cum striis quinque elevatis crenulatis, *fig. 4. 5.*

ROOT biennial, the thickness of ones finger, from six inches to a foot in length, frequently forked, and not unlike that of a young Paríep, of a disagreeable smell and sweetish taste: in the second year of its growth when the plant has a flowering stem, it becomes drier, more firm and solid.

STALK about six inches high, round, shining, smooth and hollow, at bottom the thickness of ones thumb, covered with a blueish kind of powder which easily wipes off, and spotted with red, towards the top branched and striated,

LEAVES. The bottom leaves large, even two feet long, of a dark green colour and shining, many times pinnated, the pinnulæ oblong and sharply cut in; the SPATHA grooved.

INFLORESCENCE. The *Universal Umbell* is composed of many striated and spreading Radii; the *Partial Umbell* similar to it.

CALYX: the *Universal Involucrum* consists of 5 or 7 leaves, which are lanceolate, turned back, and whitish at the edges, *fig. 1*: the *Partial Involucrum* is composed of 3 or 4 leaves, which surround one half of the stalk only, and spread outward, *fig. 2.*

COROLLA: PETALS five, white, unequal, heart-shaped, and bent in at top, *fig. 3.*

STAMINA: FILAMENTS five, white, the length of the Corolla; ANTHERÆ white, *fig. 3.*

PISTILLUM: GERMIN beneath the Corolla, striated, *fig. 3. 4*; STYLES two, filiform and not very short; STIGMATA round, *fig. 3.*

FRUIT is roundish, and composed of two brownish seeds, flattish on one side and round on the other, with five notched and elevated ridges, *fig. 4. 5.*

THE powerfull deleterious properties of this herb have been long known and acknowledged by all botanic writers; whence it has been commonly ranged in the class of Vegetable Poisons. And as such active principles under skilful management, are likely to accord the most efficacious remedies, this plant has been also admitted as an article of the *Materia Medica*. Until lately however, the use of it was chiefly confined to external applications, where its narcotic qualities may undoubtedly assist in assuaging pain, staving off supuration, &c. But in the year 1760, Dr. Störck, a famous Practitioner at Vienna, published a treatise on Hemlock, recommending an extract made of the inspissated juice of the herb to be taken internally, from four grains to sixty, or upwards, every day, as a cure for the *Schrophula*, *Cancer*, and others of the most terrible and inveterate disorders incident to the human body.

Our Physicians though laudably cautious of admitting or trusting to novelties, received Dr. Störck's publication with uncommon ardour, and perhaps no new medicine was ever more immediately or generally tried than this *Extractum Cicute*. The success however not answering their expectation, led some to think they had mistaken the plant. The Author was applied to, and this produced a supplement (printed 1764) wherein the species is figured, and clearly shewn to be the *Conium maculatum* of LINNÆUS. It were to be wished this had cleared up all difficulties. In his first treatise the Doctor tells us that the fresh root sliced, yielded a bitter acrid milk, of which a single drop or two being applied to the tip of his tongue, presently rendered it painful, rigid, and so much swelled that he could not speak. Yet it is certain that the roots of our *Hemlock* may be chewed and swallowed in considerable quantities without producing any sensible effect. Mr. ALCHORNE (who I believe was the first that laudably exerted himself in investigating this matter,) assures me that he has tried this in every season of the Year, and in most parts of our Island, without finding any material difference: and that

he

he has also been well informed both from *Berlin* and *Vienna*, that the *Hemlock* Roots in those countries, are no more virulent than ours about *London*. Mr. TIMOTHY LANE informs me, that he also with great caution made some experiments of the like kind, and in a short time found he could venture to eat a considerable part of a root without any inconvenience; after that, he had some large roots boiled, and found them as agreeable eating at dinner with meat, as Carrots, which they in taste somewhat resembled: and as far as his experience, joined with that of others informed him, the Roots might be cultivated in Gardens, and either eaten raw like Celery, or boiled as Parsneps or Carrots. That in Spring and Winter they are not woody as in Summer: that he has eaten them from different places and in all seasons; and that he perceived some roots were more pungent than others, but not in any degree worthy notice.

The experiments of these ingenious Gentlemen sufficiently evince the innocence of the roots of this plant, contrary to what has been asserted by Dr. STÖACK, and hence we may infer that whatever accounts have been related by Authors of their poisonous qualities, the Roots of some other Plant must have been made use of. In the poisonous quality of the *Herb* however all Authors seem agreed, but with respect to its efficacy as a medicine they very much differ. If we may believe Dr. STÖACK, there is scarce a disease incident to the human body which it either does not cure, or relieve; but it is remarkable that a copious experience of fifteen years, as well in the great Hospitals of this Metropolis as in the private practice of the whole Kingdom, should not have afforded one instance of a perfect cure by the Extract, at least none such has appeared among the valuable collections of cases published by our College of Physicians and other Medical Societies. Both Dr. FOTHERGILL of London, and the late Dr. RUTTY of Ireland, men of the greatest eminence in their profession, have declared that the success attending it has not been equal to what they had reason to expect from Dr. STÖACK's account of it; (*vid. medical observations and enquiries, vol. 3.*) yet tho' it had failed them in the cure of many of those diseases which unfortunately were the *opprobria medicorum*, it had proved beneficial in various obstinate complaints; Scrophulous tumours were to appearance dissolved by it; the progress both of occult and ulcerated Cancers was retarded, the pain alleviated, and the discharge changed for the better in every respect; divers putrid and fœtid Ulcers were by the use of *Hemlock* remarkably mended in their discharge, and disposed to heal, in some of which the Sublimate had been given in vain; hence the Extract is still frequently used, and will probably continue to be prescribed, because its effects as an Anodyne will often afford at least a temporary relief, and because in desperate diseases a doubtful remedy seems better than none at all.

The taking of the Extract is generally attended with a giddiness and often with a pain of the head, nausea, and other disagreeable symptoms; in some however its effects are apparently anodyne, as it eases pain and promotes rest even where Opium has failed.

Physicians seem somewhat divided about the best mode of exhibiting this medicine, some recommending the extract as being most easily taken in the form of pills, others the powder, as not being subject to that variation which the extract is liable to from being made in different ways. With respect to the period likewise at which the plant should be gathered, they seem not perfectly agreed, some recommending it when in its full vigour, and just coming into bloom, others when the flowers are going off and the whole plant has acquired a yellowish hue. That the Extract might be at all times equally active, and uniformly prepared, Dr. CULLEN has for many years recommended the making it from the unripe seeds, and this mode the College of Physicians at Edinburgh has thought proper to adopt in their new Pharmacopœia.

Hemlock grows very frequently on banks by the sides of Roads, by hedge sides, and in Fields and Gardens, flowering in the month of July.


We have a common English Proverb that *what is one Mans Meat is another mans Poison*, and agreeable to this are the lines of LUCRETIVS which relate to this plant;

“*Pinguetere sæpe Cicuta
Barbigeros pecudes homini quæ est acro venenum.*”

That it affords nourishment to Birds likewise there is sufficient evidence, our learned Philosopher and accurate Naturalist Mr. RAY, found in the Crop of a Thrush abundance of *Hemlock* seeds, at a time too when other vegetable food might be had in abundance. It appears to be eaten by very few or no Insects.

The dried stems or hexes are used by Boys for various purposes.

The *Hemlock* is obviously distinguished from our other umbelliferous plants by its large and spotted stalk, by the dark and shining green colour of its bottom leaves, and particularly by their disagreeable smell when bruised, and which according to Dr. STÖACK resembles that of Mice. The *Fools Parsley* and *Scandiv* with rough seeds are the most likely to be mistaken for this poisonous plant, but may easily be distinguished if attention be paid to the descriptions and figures we have already given of them,





Euthusa cynapium.

ÆTHUSA CYNAPIUM. FOOL'S PARSLEY.

ÆTHUSA *Linnei Gen. Pl. PENTANDRIA DIGYNIA.*

Rati Syn. Gen. 17. UMBELLIFERÆ HERBÆ.

ÆTHUSA (*Cynapium*) foliis conformibus. *Linnei Syst. Vegetab. p. 236. Flor. Suecic. p. 92.*

ÆTHUSA. *Haller. hist. n. 765.*

CICUTA minor petioellino similis. *Baubin. Pin. p. 160.*

CICUTARIA Apii folio. *J. Baubin.*

CICUTARIA tenuifolia *Gerard. emac. 1063.*

CICUTA minor five fatua *Parkinson. 933. Rati Syn. p. 215. the lesser Hemlock or Fool's Parsley. Scopoli Fl. Carniol. p. 206. Hudson Fl. Angl. p. 107. Hill's British Herbal small Hemlock tab. 58. icon pessima.*

RADIX annua, fusiformis, alba, minimi digiti crassitudine, paucis fibris instructa.

CAULIS pedalis ad bipedalem, erectus, ramulos, striatus, fistulosus, glaucus, versus basin sæpe purpureus, non vero maculatus.

FOLIA radicalia et ramea conformia, lævia, superne atro-virentia, inferne pallidiora, nitentia, duplicato-pinnata, pinnis pinnatifidis, profunde incisis, pinnulis ovato-acutis, mucronatis. Vagina ad basin petiolorum parva, læves, marginibus membranaceis.

PETIOLI erecti, sulcati.

UMBELLA uniuersalis patens, radiis interioribus per gradus brevioribus, intimis brevissimis; partialis uniuersali similis.

INVOLUCRUM uniuersale nullum, partiale dimidiatum, extus positum, foliolis tribus longissimis linearibus pendulis, fig. 1.

COROLLA: PETALA quinque, alba, obcordata, inæqualia, apice inflexa, exteriora maiora, fig. 2.

STAMINA: FILAMENTA quinque, alba, longitudine corollæ, inflexa: ANTHERÆ albæ, nonnunquam rubellæ, fig. 3.

PISTILLUM: GERMEN inferum, glandulâ virescente coronatum: STYLIS duo, primum erecti, dein deflexi: STIGMATA obtusa, fig. 4.

PERICARPIUM nullum: FRUCTUS ovato-subrotundus, striatus, bipartibilis, fig. 5.

SEMINA duo, pallide fusca, hinc convexa, profunde striata, hinc plana, figurâ ovato-acutâ notata, fig. 6.

ROOT annual, tapering, of a white colour, about the thickness of the little finger, furnished with few fibres.

STALK from one to two feet high, upright, branched, striated or slightly grooved, hollow, covered with a blueish kind of powder which easily wipes off, towards the bottom frequently of a purple colour, but not spotted.

LEAVES: the bottom leaves and those of the branches similar, smooth, on the upper side of a dark green colour, underneath paler and shining, twice pinnated, the leaves pinnatifid and deeply cut in, the small leaves or pinnule oval and terminating in a fine point. The SHEATH at the base of the foot-stalks small, smooth and membranous at the edges.

FOOT-STALKS of the flowers, upright and grooved. UMBEL: the uniuersal umbel spreading, the inner radii gradually shorter, the innermost very short; the partial umbel like the uniuersal.

INVOLUCRUM: the uniuersal INVOLUCRUM wanting, the partial one placed externally, and only surrounding one half of the umbel, composed of three very long, linear, and pendulous leaves, fig. 1.

COROLLA: five unequal, heart-shaped, white PETALS, bent in at top, the outer ones largest, fig. 2.

STAMINA: five white FILAMENTS the length of the Corolla, bending in: ANTHERÆ white, sometimes reddish, fig. 3.

PISTILLUM: GERMEN placed below the corolla, and crowned by a glandular substance of a greenish colour: two STYLES first upright, afterwards bending downward: STIGMATA blunt, fig. 4.

SEED-VESSEL wanting: the FRUIT or unripe seed of an oval roundish shape, striated, and dividing into two parts, fig. 5.

SEEDS two, of a pale brown colour, convex and deeply striated on one side, flat on the other, and marked with a figure of an oval pointed shape, fig. 6.

ONE of the principal advantages resulting to mankind from Botany, is the rightly ascertaining those plants which are used for food, from those which are known to be poisonous. It not unfrequently happens that both these kinds of Herbs grow in the same soil, may often in the same bed together, and so similar are they in their general appearance, that the indiscriminating eye of the common observer readily mistakes the one for the other, and hence diseases fatal in their consequences sometimes ensue. To point out then the most obvious distinctions between such kinds of plants, is not only our business but our duty.

The Fool's Parsley seems generally allowed to be a plant which possesses poisonous qualities.


Baron HALLER has taken a great deal of pains to collect what has been said concerning it, and quotes many authorities to shew that this plant (on being eaten) has been productive of the most violent symptoms, such as anxiety, hiccough, and a delirium even for the space of three months, stupor, vomiting, convulsions and death: He suspects however that the common Hemlock may sometimes have had a share in producing these symptoms, as he finds in authors that the Fool's Parsley had been used by a whole family without any bad effect, although he imagines this might be owing to the smallness of the quantity eaten. As a corroborating proof of its deleterious quality, LINNÆUS asserts that it proves fatal to geese if they happen to eat it.

Altho' it seems rather doubtful whether it be so poisonous to mankind as is represented, yet it will perhaps be most prudent to consider it as such, until future experiments shall determine its effects with more certainty.

The plants to which this bears the greatest resemblance, are common Garden Parsley and common Hemlock, *Conium maculatum*; this similarity has been observed by most Botanic Writers, some of whom have called it a kind of Hemlock, others a kind of Parsley; it differs however considerably from both these Genera. The colour of its leaves alone, is nearly sufficient to distinguish it from Parsley; those of common Parsley are of a yellowish green colour, those of Fool's Parsley of a very dark green, and much more finely divided; the leaves of Parsley when bruised have a strong but not disagreeable smell, those of Fool's Parsley have very little smell in them. These marks if attended to are sufficient to distinguish the leaves of these two plants, and in the state of leaves they are most liable to be taken for one another, as they grow together in Gardens. Where much Parsley is used, the Mistress of the house therefore would do well to examine the Herbs previous to their being made use of; but the best precaution will be always to sow that variety called curled Parsley, which cannot be mistaken for this or any other plant.

It is distinguished from Hemlock by being in every respect smaller, and not having that strong disagreeable smell which characterizes the leaves of that plant; the stalk likewise is not spotted as in the Hemlock; and lastly it is distinguished from all our umbelliferous plants by the three long, narrow, pendulous leaves which compose its partial involucre, and which are placed at the bottom of each of the small Umbels.

It grows very common in Gardens, and all kinds of cultivated ground, and flowers in July and August.





Scandix tuberosa

SCANDIX ANTHRISCUS. SCANDIX WITH ROUGH SEEDS.

SCANDIX *Linnaei Gen. Pl. PENTANDRIA DIGYNIA*

Raii Syn. Gen. II. UMBELLIFERÆ HERBÆ

SCANDIX *Anthriscus* feminibus ovatis hispida, corollis uniformibus, caule levī. *Linnaei Syst. Vegetab. p. 237. Flor. Suecic. p. 93.*

CAUCALIS vaginis lanuginosis, foliis triplicato-pinnatis, feminibus rostratis. *Haller hist. n. 743.*


MYRRHIS sylvestris, feminibus asperis. *Babin pin. 160. Parkinson 935. Ger. emac. 1038. Raii Syn. p. 220. Small Hemlock-Chervil with rough Seeds. Hudson Fl. Angl. p. 108. Jacquin Flor. Austriac. Vol. 2. p. 35. tab. 154.*

RADIX annua, parva, albida, subinspida.	ROOT annual, small, whitish, with little taste.
CAULIS pedalis ad tripedalem, sæpe altior, tuberculosus, teres, fistulosus, lævis, ad genicula tumidus et substriatus, plerumque viridis.	STALK from one to three feet high, frequently taller, nearly upright, round, hollow, smooth, swelled and slightly striated at the joints, and most commonly green.
FOLIA. Vaginæ ad basin foliorum magnæ, marginibus lanuginosis; Folia mollia, tenera, multiplicate pinnata, hirsutula, ex luteo-virentia.	LEAVES. The sheaths formed by the base of the leaves are large and downy at the edges: the leaves soft, tender, many times pinnated, slightly hairy, and of a yellowish green colour.
INFLORESCENTIA Umbella. UMBELLÆ obliquæ, pedunculatæ: Pedunculus universalis RADII brevior, RADII univesales 3—5. glabri, partiales 2—6.	INFLORESCENCE an Umbell, the UMBELLS oblique, standing on footstalks, the general or universal footstalk shorter than the RADII; the universal RADII from 3 to 5, the partial RADII from 2 to 6.
CALYX: Involucrum universale nullum. Partiale plerumque pentaphyllum, foliolis lanceolato-acuminatis, ciliatis, persistentibus fig. 1.	CALYX. The universal Involucrum wanting, the Partial one generally composed of five leaves, which are pointed, hairy at the edges, and continue. fig. 1.
COROLLA: PETALA quinque, minima, subæqualia, alba, subcordata, apicibus inflexis. fig. 2.	COROLLA: five PETALS very minute, nearly equal, white, somewhat heart shaped, the tips bending in, fig. 2.
STAMINA: FILAMENTA quinque, petalis paulo breviora; ANTHERÆ primum virides, dein fuscæ fig. 3.	STAMINA: five FILAMENTS, a little shorter than the Petals; the ANTHERÆ first green, afterwards brown, fig. 3.
PISTILLUM: GERME oblongum, inferum, subcompressum, hirsutum, STYLI duo breves. fig. 5.	PISTILLUM: the GERME oblong, placed beneath the Corolla, flattish, and rough, two STYLES very short fig. 5.
SEMINA duo, oblonga, e fusco-nigricantia, hinc fuscato-plana, inde convexa, rostrata, pilis rigidis hamatis undique aspera fig. 6.	SEEDS two, oblong, of a dark brown colour, on one side flat and grooved, on the other convex, running out to a point, and prickly with stiff hooked hairs, fig. 6.

THE great similarity in the external appearance of a great number of umbelliferous plants, frequently hath been the cause of mistakes, which have sometimes proved hurtful to the health of individuals. At the same time that there is no class of plants which, at first sight, appears to the young Botanist more difficult of investigation than this, there is none perhaps which affords more constant, or more certain marks of generic and specific difference. Obvious distinctions may be drawn from the *Stalk* and *Leaves*: in some the stalk is smooth in others, rough; and in others, more or less deeply channelled; in some the leaves are very finely divided; and in others, but coarsely so; but the parts of *Frustrification* afford the most pleasing, and scientific, distinguishing marks. The absence, or presence of the general and partial *Involucrum* the number, shape, and situation of its leaves, the number of the *Radii* which compose the umbell, the size and equality of the *Petals*, and the very different appearances of the *Seeds*, all unite to render a knowledge of these plants easily acquired.

Some of the *Umbelliferi* are used in food, and others in medicine; the greatest care will therefore be necessary in the drawing and description of these; and in this, no one seems to have succeeded so well as the celebrated JACQUIN. In the first and second volumes of his *Flora Austriaca*, lately published, and which indeed are a most valuable addition to the stock of botanic knowledge, a great number of these plants are figured and described.

This plant grows very common on dry banks and in hedges, flowers from the beginning to the end of May, and the seeds are ripe in June. When it becomes luxuriant, as it sometimes will from growing in a moist situation, it puts on somewhat the appearance of the common Hemlock, but may easily be distinguished from that poisonous plant, if attention be paid to the following particulars: The leaves of the Hemlock are perfectly smooth; these have a flight hairiness, are more finely divided, and of a paler green; the stalk of the Hemlock is spotted; this is not; the Hemlock has a general involucrium, which in this plant is wanting; the seeds of the Hemlock are smooth, and these are rough; the Hemlock has a strong disagreeable smell; this not disagreeable, but more like Chervil, to which in its virtues it should seem nearest allied.





Have you

ALSINE MEDIA. COMMON CHICKWEED.

ALSINE *Linnaei Gen. Pl.* PENTANDRIA TRIGYNIA.

Cal. 5-phyllus. *Petala* 5-aequalia. *Caps.* 1-locularis, 3-valvis.

Raii Syn. Gen. 24. HERBÆ PENTAPETALÆ VASCULIFERÆ.

ALSINE *media. Linnaei Syst. Vegetab. p.* 246. *Flora Suecic. p.* 37.

ALSINE *foliis petiolatis, ovato lanceolatis, petalis bipartitis. Haller hift. herb. n.* 880.

ALSINE *media. Scopoli Fl. Carniol. n.* 376.

ALSINE *media. Baubin pin. p.* 250.

ALSINE *media seu minor. Gerard emac. 611. Raii Syn. p.* 347. Common Chickweed. *Hudson Fl. Angl. p.* 113. *Oeder Fl. Dan. 525, 438.*

RADIX annua, fibrosa, capillacea.

CAULES plures, tenelli, teretes, subrepentes, ramosi, viticulis geniculati, *unifariam hirsuti*, apicibus sensim incrassatis.

FOLIA ovato-acuta, glabra, leviter ciliata; inferiora petiolata, superiora sessilia, connata.

PETIOLI ad basin latiora, hirsuti.

PEDUNCULI uniflori, axillares, hirsuti, peractâ florescentiâ penduli, demum erecti.

CALYX: PERIANTHIUM pentaphyllum, foliolis lanceolatis, concavis, subcarinatis, marginatis, hirsutis, *Petalis longioribus, fig. 1.*

COROLLA: PETALA quinque, alba, nitida, ad basin fere partita, *fig. 3, 4, 5.*

STAMINA: FILAMENTA quinque, alba, inter Petala locata, Glandulâ ad basin instructa; ANTHERÆ subrotundæ, purpurascetes, *fig. 5, 6.*

PISTILLUM: GERMEN subovatum; STYLI tres filiformes; STIGMATA simplicia, *fig. 7.*

PERICARPIUM: CAPSULA unilocularis, in valvulas sex dehiscetes, *fig. 8.*

SEMINA octo ad quindecim, subreniformia, aspera, e fusco-aurantiaca, pedicellis receptaculo connexa, *fig. 9, 10, auct.*

ROOT annual, fibrous, capillary.

STALKS numerous, tender, round, striking root here and there, branched, jointed and stringy, *hairy on one side only*, growing thicker towards the top.

LEAVES of a pointed oval shape, smooth, slightly hairy at the edges, the lowermost standing on foot-stalks, the uppermost sessile, connate.

FOOT-STALKS of the leaves broadest at bottom, and hairy.

FOOT-STALKS of the flowers, each sustaining one flower, proceeding from the bosoms of the leaves, hairy, when the flowering is over hanging down, finally becoming upright.

CALYX: a PERIANTHIUM of five leaves, each of which is lanceolate, concave, slightly keel-shaped at bottom, with a margin at the edge, hairy, and longer than the Petals, *fig. 1.*

COROLLA consists of five white shining PETALS, divided nearly to the base, *fig. 3, 4, 5.*

STAMINA: five white FILAMENTS, placed betwixt the Petals, furnished at bottom with a little Gland; ANTHERÆ roundish, of a purplish colour, *fig. 5, 6.*

PISTILLUM: GERMEN somewhat oval; STYLES three, filiform; STIGMATA simple, *fig. 7.*

SEED-VESSEL a CAPSULE of one cavity, splitting into six valves, *fig. 8.*

SEEDS from eight to fifteen, somewhat kidney-shaped, of a brownish orange colour, with a rough surface, connected to the receptacle by little foot-stalks, *fig. 9, 10, magnified.*

CHICKWEED being a plant which will grow in almost any situation, is consequently liable to assume many different appearances: when it grows in a rich soil, and shady situation, it will frequently become so large as to resemble the *Cerastium aquaticum*; while at other times, on a dry barren wall, its leaves and stalks will be so minute, as to make the young botanist take it for some species different from the common Chickweed: happily however it affords marks which if attended to, will readily distinguish it from the *Cerastium*, and every other plant: exclusive of its differing from the *Cerastium* in its generic character, its Petals are shorter than the leaves of its Calyx; while in the *Cerastium* they are longer; hence a considerable difference will be observable at first sight in the size of the flowers of these two plants: and from all other plants related to it, it may be distinguished by the singular appearance of its stalk, which is *alternately hairy on one side only*.

The most common number of its *Stamina* with us is five; yet I have often seen it with less, and sometimes with more; and this inconstancy in the number of its *Stamina* has been noticed by most botanic writers: GOUAN, in his *Flor. Monspel.* mentions from 3 to 10, with as many Pistilla; this circumstance with respect to the number of its *Stamina*, unfortunately separates it from other plants with which it appears to have by nature a very near relation: but as five *Stamina* appear to be its most constant number, LINNÆUS could not have placed it amongst those plants with ten *Stamina*, without doing violence to his system.

Of annual plants there are few more troublesome: it sows itself plentifully in the summer, and remains green throughout the winter, flowering during the whole time, if the weather be mild: but its chief season for flowering is in the spring. In rich garden mould, where the ground is highly cultivated, and in the fields about town, it does a deal of mischief: by the quickness of its growth and the great number of its shoots, it covers and chokes many young plants; hence it should be carefully weeded from dunghills.


The seeds are very beautiful, and have the greatest affinity to those of the *Cerastium aquaticum*.

When the flowers first open, the foot-stalks which support them are upright; as the flowers go off they hang down; and when the seeds become ripe, they again become erect.

LINNÆUS has observed that the flowers open from nine in the morning till noon, unless rain falls on the same day, in which case they do not open: from what little observations I have made on this plant, it is not subject to be affected precisely in the same manner here, having seen in the month of March, the blossoms continue rather widely expanded after repeated showers of rain.

It is considered as a wholesome food for Chicken and small Birds, whence, as RAY observes, it has obtained its name: boiled it resembles Spinach so exactly as scarcely to be distinguished from it, and is equally wholesome, being a plant which may be procured almost any where very early in the spring, it may be no bad substitute where Spinach or other greens are not to be had in plenty, and much preferable to Nettle-tops and other plants which the lower sort of people seek after in the spring with so much avidity. Swine are very fond of it, and prefer it to Turnep-tops. It is eaten by many insects, particularly by the Caterpillar of the *Phalena Villica* or *Cream Spot Tiger Moth*, and other hairy Caterpillars of the *Tyger* kind.

As a medicine it contains no active principle; but is frequently applied to hot, painful, and inflammatory swellings, either by itself, bruised, or mixed with poultices, with good success.



ERICA TETRALIX. CROSS-LEAVED HEATH.

ERICA *Linnæi* Gen. Pl. OCTANDRIA MONOGYNIA.

Cal. 4-phyllus. Cor. 4-fida. Filamenta receptaculo inserta. Antheræ bifidæ.

Caps. 4-locularis.

Raii Syn. ARBORES ET FRUTICES.

ERICA *tetralix* foliis quaternis ciliatis, floribus capitatis imbricatis.

ERICA *tetralix*, antheris aristatis, corollis ovatis, stylo incluso, foliis quaternis ciliatis, floribus capitatis.

Linn. Syst. Vegetab. p. 302. Fl. Suecic. n. 337.

ERICA ex rubro nigricans scoparia. Baubini Pin. 486.

ERICA Brabantica folio Coridis hirsuto quaterno. I. B. 1. 358.

ERICA pumila Belgarum Lobelio, scoparia nostras. Parkinson. 1482.

ERICA major flore purpureo. Gerard. emac. 1382. Raii Syn. p. 471, Low Dutch Heath or Besome Heath.

Hudson Fl. Angl. p. 144. Oeder Fl. Dan. icon. 81.

CAULES fruticosi, dodrantes aut pedales, ramosi, fusci, scabriusculi ex reliquantis foliorum.

STALKS shrubby, about nine or twelve inches high, branched, roughish from the remains of the leaves which have fallen off.

FOLIA quaterna, ovato-linear, patentia, prope flores cauli adpressa, marginibus inflexis, ciliatis, ciliis glandulâ terminatis, superficie superiore plana, inferiore concava.

LEAVES growing by fours, of an oval-linear shape, spreading, near the flowers pressed close to the stalk, the edges turned in and ciliated or hairy, each of the hairs terminating in a small round globule, the upper surface flat, the inferior surface concave.

FLORES secundi, imbricati, in capitulum congesti, carnei.

FLOWERS hanging down one over another all one way, forming a little head, of a pale red colour.

CALYX: PERIANTHIUM hexaphyllum, foliolis hirsutis, duo inferiora ovato-lanceolata, cætera linearia, fig. 2.

CALYX: a PERIANTHIUM of six leaves, the leaves hairy, the two lowermost of an oval-pointed shape, the rest linear, fig. 2.

COROLLA ovata, monopetala, ore quadrifido, laciniiis reflexis, fig. 3.

COROLLA oval, monopetalous, the mouth divided into four segments, which turn back, fig. 3.

STAMINA: FILAMENTA octo, subulata, alba, corollâ breviora, receptaculo inserta; ANTHERÆ sagittatæ, conniventes, purpureæ, biforaminosæ, bicornes, fig. 4, 5, 6.

STAMINA: eight FILAMENTS, tapering, white, shorter than the Corolla, inserted into the receptacle; ANTHERÆ arrow-shaped, closing together, purple, having two apertures for the discharge of the Pollen, and two little horns, fig. 4, 5, 6.

PISTILLUM: GERMEN cylindraceum, subfulcatum, villosum, glandulâ ad basin cinctum, fig. 7, 8; STYLUS filiformis, purpurascens, fig. 9; STIGMA, obtusum, fig. 10.

PISTILLUM: GERMEN cylindrical, slightly grooved, villous, surrounded at bottom by a gland, fig. 7, 8; STYLE filiform, purplish, fig. 9. STIGMA blunt, fig. 10.

PERICARPIUM: CAPSULA subrotunda, villosa, apice truncata, quadrivalvis, fig. 11, 12.

SEED-VESSEL: a roundish CAPSULE covered with a kind of down, cut off as it were at top, having four valves, fig. 11, 12.

SEMINA plurima, minuta, flavescencia, fig. 13, 14.


SEEDS numerous, minute, and yellowish, fig. 13, 14.


THIS species of *Heath*, though not applicable to such a variety of uses as some of the others, is not inferior to any of them in the beauty and delicacy of its flowers, which in general are of a pale red colour, but sometimes they occur entirely white.

It is obviously enough distinguished from the rest, not only by its flowers growing in a kind of pendulous cluster on the tops of the stalks, but by its leaves also, which growing by fours on the stalk, form a kind of cross; these are edged with little stiff hairs, each of which has a small globule at its extremity.

At the latter end of the Summer it contributes its share with the others to decorate and enliven those large tracts of barren land which too often meet the eye in many parts of this kingdom.

It delights to grow in a moister situation than some of the others, and will thrive well enough in gardens, if taken up either in Spring or Autumn with a quantity of earth about its roots: this is necessary, as the *Heath* in general bear transplanting ill.





POLYGONUM BISTORTA. THE GREATER BISTORT OR SNAKE-WEED.

POLYGONUM *Linnei Gen. Pl.* OCTANDRIA TRIGYNIA.

Raii Synopsi, Genus quintum. HERBÆ FLORE IMPERFECTO SEU STAMINEO, (VEL APETALO POTIUS.)

POLYGONUM *Bistorta* caule simplicissimo, monostachyo, foliis ovatis in petiolum decurrentibus. *Linnei Syst. Vegetab. p.* 311.

POLYGONUM radice lignosa contorta, spica ovata, foliorum petioliis alatis. *Haller. Hist. v.* 2. 258.

COLUBRINA Seu Serpentina femina. *Fuschii icon.* 774.

SERPENTARIA mas five Bistorta. *Fuschii icon.* 773.

BISTORTA major radice minus intorta. *Baubin. Pin.* 192.

BISTORTA major radice magis intorta. *Baubin. Pin.* 192.

BISTORTA major *Gerard emac* 399. major vulgaris *Parkinson* 391. *Raii Synopsi* 147. *Hudson. Fl. Angl.* 146. *Flor. Dan. Ic.* 421.

RADIX perennis, crassitie digiti, plus minusve intorta, externe castanea, interne carnea, fibris et stolonibus plurimis instructa.

CAULIS pedalis aut bipedalis, simplex, suberectus, foliis, articulatus, (geniculi tumidi) teres, laevis.

STIPULÆ vaginantes, apice membranaceæ, marcescentes, ore obliquo.

FOLIA cordato-lanceolata, undulata, subtus cærulescentia, glabra, inferiora in petiolo decurrentia, superiora amplexicaulia in stipulis desinentia.

FLORES spicati, spica oblongo-ovata, densa.

BRACTEÆ membranaceæ, marcescentes, biflores, bivalves, valvula inferiore tricuspidata cuspidata medio longiore quasi aristata, flores pedicellati, pedicellis calyce longioribus.

CALYX five COROLLA subovata, quinquepartita, carnea, laciniis ovatis, obtusis, concavis. *fig.* 1. 3.

STAMINA: FILAMENTA octo, subulata, alba, corollâ longiora, ANTHERÆ biloculares, purpurascens, incumbentes. *fig.* 2.

PISTILLUM: GERMEN triquetrum, sanguineum, STYLIS tres longitudine staminum; STIGMATA parva, rotunda. *fig.* 5. 6. 7.

NECTARIUM. glandulæ rubræ in fundo calycis. *fig.* 4.

SEMEN triquetrum, fuscum, mucronatum, nitens, vernice quasi obductum. *fig.* 8.

ROOT perennial, the thickness of one's finger, more or less crooked, externally of a chestnut, internally of a flesh colour, furnished with numerous fibres and creepers.

STALK from one to two feet high, simple, nearly upright, solid, jointed, (the joints swelled,) round and smooth.

STIPULÆ enclosing the Stalk as in a sheath, at top membranous, withered, the mouth oblique.

LEAVES: the bottom leaves somewhat heart shaped and pointed; waved at the edges, smooth, underneath blueish and continued down the footstalks, the upper leaves embracing the stalk, and terminating in the stipule.

FLOWERS growing thickly in a spike, the spike of an oblong oval shape.

FLORAL LEAVES membranous, and withered, containing two flowers and having two valves, the lower valve three pointed, the middle point running out into a kind of arista or beard, the flowers growing on footstalks which are longer than the Calyx.

CALYX or COROLLA, of an oval shape and flesh coloured, divided into five segments, which are oval, obtuse, and concave. *fig.* 1. 3.

STAMINA: eight FILAMENTS, tapering, white, and longer than the Calyx; the ANTHERÆ bilocular, purplish, and laying across the filaments. *fig.* 2.

PISTILLUM the GERMEN three square, of a deep red colour, three STYLES the length of the Stamina; the STIGMATA small and round. *fig.* 5. 6. 7.

NECTARIUM: several small red glands in the bottom of the Calyx. *fig.* 4.

SEED: triangular, brown, pointed, and shining as if varnished. *fig.* 8.

WHEN a Plant not intended to be cultivated, in any respect prevents the growth of one which is the object or Cultivation, such a plant, however beautiful, may with propriety be called a Weed; nor will the elegance or utility of the Bistort, secure it in the estimation of the Farmer, from that appellation.

This Plant generally grows in moist Meadows, and flowers in May and June; when it has once taken root, it propagates very fast, and frequently will form large patches, to the exclusion of a considerable portion of the Grass; nor is it destroyed but with the greatest difficulty. Happily, our Farmers about Town are pretty much strangers to this Plant, as it is met with but rarely. It grows plentifully in a Meadow by the side of *Bishop's Wood* near *Hamstead*, and my obliging Friend Dr. ALLEN informs me he has found it about *Battersea*.


As an astringent Medicine, the Bistort appears to possess considerable virtue, and as such may with propriety be made use of in all cases where astringents are required; but more particularly in long continued evacuations from the Bowels, and other discharges both serous and sanguinous. It is recommended also to taken teeth which are loose, and may be used either in powder, infusion, or extract. It could be procured in sufficient quantity to make it answer. It might well be applied to the purpose of tanning Leather.

In some parts of England the leaves are eat as a Pot-herb.



— *Polygonum* — *Bistorta* —







Polygonum Persicaria.

POLYGONUM PERSICARIA. COMMON SPOTTED PERSICARIA.

- POLYGONUM *Linnaei Gen. Pl.* OCTANDRIA TRIGYNIA.
Raii Syn. Gen. 5 HERBÆ FLORE IMPERFECTO SEU STAMINEO, VEL APETALO POTIUS.
 POLYGONUM *Persicaria floribus hexandris semidigynis, pedunculis lævibus, stipulis ciliatis, spicis ovato-oblongis erectis.*
 POLYGONUM *Persicaria floribus hexandris digynis, spicis ovato-oblongis, foliis lanceolatis, stipulis ciliatis, Lin. Syst. Vegetab. p. 312. Flor. Suecic. p. 130.*
 POLYGONUM *foliis ovato-lanceolatis, subhirsutis, spicis ovatis, vaginis ciliatis. Haller. hist. Helv. v. 2. p. 257.*
 PERSICARIA *mitis maculosa et non maculosa. Bauhin. Pin. p. 101.*
 PERSICARIA *maculosa Gerard. emac. 445. vulgaris mitis seu maculosa. Parkinson. 856. Raii Syn. ed. 3. p. 145. n. 4. Dead or spotted Arimart. Hudson Flor. Angl. p. 147. n. 4. Scopoli Fl. Carniol. p. 279.*

RADIX simplex, fibrosa.

CAULIS erectus, ad basin aliquando repens, pedalis ad tripedalem, ramifusus, teres, glaber, ad geniculos sensim incrassatus, sæpe rubens: sub geniculis puncta radicalia discernantur quamvis huic speciei non propria.

RAMI alterni, e singulo geniculo prodeuntes, patentes, sæpe diffusi.

STIPULÆ vaginantes, liquore viscido sæpe repletæ, ciliatæ.

FOLIA lanceolata, subpetiolata, margine nervoque medio subhirsutis, utrinque lævia, maculâ ferrugineam quodammodo referente sæpius notata.

PEDUNCULI læves.

FLORES spicati, rosei, Spicæ terminales, erectæ, subovatae.

CALYX: PERIANTHIUM quiquepartitum, coloratum, persistens, segmentis ovatis obtusis, fig. 1, 2.

COROLLA nulla.

STAMINA: FILAMENTA sex fundo calycis inserta longitudine corollæ ANTHERÆ rubentes, fig. 2.

PISTILLUM: GERME ovatum, compressum, aut triquetrum, fig. 3, 6. STYLUS ad medium usque bifidus sæpe trifidus, fig. 5, 8. STIGMATA duo aut tria subrotunda, fig. 4, 7.

SEMEN unicum, nitidum, aut subovatum, acuminatum, ad unum latus leviter convexum; fig. 9, 11, aut trigonum, fig. 10, 12.

ROOT simple and fibrous.

STALK upright, sometimes creeping at bottom, from one to three feet high, branched, round, smooth, gradually thicker at the joints, often of a red colour: a little beneath each joint some radical points are observable, which however are not peculiar to this species.

BRANCHES alternate, proceeding from each joint, spreading, frequently very much so.

STIPULÆ embracing the stalk, frequently full of a viscid liquid, and terminated by long ciliae or hairs.

LEAVES lanceolate, with short foot-stalks, the edge and midrib slightly hairy, smooth on both sides, in general having a large spot on the middle of the leaf somewhat like a horse shoe.

FOOT-STALKS of the flowers, smooth.

FLOWERS growing in spikes, of a bright rose colour, the spikes terminal, upright, of a somewhat oval shape.

CALYX: a PERIANTHIUM divided into five segments, coloured, and persisting, the segments oval and obtuse, fig. 1, 2.

COROLLA wanting.

STAMINA: six FILAMENTS inserted into the bottom of the Calyx, the length of the Corolla; the ANTHERÆ redish, fig. 2.

PISTILLUM: GERME oval and flat, or three-square, fig. 3, 6. STYLE divided down to the middle into two, often into three parts, fig. 5, 8. STIGMATA two or three, and round, fig. 4, 7.

SEED one, shining, either of an oval pointed shape and slightly convex on one side, fig. 9, 11. or three-square, fig. 10, 12.

The very great similarity which exists between the several species of the Polygonums, has occasioned no small degree of trouble to Botanists, in rightly ascertaining the limits of each Species and Variety; a difficulty not to be overcome while Books are consulted more than Nature. Sensible of the truth of this observation, and earnestly desirous of arriving at some certainty on this subject, we have examined a vast number of all the different Species and Varieties of Polygonum which our neighbourhood affords, compared them with one another, sown the seeds, and cultivated many of them; and if we do not deceive ourselves, have reduced some of the more difficult ones to their true Species and Varieties.

As what we relate concerning these plants is no more than the result of the most accurate and repeated investigation, assisted by the microscope, we shall be the less concerned because we differ from Authors of the most respectable Authority.

The writer who gives an account of all the known plants in the universe, cannot be supposed to have the opportunity of being so minute in his enquiries as one who describes the plants of a particular spot, which as they grow are constantly the objects of his attention.

We have ventured to alter Linnaeus's Specific description of this plant, which stands thus.

Polygonum floribus hexandris digynis, spicis ovato-oblongis, foliis lanceolatis, stipulis ciliatis. to

Polygonum floribus hexandris semidigynis, pedunculis lævibus, stipulis ciliatis, spicis ovato-oblongis erectis.

We have not made this alteration from an idle desire of differing from so great a Man, whom we truly respect and revere, but solely to make the distinctions betwixt those plants more obvious, and thereby add our mite to the general stock of Botanic knowledge. In specific descriptions, the distinguishing marks should as much as possible be contrasted or opposed to each other, in these plants this does not seem to have been sufficiently attended to. What we have principally in view by altering the Specific description is to distinguish it from the *Polygonum Pensylvanicum* and its varieties, of which there are several, and to which the *Polygonum Persicaria* in its general habit is exceeding nearly allied.

In all the flowers of this Species which we have examined, the Style has been divided just half way down, hence we have called the flowers *Semidigyni*, had it been divided down to the base they would with propriety have been called *Digyni*. In most of the flowers the Style is divided into two parts, and the Germen is a little convex on each side, in some of the flowers the style is divided into three, hence those flowers might be called *Semitriginis*, and when this is the case the Germen is always triangular. In the *Polygonum Pensylvanicum* the Style is divided nearly to the base, this difference then in the division of the Style, is of considerable consequence in distinguishing the two Species and their varieties from each other.

The footstalks which support the flowers in this Species, are quite smooth, in the *Polygonum Pensylvanicum*, they are beset with a great number of minute glands, which gives them a manifest roughness, and contributes to characterize this Species.


The Stipulæ are furnished with long Ciliae or Hairs, particularly towards the top of the plant, in the *Polygonum Pensylvanicum* these are wanting. These two plants likewise differ much in the form of their seeds, of which we shall speak more fully in our account of the latter.

The flowers always grow in upright spikes of an oval shape more or less round; by these two characters this Species is at once distinguished from the *Polygonum Hydropiper*, the spikes of which are *filiform and pendulous*.

The leaves are most commonly spotted, but this is neither constant nor peculiar to this Species, and difference of size only forms the principle variety to which it is subject.

It grows exceedingly common in all our Ditches, and flowers in August and September; its blossoms are beautiful and last a considerable time, and it is not so common, it would probably be thought worthy of a place in our Gardens.

No particular virtues or uses are attributed to it.



POLYGONUM PENSYLVANICUM. PALE - FLOWERED PERSICARIA.

POLYGONUM *Linnaei Gen. Pl. OCTANDRIA TRIGYNIA.*

Raii Syn. Gen. 5. HERBÆ FLORE IMPERFECTO SEU STAMINEO (VEL APETALA POTIUS.)

POLYGONUM floribus hexandris, digynis; stipulis muticis; pedunculis scabris; seminibus utrinque depressis.

POLYGONUM floribus octandris digynis, pedunculis hispidis, foliis lanceolatis, stipulis muticis.

Linnaei Syst. Vegetab. Sp. Plant. p. 519.

PERSICARIA mitis major foliis pallidioribus. *D. Bobarti, Dead Asmart the greater with pale leaves.*

Raii Syn. ed. 3. p. 145. Hudson Fl. Angl. p. 148.

RADIX fibrosa, annua.

CAULIS tripedalis circiter, teres, glaber, fistulosus, ramosus; rami patentes, geniculis maxime incrassatis.

FOLIA ovato-lanceolata, supra glabra, subtus glandulis punctata, sæpe pubescentia, ciliata, nunc maculata nunc immaculata.

PETIOLI subtus hirsuti, scabriusculi.

STIPULÆ basi nervosæ, muticæ.

PEDUNCULI pilis brevibus glanduliferis scabri. *fig. 1.*

FLORES herbacei, pedunculis brevibus insidentes, dense glomerati, spicæ ovatæ, seminibus maturis subnantes.

CALYX: PERIANTHIUM quinquepartitum, laciniiis ovatis, obtusis, *fig. 2, 3.*

COROLLA nulla.

STAMINA: FILAMENTA sex, subulata, alba, Corollâ paulo breviora; ANTHERÆ biloculares; POLLEN globosum, *fig. 4.*

PISTILLUM: GERME subovatum; STYLUS fere ad basin divisus; STIGMATA duo subrotunda, *fig. 5, 6.*

SEMEN cordatum, acuminatum, compressum, medio depressum, nitidum, *fig. 9, 10,* magnit. nat. *fig. 7, 8,* lente auct. subinde obtuse triquetrum, *fig. 12.*

ROOT fibrous and annual.

STALK about three feet high, round, smooth, hollow, branched, the branches spreading, and the joints very much swelled.

LEAVES of an oval pointed shape, smooth on their upper surface, underneath dotted with small glands, and often downy, edged with little hairs, sometimes with and sometimes without spines.

FOOT-STALKS of the leaves hairy underneath, with a slight roughness to the touch.

STIPULÆ rib'd at bottom, and not terminated by any hairs.

FOOT-STALKS of the flowers rough with little glands. *fig. 1.*

FLOWERS of a greenish colour, sitting on short foot-stalks, and growing thickly together; spikes oval, and when the seeds are ripe drooping a little.

CALYX: a PERIANTHIUM divided into five segments, which are oval and obtuse, *fig. 2, 3.*

COROLLA wanting.

STAMINA: six FILAMENTS, tapering, white, a little shorter than the Corolla; ANTHERÆ bilocular; POLLEN globular, *fig. 4.*

PISTILLUM: GERME somewhat oval; STYLE divided nearly down to the base; STIGMATA two, roundish, *fig. 5, 6.*

SEED heart-shaped, pointed, flat, with a depression in the middle, shining, *fig. 9, 10,* of its natural size, *fig. 7, 8,* magnified, sometimes obtusely triangular, *fig. 12.*

The plant here figured, is the *Persicaria mitis major foliis pallidioribus*, *D. Bobarti*, and which is particularly described in the 3d. edition of *RAY's Synopsis*, p. 145; from the consonancy of this description, with that which LINNÆUS had given of the *Polygonum Pensylvanicum*, in the 3d. edition of his *Species Plantarum*, Mr. HUDSON set it down in his *Flora* as that species; and LINNÆUS, in the last edition of his *Systema Vegetab.* as a confirmation of our English *Polygonum's*, being the same with his *Pensylvanicum*, quotes BOBART's descriptive name.

By *RAY*, LINNÆUS, and HUDSON, then, it is made a distinct species; by *HALLER* it is considered as a variety of the *Polygonum Persicaria*; but as the Baron forms his judgment from dried specimens that were sent him, in which many of the distinguishing characters of this plant would be unavoidably lost, he seems the most likely to be mistaken: I shall therefore join in making it a distinct species; and I trust shall give such striking additional characters, as will settle this matter beyond dispute.


The true *Polygonum Pensylvanicum* (for there are several varieties of it) has the greatest affinity with the *Polygonum Persicaria*, but differs from it in the following particulars, viz. place of growth, size, stipulæ, leaves, foot-stalks of the leaves, foot-stalks of the flowers, style, and seeds.


While the *Polygonum Persicaria* usually delights to grow by the sides of moist ditches, the *Pensylvanicum* prefers a richer and more luxuriant soil; and so common is it with us about town, that there is scarce a dunghill on which it may not be found: indeed in its attachment to this particular soil, it resembles many of the *Chenopodium* or *Oraches*. Was it never to occur in other situations, some might be ready to suspect that it was a variety of the *Persicaria* arising from richness of soil; but it is frequently found in other places: and I remember once to have seen the *Polygonum Persicaria*, *Hydropiper*, and *Pensylvanicum*, all growing by the side of a stream within six inches of each other.

In its most common state it is much larger than the *Polygonum Persicaria*, and its joints in particular are more swelled; its Stipulæ are much more strongly rib'd at bottom, and have no Ciliæ; its leaves are broader, the veins somewhat deeper and more strongly marked; the hairs on the edges of the leaves more visible, but particularly so under the foot-stalk of the leaf, to which they give a manifest roughness: in the uppermost leaves the under side is generally dotted with very minute glands, while in the lowermost it is covered with a kind of down: this last character, though contrary to what LINNÆUS asserts, is never seen in the *Polygonum Persicaria*; but in this species it is always more or less predominant. The foot-stalks of the flowers are thickly beset with little yellowish glands, standing on short foot-stalks, which sometimes extend half way down the plant; this appearance never or exceeding rarely occurs in the *Polygonum Persicaria*: the flowers are of a pale or greenish hue, and form thicker and larger spikes than in the *Polygonum Persicaria*, and when ripe are so heavy as frequently to hang down a little: the Style is divided very nearly down to the Germen, while in the *Polygonum Persicaria* it is divided only half way; and this division of the Style, I look upon as one of the most constant and certain criteria of this species: lastly the form of the seeds contributes not a little to the farther ascertaining and fixing it; in the *Persicaria* the seeds are either triangular, or of a pointed oval shape, with a little convexity on each side; in this species it is in general flat, with a depression on each side; it is also larger and broader; now and then a seed occurs forming an unequal triangle, but these are very rare, while the triangular seed is most frequent in the *Polygonum Persicaria*.



Polygnum Pennsylvanicum







Polygonum Pensylvanicum var. *caule maculato*.

POLYGONUM PENNSYLVANICUM. VAR. CAULE MACULATO. SPOTTED-STALK'D PERSICARIA.

PERSICARIA latifolia geniculata, caulibus maculatis. *D. Rand. Raii Syn. p. 145.*

PERSICARIA maculosa procumbens foliis subtus incanis. *Raii Syn. p. 146. eadem est planta solo autem minus læto proveniens.*

Such then is the difference, which from repeated examinations, I have been able to discover betwixt the *Polygonum Persicaria* and the *Pennsylvanicum* in its most common state; in this state however it does not always occur, but is subject to more Varieties than any of our other *Persicaria's*: without any desire of multiplying them, I make the following, having found them all about *London*:

- 1 *Polygonum Pennsylvanicum. var. caule et floribus rubris.*
- 2 *caule maculato.*
- 3 *foliis subtus incanis.*

The first of these varieties is very often found with the true species on dunghills, as also in corn-fields, and is like it in every respect excepting its colour, the stalks and flowers being red, but not so beautifully bright as those of the *Polygonum Persicaria*.

The second variety here figured, which indeed comes near to a distinct species, grows much in the same situations, and oftentimes with the *Polygonum Persicaria* in the ditches about *St. Georges-fields*, particularly in a large ditch on the right-hand side of the road between the end of *Blackman-Street* and *Newington*, where it is very common in the month of *September*. It not only differs from the other in having its stalk spotted with red, a character which it keeps very constantly, but its spikes are much slenderer, rather more so even than those of the *Persicaria*, of a red colour, but not quite so bright as those of that plant: the under side of the foot-stalk of the leaves is remarkably rough; the little glands on the foot-stalks of the flowers, and the parts of the fructification are similar to those of the true species, but the seeds are smaller: when this variety grows in the rich soil abovementioned, it is full as large as the *Pennsylvanicum* itself; but when it grows in a different soil and situation, as on the watery parts of *Blackheath* and *Peckham-Rye*, it becomes much smaller, generally has its leaves white underneath, and will certainly be taken for the *Polygonum Persicaria* if not attentively examined: its spotted stalk and the roughness of the foot-stalks of the leaves will however readily discover it.

The third variety, with leaves hoary on the under side, is found here and there in corn-fields and other places, where the soil is not very rich, and is obviously enough distinguished.


Besides these striking varieties, it is subject, like all other plants, to vary in size according to the richness or poverty of the ground on which it grows, and like the *Polygonum Persicaria*, its leaves are sometimes spotted and sometimes not.

This descriptive account will perhaps appear tedious and uninteresting to some; if however by these practical observations, the obscurity which has hitherto dwelt on this difficult Genus, shall in some degree be removed, and the road of investigation made easier to the young Botanist, I shall think my time usefully employed; I would not however wish him to take upon trust what is here advanced, but to examine each plant and its several parts for himself; thus he will become improved, and be able perhaps to throw a still greater light on the subject.

The Sparrow and other small Birds are very fond of the seeds of this species and its varieties: but the Farmer should carefully weed them from his dunghills.

The first of these is the fact that the
 number of cases of disease is not
 proportional to the number of persons
 exposed to the disease.

The second is the fact that the
 number of cases of disease is not
 proportional to the number of persons
 exposed to the disease.



POLYGONUM HYDROPIPER. BITING PERSICARIA OR WATER PEPPER.

POLYGONUM *Linnaei Gen. Pl.* OCTANDRIA TRIGYNIA.

Cal. o. Cor. 5-partita, calycina. Sem. 1, angulatum.

Rai Syn. Gen. HERBÆ FLORE IMPERFECTO SEU STAMINEO VEL APETALO POTIUS.

POLYGONUM *Hydropiper floribus hexandris femidigynis; foliis lanceolatis, undulatis, immaculatis; spicis filiformibus nutantibus.*

POLYGONUM *Hydropiper floribus hexandris femidigynis, foliis lanceolatis, stipulis submuticis. Linn. Syst. Vegetab. p. 312.*

POLYGONUM *foliis ovato lanceolatis, spicis florigeris, vaginis calvis. Haller. bist. p. 256. n. 1554.*

POLYGONUM *Hydropiper. Scopoli Fl. Carniol. n. 467.*

PERSICARIA *urens seu Hydropiper. Baubin. pin. 101.*

PERSICARIA *vulgaris acris seu minor. Parkinson. 856.*

HYDROPIPER. *Gerard. emac. 445. Raii Syn. p. 144. Water-pepper, Lakeweed or Arsmart. Hudson Fl. Angl. p. 148.*

RADIX annua, fibrosa.

CAULIS erectus, ramosus, basi nonnunquam repens, pedalis ad tripedalem, geniculis incrassatis, demum ruberrimus.

FOLIA lanceolata, undulata, c viridi flavescentia, glabra.

STIPULÆ ciliatæ.

FLORES spicati, spicæ tenues, demum nutantes.

CALYX: PERIANTHIUM quadripartitum, glandulis minimis adpersum, laciniis obtusis, concavis, fig. 1, 2, 3.

COROLLA nulla.

STAMINA: FILAMENTA sex alba; ANTHERÆ albæ biloculares, fig. 3.

PISTILLUM: GERMEN ovatum; STYLUS bifidus, ad medium usque divisis; STIGMATA duo, rotunda, fig. 4, 5.

SEMEN ovato-acuminatum, castaneum, fig. 6.

ROOT annual and fibrous.

STALK upright, branched, sometimes creeping at bottom, from one to three feet high, the joints swelled, finally becoming very red.

LEAVES lanceolate, waved, of a yellowish green colour and smooth.

STIPULÆ ciliated.

FLOWERS growing in spikes, which are slender and finally drooping.

CALYX: a PERIANTHIUM divided into four segments, sprinkled with very minute glands, the segments blunt and hollow, fig. 1, 2, 3.

COROLLA wanting

STAMINA six white FILAMENTS; ANTHERÆ white and bilocular, fig. 3.

PISTILLUM: GERMEN oval; STYLE bifid, divided down to the middle; two round STIGMATA, fig. 4, 5.

SEEDS of an oval pointed shape, and chestnut colour, fig. 6.

It is one of the maxims laid down by the Author of that system of Botany which at present is so deservedly held in esteem, and which I trust for the sake of this delightful science will for ever withstand the attempts of all those who frame systems merely to raise themselves into consequence, that in all specific descriptions taste is to be excluded: some may perhaps be ready to treat this as too dogmatical, but when they come to find that both the *Hydropiper* and *Sedum acre*, plants which in general are very hot and biting, sometimes are found insipid, they will readily adopt it as founded in strict propriety.

The present species of *Polygonum* very properly receives its name of *Hydropiper* from its hot and biting taste, which appears to arise from its essential oil dispersed in little cells or glands all over the plant, but more particularly observable on the Calyx with a small magnifier, and which, if tasted, will be found to be more biting than any other part of the plant: this quality which is peculiar to the *Hydropiper*, generally leaves a strong idea of the plant on the mind of the Tyro: but it has other more invariable characters whereby it may be distinguished. Notwithstanding its obvious difference from the other plants of this genus, apparent even to such as know very little of Botany, both *Scopoli* and *HALLER* seem to entertain doubts whether it be really distinct from the *P. Persicaria* and *P. minus*.

The three plants as they usually grow, and I have seen them all three grow together, are certainly distinct enough: but there are some intermediate varieties which bring them very near together, and perhaps justify such suspicions: a variety of the *Hydropiper*, scarce differing in any other respect but its insipidity, I have now and then met with in the same situation as we usually find the true species: from the *P. Persicaria* it differs principally in its leaves, spikes, form and size of its seeds; and first its leaves are of a yellower hue, more undulated, and never marked with any spots; its spikes are slender, and when the seeds are ripe they bend and hang down; the seeds are much larger, more acuminate, and of a chestnut colour; its stipule are very evidently ciliated; though *HALLER* makes their mislead.


It is the only *Persicaria* that has any pretensions to be an active medicine: given in infusion or decoction it proves diuretic, hence it is made use of in the Dropsy and Jaundice; and the distilled water of it is recommended by *BOYLE* as efficacious in the Stone and Gravel: *LINNÆUS* informs us, that the plant will dye Woollen cloth of a yellow colour.

Although the herb is so acrid, the seeds are insipid and nutritive.

It is found in great abundance in all those places which lie under water during the Winter, flowers in September, generally a month later than the *P. Persicaria*: in exposed places it becomes very red in going off.



Polygonum Hydropiper.





Polygonum aviculare.

POLYGONUM AVICULARE. BIRDS POLYGONUM OR KNOT-GRASS.

POLYGONUM *Linnei Gen Pl. OCTANDRIA TRIGYNIA.*

Cal. o. Cor. 5-partita, Calycina. Sem. 1. angulatum.

Raii Syn. Gen. 5. HERBÆ FLORE IMPERFECTO SEU STAMINEO. (VEL APETALA POTIUS.)

POLYGONUM *aviculare* floribus oclandris trigynis axillaribus, foliis lanceolatis, caule procumbente herbaceo. *Linn. Syst. Vegetab. p. 312. Sp. Pl. 519. Fl. Suecic. n. 339.*

POLYGONUM procumbens, foliis linearibus, acutis, floribus solitariis. *Haller hist. n. 1560.*

POLYGONUM *aviculare. Scopoli Fl. Carniol. n. 471.*

POLYGONUM mas vulgare. *Gerard emac. 451.*

POLYGONUM mas vulgare majus. *Parkinson 443.*

POLYGONUM seu Centinodia. *I. Baubin 3. 374. Raii Syn. p. 146. Hudson Fl. Angl. p. 149.*

RADIX annua, simplex, lignosa, multis fibris donata, terram firmiter apprehendens ut extirpatu difficilis sit, sapore adstringente.

ROOT annual, simple, woody, furnished with many fibres, taking strong hold of the earth, so as to be with difficulty pulled up, and of an astringent taste.

CAULES plures, plerumque procumbentes, interdum vero suberecti, dodrantales, ramosi, tenues, striati, laeves, teretes, geniculati, ad geniculos paululum incrassati.

STALKS several, generally procumbent, sometimes nearly upright, about nine inches in length, branched, slender, striated, smooth, round, jointed, the joints a little swelled.

FOLIA quam maxime variantia, ovata, lanceolata aut etiam linearia, alterna, lævia, e vaginis stipularum prodeuntia.

LEAVES varying exceedingly, oval, lanceolate, or sometimes even linear, alternate, smooth, proceeding from the sheaths of the Stipulae.

STIPULÆ vaginantes, membranaceæ, albida, nitida, apice fibrosa.

STIPULÆ forming a sheath round the joints, membranous, white, shining, at top fibrous.

FLORES axillares, e vaginis stipularum cum foliis prodeuntia.

FLOWERS axillary, proceeding with the leaves from the sheaths of the Stipulae.

CALYX: PERIANTHIUM quinquepartitum, laciniis ovatis concavis, patentibus, dimidio inferiore viridi, superiore albo, sæpe colorato, fig. 1, 2.

CALYX: a PERIANTHIUM divided into five segments, the laciniæ oval, concave and spreading, the lower half green, the upper half white and often coloured, fig. 1, 2.

COROLLA nulla.

COROLLA wanting.

STAMINA: FILAMENTA octo corolla breviora; ANTHERÆ flavæ, fig. 2, auct.

STAMINA: eight FILAMENTS shorter than the Corolla, ANTHERÆ yellow, fig. 2, magnified.

PISTILLUM: GERMEN triquetrum; STYLUS longitudine staminum, trifidus; STIGMATA tria, rotunda, fig. 3, auct.

PISTILLUM: GERMEN triangular; STYLE the length of the Stamina, trifid; STIGMATA three, round, fig. 3, magnified.

SEMEN triquetrum, nigricans, intra calycem, fig. 4.


SEED triangular, of a blackish colour, contained within the Calyx, fig. 4.

Those plants which have been observed to be eaten by cattle, have often obtained the name of *Grass*, although they have not possessed the least similitude to those which are real Grasses, and the present plant is one of these. Cattle in general are fond of it, and hogs in particular eat it with great avidity. The seeds afford sustenance to many of the small birds, whence it has acquired the name of *aviculare*. The Caterpillar of the *Phalena runcicis* (with us the *Knot-grass Moth*;) I have frequently found feeding on its leaves, although it is by no means confined to this plant: in *Sweden*, LINNÆUS informs us it feeds on the *Dock* (*Rumex*;) and *Scotibible*.

This species of *Polygonum* may be considered as one of our most common plants; it delights to grow in a sandy or gravelly soil, on banks, and by the sides of roads and paths, being of quick growth, and spreading a great deal of ground; it often covers whole fields, that by turning in of cattle, have had their natural coat of grass destroyed.

Where a plant of this species happens to grow singly in a rich soil, it will often cover the space of a yard or more in diameter, and the leaves become broad, and large; but when it grows very thick together, by the sides of paths, it is in every respect smaller, and the stalks are more upright. It is subject, like most other plants, to several varieties, and of these are the *Polygonum brevi angustaque folio*, and the *Polygonum oblongo angustaque folio* of *C. Baubine*.

It has been considered by ancient writers, as possessing some medical virtue, particularly as an Astringent, and is by them recommended in Diarrhæas, Dysenteries, Bleeding at the nose, and other Hemorrhages; but in the present practice, its use seems justly superseded by more efficacious medicines.





Polygonum minus.



POLYGONUM MINUS. SMALL, CREEPING, NARROW-LEAVED

PERSICARIA.

POLYGONUM *Linnaei* Gen. Pl. OCTANDRIA TRIGYNIA.

Cal. o. Cor. 5-partita calycina. Sem. 1, angulatum.

Raii Syn. Gen. 5. HERBÆ FLORE IMPERFECTO SEU STAMINEO VEL APETALA POTIUS.

POLYGONUM *minus* floribus hexandris, submonogynis, foliis lineari-lanceolatis, caule basi repente.

POLYGONUM *minus* hexandris digynis foliis lanceolatis, stipulis ciliatis, caule divaricato patulo.

Hudson Fl. Angl. p. 148.

POLYGONUM foliis ovato-lanceolatis, glabris, spicis strigosis, vaginis ciliatis. *Haller. hif. p. 257.*

n. 1555.

PERSICARIA *minor. Bauhin Pin. 1014? angustifolia. Bauhin Pin. 101. 3?*

PERSICARIA *puflillarepens Ger. emac. 446. Parkinson. 857. Raii Syn. 145. 2. Small Creeping Arismart.*

PERSICARIA *angustifolia ex singulis geniculis florens. Mer. Pin. 90? Raii Syn. 145. 3. Narrow-leaved Lakeweed.*

RADIX annua, fibrosa.

CAULES plures, dodrantes, aut pedales, *basi repentes*, demum suberecti, geniculati, (geniculis paululum incrassatis,) laeves, rubicundi.

FOLIA *lineari-lanceolata*, pene aevia, superne glabra.

STIPULÆ vaginantes, ciliatæ.

SPICÆ tenues, parum nutantes, e singulis geniculis prodeuntes.

CALYX: PERIANTHIUM quinquepartitum, persistens, coloratum, laciniis obtusis concavis, *fig. 1.*

COROLLA nulla.

STAMINA: FILAMENTA sex; ANTHERÆ biloculares, albae, intra Corollam.

PISTILLUM: GERME ovatum aut triangulare; STYLUS filiformis, apice bifidus aut trifidus; STIGMATA duo aut tria rotunda, reflexa, *fig. 2, 3.*

SEMEN aut ovato-acutum aut triangulare, cauleanum; magnitudinis fere et formæ feminis Polygoni Persicariæ, *fig. 4, 5.*

N. B. Omnes partes fructificationis lente augentur.

* ROOT annual, and fibrous.

STALKS several, about nine inches or a foot high, *creeping at bottom*, then becoming nearly upright, jointed, (the joints somewhat thickened,) smooth, of a reddish colour.

LEAVES *between linear and lanceolate*, scarcely any appearance of veins, on their upper surface smooth. STIPULÆ forming sheaths round the joints, and ciliated.

SPIKES slender and a little drooping, proceeding from each joint of the stalk.

CALYX: a PERIANTHIUM divided into five segments, which are obtuse and hollow, *fig. 1.*

COROLLA wanting.

STAMINA six FILAMENTS; ANTHERÆ bilocular, and white, within the Corolla.

PISTILLUM: GERME oval or triangular; STYLE filiform, at top bifid or trifid; STIGMATA two or three, round and turned back, *fig. 2, 3.*

SEEDS oval or triangular, of a chestnut colour, nearly of the same size and shape as the Polygonum Persicariæ, *fig. 4, 5.*

* N. B. All the parts of the fructification are magnified.

If the opportunity of seeing this plant growing wild had ever occurred to the celebrated Swedish Botanist, he would doubtless have considered it as a distinct species; at present he has placed it in the last edition of his works, the *Systema Vegetabilium*, as a variety of the *Polygonum Persicaria*, probably misled by dried specimens of the plant: those who trust to such are exceeding liable to deceive both themselves and others, particularly in plants whose parts of fructification (from which it is sometimes necessary to draw specific differences) are very minute—those in the living plants are with difficulty enough distinguished, and in dried specimens not to be investigated.

Whoever has observed the appearance which the *Polygonum minus* and *Persicaria* usually put on, must have been struck with the great dissimilarity of the two in their general habits; and if they have taken the pains to examine the parts of fructification, they will, I am persuaded, be convinced that both Mr. RAY and HUDSON are justifiable in making them distinct species.

It differs from the *Polygonum Persicaria* in its size, growth of its stalk, shape of its leaves, form of its spikes, and division of its Pistillum. In height it seldom exceeds a foot, whereas the *Persicaria* often occurs a yard high; the stalk of this species creeps at bottom, in the *Persicaria* it never does: it is true in the *Persicaria*, and most of the *Polygonums*, a number of little roots push themselves out at the joints, which are next the ground; but in this species the stalk at bottom is absolutely procumbent, whilst in the *Persicaria* it is always upright; the leaves are much narrower, approaching rather to linear than lanceolate, and on their upper surface have much less appearance of veins, than in the *Persicaria*; the spikes, instead of being oval or nearly round, and upright, as in the *Persicaria*, are slender and a little drooping: the Pistillum, which is a part of very great consequence in determining many of the species and varieties of this genus, is slightly divided at top only; while that of the *Persicaria* is divided half way down; hence as I have called this species *submonogynum*, I have called this *submonogynum*.

Hitherto I have met with this plant growing wild nowhere but in *Tat. H. field's, H. field's*, where it makes ample amends for its scarcity elsewhere, being found in the greatest abundance in the watery parts of those fields, along with the *Sisymbrium sylvestre* in the month of September, when it is in full bloom.

At present it does not appear that it has any thing more than its scarcity to recommend it to our notice.

WILSON'S HISTORY OF

THE UNITED STATES OF AMERICA

BY

JOHN WILSON

NEW YORK

Published by the

Author

NEW YORK

1884

Copyright, 1884, by

Author

BUTOMUS UMBELLATUS. FLOWERING RUSH, OR WATER GLADIOLE.

BUTOMUS *Linnaei Gen. Pl. ENNEANDRIA HEXAGYNIA.*

Raii Gen. 17. HERBÆ MULTISILIQUÆ seu CORNICULATÆ.

BUTOMUS umbellatus. *Lin. Spec. plant. 532.*

JUNCUS floridus major. *Bauhin. pin. p. 12.*


GLADIOLUS palustris Cordi *Gerard. emac. 29.*


Raii Synopsi ed. 3. 273. Hudson. Fl. Angl. 152. Scopoli Fl. Carn. ed. 2. p. 283. Halleri hist. pl. Helv. Vol. 2. 81.

RADIX perennis, alba, tuberculosa, transversa, edulis : ex inferiore parte radículas prælongas dimittens.	ROOT perennial, white, knobby, transverse, eatable ? from its under side sending down a great number of very long fibres.
SCAPUS pedalis ad orgyalem, teres, glaber.	STALK round, smooth, from one to five or six feet high, according to its place of growth.
FOLIA triquetra, spongiosa, fig. 1. scapo breviora ; ad basin spathacea, apicibus compressis, tortuosis.	LEAVES triangular, spongy, fig. 1. shorter than the stalk, at bottom sheathy, at top flat, and twist- ed.
FLORES in UMBELLA, ad triginta ; pedunculi digitales, c vaginis membranaceis prodeuntibus.	FLOWERS numerous, to thirty, each on a single pe- duncle of about a finger's length, forming an UMBELL, surrounded at bottom by withered membranous sheaths.
CALYX. INVOLUCRUM triphyllum, foliolis lanceolatis, marcescentibus.	CALYX. An INVOLUCRUM of three leaves, spear shaped and withered.
COROLLA. PETALA sex inæqualia, subrotunda, concava, rosea, fig. 2, alternis minoribus, acutioribus. fig. 3.	COROLLA. Composed of six PETALS, which are round- ish, concave, and most commonly of a bright red ; fig. 2. the three exterior smaller, and more pointed, fig. 3.
STAMINA. FILAMENTA novem, subulata, fig. 4, 5. ANTHERÆ insidentes, dum pollinem involvunt oblongæ, rubræ, quadrifidatæ, mucrone brevi terminatæ, fig. 6, 7, emissio polline subcordatæ, compressæ, bilamellosæ, fig. 4. POLLEN flavissi- mum.	STAMINA. Nine FILAMENTS tapering, fig. 4. 5. the ANTHERÆ sitting on the filaments, before the shedding of the Pollen, oblong, reddish, having four grooves, and terminated by a short point, fig. 6. 7. appearing afterwards somewhat heart shaped, flat, and as if composed of two lamellæ, fig. 4. the POLLEN is of a bright yellow colour.
PISTILLUM. GERMEN subtriangulare, latere exteriore latiore, convexo, fig. 9. 10. STYLI sex subulati ; fig. 8. STIGMA canaliculatum.	PISTILLUM. the GERMEN nearly triangular, the outer side broad and roundish, fig. 9. 10. six STYLES, tapering, the STIGMA has a small channel in it, which afterwards spreads into two lips, fig. 11. 12.
PERICARPium. CAPSULÆ sex, oblongæ, attenuatæ, erectæ, univalves, apice bilabiata, introrsum dehiscentes, fig. 11. 12.	SEED-VESSEL. Six CAPSULES, oblong, tapering, up- right, of one valve, opening inwards, fig. 11. 12.
SEMINA plurima, minuta, oblonga, fusca, fig. 13.	SEEDS numerous, small, oblong, brown, fig. 13.

WE find this stately Plant in, and by the sides of our watery ditches flowering from July to September. A few years since it was found growing in *St. George's-Fields*, but the improvements making in that, and other parts adjacent to *London*, now oblige us to go farther in search of this and many other curious Plants. About the *Island of St. Helena*, near *Deptford*, and the *Marishes by Blackwall*, it is found in great abundance, although very scarce in many other parts of *Great-Britain*. Fish-ponds, or other pieces of water, would derive great beauty from the introduction of this elegant native of our Isle, the handsome appearance of which did not escape our countryman, old *GERARD*, who describes it thus : " This Water Gladiole, or Grassy-Rush, of all others, is the fairest and most pleasant to behold, and serveth very well for the decking and trimming up of houses, because of the beaute and braverie thereof." That accurate observer *RAY* describes its nine STAMINA, although in his time they were not viewed in that consequential light which they are in our present Systems of Botany. It is the only Plant of the class *ENNEANDRIA* which grows wild in this kingdom. If vegetables were classed according to their natural affinities, this would rank among the lilies. Cattle do not eat it. It is so hardy as to bear the cold of *Lapland*.







SAXIFRAGA GRANULATA.

WHITE SAXIFRAGE.

SAXIFRAGA *Linnæi* Gen. Pl. DECANDRIA DIGYNIA.

Calyx quinquepartitus. *Corolla* pentapetala. *Capsula* birostris, unilocularis, polysperma.

Raii Syn. HERBÆ PENTAPETALÆ VASCULIFERÆ.

SAXIFRAGA *granulata* foliis caulinis reniformibus lobatis, caule ramofo, radice granulata. *Linn. Syst.*

Vegetab. p. 344. Fl. Suecic. n. 372.

SAXIFRAGA foliis radicalibus reniformibus, obtuse dentatis, caulinis palmatis. *Holboell. hist. belg. n. 976.*

SAXIFRAGA rotundifolia alba. *Bauhin Pin. 309.*

SAXIFRAGA alba. *Gerard emac. 841.*

SAXIFRAGA alba vulgaris. *Parkinson 424. Raii Syn. 354. Hudson Fl. Angl. p. 159. Oeder. Flor.*

Dan. 514.

RADIX. Fibris hujus radices glomeratim adnascuntur plurimi bulbilli, extus rubefcentes aut flavescerentes, intus albi, saporis primum adstringentis, postea amari et ingrati.

CAULIS plerumque simplex, pedalis, subramosus, teres, hirsutus, præsertim ad basin, parum foliosus.

FOLIJA radicalia petiolis longis, hirsutis, basi latis infidentia, reniformia, hirsutula, lobata, lobis obtusis; caulina sicut adscendunt petiolis brevioribus gaudent donec tandem sessilia fiunt, lobis foliorum acutiora evadunt, apicibus rufescentibus.

CALYX: PERIANTHIUM quinquepartitum, hirsutululum, subviscidum, lacinias ovato-acutis apice rufis, fig. 1.

COROLLA: PETALA quinque alba, patentia, apice rotundata, basi angustiora et venis flavescens notata, fig. 2.

STAMINA: FILAMENTA decem subulata; ANTHERÆ ovatæ, compressæ, infidentes, flavæ, biloculares, quorum quinque Pollen primum emittunt, hinc longiores, fig. 3, 4.

PISTILLUM: GERMEN subrotundum, inferum, glandulâ saturate viridi cinctum, fig. 7; STYLI duo Staminiibus breviores, incurvati, fig. 5; STIGMA concavum, fig. 5, demum expandens, fig. 6.

PERICARPIUM: CAPSULA subovata, birostris, bilocularis, pallide fusca, fig. 8.

SEMINA numerosa, minutissima, nigra, fig. 9.

ROOT. To the fibres of the root of this plant, adhere in clusters a number of small bulbs, externally red or yellowish, internally white, of a taste at first astringent, afterwards bitter and disagreeable.

STALK generally simple, about a foot high, a little branched, round, hirsute particularly at bottom, furnished with but few leaves.

LEAVES which grow next the root placed on long hairy foot-stalks with a broad base, kidney-shaped, slightly hairy, divided into obtuse lobes, those of the stalk, as they ascend, are furnished with shorter foot-stalks, till they gradually become sessile, the lobes more acute, and the tips of a reddish colour.

CALYX: a PERIANTHIUM divided into five segments hirsute and somewhat viscid, the laciniae of an oval pointed shape, and red at the tips, fig. 1.

COROLLA: five PETALS, white, spreading, round at top, at bottom narrower, and striped with yellowish veins, fig. 2.

STAMINA: ten FILAMENTS tapering; ANTHERÆ oval, flat, sitting on the Filaments, yellow, bilocular, five of them shed the Pollen first, hence they become longer than the others, fig. 3, 4.

PISTILLUM: GERMEN roundish, placed below the Calyx, surrounded by a gland of a deep green colour, fig. 7; STYLES, two, shorter than the Stamina, bending inward, fig. 5; STIGMA hollow, fig. 5, finally expanding, fig. 6.

SEED-VESSEL: a CAPSULE of a shape somewhat oval, and pale brown colour, having two beaks or horns, and two cavities, fig. 8.

SEEDS numerous, very minute, and black. fig. 9.

THE Root of this species of *Saxifrage*, by means of which it is chiefly propagated, affords the young Botanist a very good example of the *Radix granulata*, being composed of a number of little grains or bulbs, connected together in clusters by the fibres; some of these bulbs are solid and entire, not unaptly resembling in shape the bulbs of Onions; others spread open at top, and seem to be composed of a number of squamulæ or lesser bulbs, these are often of a bright red colour: the upper part of the stalk, the foot-stalks of the flowers, and calyx, are covered with a kind of hairs, which terminate in a viscid globule, and which seem to accompany most of the plants of this Genus. The two Styles, which at first are short, with a hollow Stigma, fig. 5, quickly grow much longer; the Stigmata spread open, so that they resemble in some degree a pair of tea-tongs, fig. 6.


This plant does not occur so frequently with us as many others: according to Mr. HUDSON, it is common about *Wandsworth*; I have frequently gathered it in the fields about *Peckham*, and lately have found it in great abundance much nearer town, viz. in the fields called *Lack-fields*, on the right hand side of *Kent-street Road*, at the back of, and contiguous to Mr. DRIVER'S Nursery Gardens: it delights to grow in dry pastures which have a gravelly bottom; flowers in May, and produces its seeds in the month following. When double, it serves, with many other British plants, to ornament the gardens of the curious.

Like many other plants, this seems to owe what little importance it has in medicine to the doctrine of signatures, which has most unphilosophically introduced a number of plants into our *Materia Medica*. As the root bore so great a resemblance to little stones, it was concluded it must be efficacious in the stone and gravel, for which diseases it has been recommended, but there are no accounts of its success to be depended on. If it does possess any medical virtue, it should appear from the taste of the root to be that of an astringent.



Ant. et pist.

Saxifraga granulata.



SEDUM ALBUM. WHITE-FLOWERED STONECROP.

SEDUM *Linnei Gen. Pl.* DECANDRIA PENTAGYNIA.

Cal. 5 fidus. *Cor.* 5-petala. *Squamæ* nectariferæ 5, ad basin germinis. *Caps.* 5.

Raii Syn. Gen. 17. HERBÆ MULTISILIQUÆ SEU CORNICULATÆ.

SEDUM *album* foliis oblongis obtusis teretiusculis sessilibus patentibus, cyma ramosa. *Lim. Syst. Vegetab.* p. 359. *Sp. Pl.* p. 619. *Fl. Suecic.* 153.

SEDUM caule glabro, foliis teretibus; umbellis ramosis; floribus petiolatis. *Haller hist. helv.* n. 959.

SEDUM *album.* *Scopoli Fl. Carn.* p. 324.

SEDUM minus teretifolium album. *Baubin.* p. 283.

SEDUM minus officinarum. *Gerard emac.* 512.

VERMICULARIS five crassula minor vulgaris. *Parkinson* 734. *Raii Syn.* 271. *Hudson Fl. Angl.* p.

171. *Oeder. Fl. Dan. Icon.* 66.

RADIX perennis, fibrosa.

CAULES flexuose super muros repent, dein eriguntur, triunciales circiter, foliosi, rubri.

FOLIA sessilia, oblonga et fere cylindracea, obtusa, non admodum conferta, patentia, carnosâ, glabra, sæpius rubicunda.

INFLORESCENTIA : Flores petiolati, in CYMAM ramosam confertam dispositi.

CALYX : PERIANTHIUM pentaphyllum, foliolis brevibus, obtusis, fig. 1.

COROLLA : PETALA quinque alba, acuminata, lineâ longitudinali rubra sæpius notata, fig. 2.

NECTARIUM glandula minima squamiformis ad basin singuli Germinis. fig. 6.

STAMINA : FILAMENTA decem alba, fig. 2, 3; ANTHERÆ rubræ.

PISTILLUM : GERMINA quinque, in STYLOS totidem acuminatos definitia; STIGMATA simplicia, fig. 4, 5.

PERICARPIUM : CAPSULÆ quinque minimæ acuminatæ introrsum dehiscentes, fig. 7.

SEMINA parva, oblonga, fig. 8.

ROOT perennial and fibrous.

STALKS creep on the walls in a crooked form, then grow upright, about three inches high, leafy, and red.

LEAVES sessile, oblong and almost cylindrical, obtuse, but thinly placed on the stalk, spreading, fleshy, smooth, and generally of a reddish colour.

INFLORESCENCE : Flowers standing on foot-stalks, and disposed in a thick branched CYMA.

CALYX : a PERIANTHIUM of five leaves, which are short and obtuse, fig. 1.

COROLLA : five white PETALS, acuminated and generally marked with a longitudinal red streak, fig. 2.

NECTARY a very minute squamiform gland at the base of each of the Germina, fig. 6.

STAMINA : ten white FILAMENTS, fig. 2, 3; ANTHERÆ deep red.

PISTILLUM : five GERMINA, terminating in so many acuminated STYLES; the STIGMATA simple, fig. 4, 5.

SEED-VESSEL : five small acuminated CAPSULES, opening inwardly, fig. 7.

SEEDS small and oblong, fig. 8.

The *Sedum album* may be considered with us as rather a scarce plant; it is found here and there on the Walls about Town, particularly on the Chapel-wall in *Kentish-Town*, where it has grown for many years; also upon a Wall on the left-hand side leading from *Bromley* to *Bromley-Hall*, in *Middlesex*. It has been thought to possess sufficient beauty to recommend it as a garden plant, and is accordingly, with very little trouble, cultivated in many of the gardens of the curious, nothing more being necessary than placing it in a pot filled with gravel or mould: in such a situation it will grow, flourish, and propagate itself very fast.


It has been called *album* from the colour of its flowers, which generally however have a tinge of red in them. It flowers in July. The round and oblong shape of its leaves readily distinguishes it from our other *Stonecrops*.

HALLER informs us that it possesses all the virtues of the large *Housleek*, and that he has used the juice of it in uterine hæmorrhages, but does not inform us with what success. By way of cataplasm it is applied to the piles when in a painful state, and is said to have sometimes been made the same use of in cancers with success. By some it is eaten as a pickle.



a b c d e f g h i j k l m n o p q r s t u v w x y z

Sedum album.



SEDUM ACRE. COMMON YELLOW STONECROP, OR WALL-PEPPER.

SEDUM. *Linnaei Gen. Pl.* DECANDRIA PENTAGYNIA.

Raii Synopsis Gen. 17. HERBÆ MULTISILIQUÆ SEU CORNICULATÆ.

SEDUM *acre* foliis subovatis, adnato-fessilibus, gibbis, erectiusculis, alternis; cyma trifida. *Lin. Syst. Vegetab.* p. 359. *Fl. Suecic.* p. 153.

SEDUM foliis conicis confertis, caulibus ramosis, fummis trifidis. *Haller. bist. v. 1. n. 966.*

SEMPERVIVUM minus vermiculatum *acre*. *Baubin. pin.* 283.

VERMICULARIS seu Illecebra minor *acris*. *Ger. emac.* 517.

ILLECEBRA minor seu Sedum tertium Dioscoridis. *Parkinson 735. Raii Synop.* 270. *Hudson. Fl. Angl.* p. 171.

RADIX perennis, fibrosa.

CAULES numerosi, caespitosi, ramosissimi, palmares, ad basin repentes, dein erecti, teretes, foliosissimi.

FOLIA alterna, conferta, imbricata, subrecta, adnato-fessilia, ovata, obtusa, breviter, carnosa, margine paululum compressa, glabra, sapore acri. *fig. 1.*

FLORES sessiles, lutei, in Cymas subtrifidas dispositi.

CALYX: PERIANTHIUM quinquepartitum, persistens, lacinis crassis obtusiusculis, *fig. 2.*

COROLLA: PETALA quinque lanceolato-acuminata, plana, patentia, Calyce duplo longiora, *fig. 3.*

NECTARIUM: Squamula minima, alba, ad basin, singuli germinis extrorsum posita, *fig. 7.*

STAMINA: FILAMENTA decem subulata, longitudine Corollæ. ANTHERÆ flavæ, *fig. 4.*

PISTILLUM: GERMINA quinque oblonga, flava, in Stylis acuminatis definitia. STIGMATA simplicia, *fig. 6.*

PERICARPIUM: CAPSULÆ quinque patentia, acuminata, compressæ, longitudinaliter futura introrsum dehiscences, *fig. 8.*

SEMINA minima, ovata, rufa, *fig. 9.*

ROOT perennial, and fibrous.

STALKS numerous, growing in tufts, very much branched, three inches high, creeping at their base, but afterwards growing upright, round, and very leafy.

LEAVES alternate, growing very thick together, and laying one over another, nearly upright, growing to the stalk, oval, blunt, short, fleshy, flattened a little at the edges, smooth, and of a very biting taste, *fig. 1.*

FLOWERS sessile, yellow, growing in Cymæ somewhat trifid.

CALYX: a PERIANTHIUM divided into five segments, and continuing, the segments thick and bluntish, *fig. 2.*

COROLLA: composed of five long-pointed PETALS which are flat, spreading, and twice the length of the Calyx, *fig. 3.*

NECTARY: a very minute scale or gland placed externally at the bottom of each Germen, *fig. 7.*

STAMINA: ten FILAMENTS, tapering, the length of the Corolla, the ANTHERÆ yellow, *fig. 4.*

PISTILLUM: five GERMINA, oblong, yellow, terminating in five long-pointed STYLES. The STIGMATA simple, *fig. 6.*

SEED-VESSEL: five CAPSULES, spreading, long-pointed, flat, opening internally by a longitudinal suture, *fig. 8.*

SEEDS very minute, oval, and reddish brown, *fig. 9.*

According to the account which some medical Writers give of this Plant it appears to possess considerable virtues, while others, from the durability of its acrimony, and the violence of its operation, have thought it scarce safe to be administered. Chewed in the mouth it has a very hot and biting taste, whence its name of *Wall-Pepper*. Applied to the skin it excoriates and exulcerates it, taken internally it proves emetic and diuretic.


The Diseases in which it has been chiefly recommended are the Scurvy and Dropsy, in both of which, according to Linnaeus, it is an excellent remedy; and some instances are brought of the efficacy of its juice in Cancers, but these perhaps, stand in need of farther confirmation.

It grows very common on Houses, Walls, and gravelly Banks, and flowers in June; it continues but a short time in blossom, but while it lasts its lively yellow colour gives a very pretty appearance to those Houses and Walls which are covered with it.



Sedum acre

1852





LYCHNIS FLOS CUCULI. MEADOW LYCHNIS.

LYCHNIS *Linnaei* Gen. Pl. DECANDRIA PENTAGYNIA.

Raii Synopsi Gen. 24. HERBÆ PENTAPETALÆ VASCULIFERÆ.

LYCHNIS *Flos Cuculi* petalis quadrifidis fructu subrotundo. *Lin. Syst. Vegetab.* p. 361. *Sp. Pl.* 625.

LYCHNIS petalis quadrifidis. *Haller. hist. v. 1. n. 921.*

CARYOPHYLLUS pratensis, laciniato flore simplici, five Flos cuculi. *Baubin. pin.* 218.

LYCHNIS plumaria sylvestris simplex. *Parkinson. parad.* 253.

ARMERIUS pratensis mas et femina. *Gerard. Emac.* 600.

Raii Synop. ed. 3. 338. *Hudson. Fl. Angl.* 174. *Oeder. Flor. Dan. tab.* 590. *Scopoli. Fl.*

Carniol. ed. 2. p. 311.

RADIX perennis, fibrosa, ex albedo fusca, saporis sub-acris.

CAULIS pedalis ad tripedalem, erectus, fulcato-angulatus, articulatus, geniculi tumidi, scabriusculus, purpurascens.

FOLIA Caulis, opposita, connata, lanceolata, carinata, suberecta, laviâ.

PEDUNCULI oppositi, plerumque unico intermedio.

CALYX: PERIANTHIUM monophyllum, tubulatum quinquedentatum, decangulatum, purpureum, persistens. *fig. 1.*

COROLLA PETALA quinque, unguis longitudine calycis, *fig. 2.* LIMBUS quadrifidus, laciniis exterioribus brevioribus, et angustioribus, *fig. 4.* ad basin limbi laminæ duæ erectæ acutæ. *fig. 3.*

STAMINA: FILAMENTA decem, subulata, quorum quinque breviora, *fig. 5,* brevioribus ungui petalorum affixis. *fig. 6.* ANTHERÆ oblongæ, biloculares, *fig. 7.* incumbentes, purpurascens.

PISTILLUM GERMEN subovatum, *fig. 8.* STYLI quinque subulati, subincurvati, *fig. 10.* STIGMATA simplicia. *fig. 10.*

PERICARPIUM CAPSULA ovata, unilocularis, ore quinquedentato, dentibus reflexis. *fig. 9.*

SEMINA numerosa, subcompressa, scabriuscula, ex cinereo-fusca. *fig. 11. 12.*

ROOT perennial, fibrous, of a brownish white colour, and somewhat biting taste.

STALK from one to three feet high, upright, somewhat angular and grooved, jointed, the joints swelled, roughish, and of a purplish colour.

LEAVES of the Stalk opposite, connate, lanceolate, the midrib prominent underneath, upright and smooth.

PEDUNCLES opposite, one generally intermediate.

CALYX a PERIANTHIUM of one leaf, tubular, quinque-dentate, having ten angles, or ridges, and of a deep purple colour.

COROLLA: five petals, the claw the length of the Calyx, *fig. 2.* the LIMB divided into four laciniæ, the exterior shortest and narrowest, *fig. 4.* at the bottom of the limb are placed two small upright laminæ, *fig. 3.*

STAMINA: ten FILAMENTS, tapering, five long and five short, *fig. 5.* the shorter filaments affixed to the claw of each petal, *fig. 6.* the ANTHERÆ oblong, bilocular, *fig. 7.* laying across the filaments, and of a purplish hue.

PISTILLUM: the GERMEN somewhat oval, *fig. 8.* five STYLES tapering and bending a little inward, *fig. 10.* STIGMATA simple. *fig. 10.*

SEED-VESSEL: a CAPSULE, oval, of one cavity, the mouth having five teeth which turn back. *fig. 9.*


SEEDS numerous, flattish, rough, and of a brown ash colour. *fig. 11. 12.*

A variety of names hath been given to this Plant, as Meadow Pink, Cuckow Flower, Wild Williams, Ragged Robin, &c. Meadow Lychnis however seems to us the most eligible. It abounds in moist Meadows, where it flowers in May and June, and is included amongst the great number of which our Meadow hay is compounded. Goats, Sheep, and Horses are said to feed on it. The use to which it is applied, seems to be chiefly ornamental; the beauty of its flowers justly entitles it (with many other neglected British Plants) to a place in the Gardens of the curious: where it is frequently found with a double flower, making a beautiful appearance, and requiring little more care in its culture, than to be placed in a moist situation: It may be propagated either by seeds or slips; the seeds may be found ripe in the latter end of June, by the sides of ditches, where the Mower's Scythe has not reached them. We sometimes find the Meadow Lychnis growing wild with a double flower, and sometimes with a white one; but this is altogether accidental.

The agreement between the blowing of flowers, and the periodical return of birds of passage, has been attended to from the earliest ages: Before the return of the seasons was exactly ascertained by Astronomy, these observations were of great consequence in pointing out stated times for the purposes of Agriculture; and still, in many a Cottage, the birds of passage and their corresponding flowers assist in regulating

"The short, and simple Annals of the Poor."

For this reason, no doubt, we have several other plants that, in different places, go by the name of Cuckow Flower. *Gerard* says, *Cardamine pratensis* (Common Ladies Smock) is the true Cuckow Flower. *Shakespeare's* Cuckow Buds are of "yellow hue." By some the Orchis, Arum, and Wood-forrel are all called after the Cuckow.





Cerastium aquaticum.

CERASTIUM AQUATICUM. MARSH CERASTIUM OR MOUSE-EAR CHICKWEED.

CERASTIUM *Linnei Gen. Pl.* DECANDRIA PENTAGYNIA.

Raii Synop. Gen: 24 HERBÆ PENTAPETALÆ VASCULIFERÆ.

CERASTIUM *aquaticum* foliis cordatis, sessilibus, floribus solitariis, fructibus pendulis. *Linnei Syst.*

Vegetab. p. 363. *Fl. Suecic. p.* 157.

ALSINE foliis ovato-cordatis, imis petiolatis, tubis quinis. *Haller. hyst. n.* 885.

STELLARIA *aquatica. Scopoli Fl. Carniol. p.* 320.

ALSINE *aquatica major. Baubin. pin.* 254.

ALSINE *major. Gerard emac.* 611. *maxima Parkinson 759. Raii Syn. p.* 347. *Hudson Fl. Angl. p.* 177.

RADIX perennis, fibrosa, repens.

ROOT perennial, fibrous, and creeping.

CAULES bipedales, debiles, pene teretes, teneri, filiosi, hirsuti, ramosi, rami alterni.

STALKS about two feet in length, weak, almost round, tender, stringy, hirsute, and branched, the branches alternate.

FOLIA Caulis sessilia, amplexicaulia, cordato-acuminata, margine in superioribus presertim undulata, laevia, subvifida; ramorum magis undulata, petiolata.

LEAVES of the Stalk sessile, embracing the Stalk, somewhat heart shaped and acuminate, the edge particularly in the upper ones waved, smooth, and somewhat viscid; those of the branches more waved with short footstalks.

PEDUNCULI alterni, e dichotomia caulis, uniflori, post florescentiam penduli.

FOOTSTALKS alternate, from the forking of the Stalk, uniflorous, after the blossom is gone off pendulous.

CALYX: PERIANTHIUM pentaphyllum, persistens, foliolis lanceolatis, concavis, fubcarinatis, apice obtusiusculis, hirsutis, margine membranaceis, petalis paulo brevioribus. *fig. 1.*

CALYX: a PERIANTHIUM of five leaves, persisting, the leaves lanceolate, concave, slightly keel-shaped, bluntish at top, hirsute, at the edge membranous, and a little shorter than the Petals, *fig. 1.*

COROLLA: PETALA quinque alba, patentia, bipartita, lacinis oblongis, nervosis, divaricantes, *fig. 2. 3.*

COROLLA: five PETALS white, spreading, divided almost to the bottom, the laciniae or segments oblong, nervous, and divaricating, *fig. 2. 3.*

STAMINA: FILAMENTA decem, subulata, alba, receptaculo inserta, ad basin et inter petala alterne locata, *fig. 4.* quæ inter petala locantur paulo longiora sunt et glandula ad basin instruuntur *fig. 5.* ANTHERÆ insidentes, biloculares, albae, *fig. 4.*

STAMINA: ten FILAMENTS, tapering, white, fixed to the receptacle, placed alternately, one at the base and one betwixt each petal, *fig. 4;* those placed between the petals are a little longer than the others, and furnished at bottom with a gland, *fig. 5.* ANTHERÆ white and bilocular, *fig. 4.*

PISTILLUM: GERMEN subrotundum, apice fuscum, STYLII quinque albi, filiformes, longitudine Germinis. STIGMATA simplicia, *fig. 6.*

PISTILLUM: GERMEN roundish, at top grooved, five STYLES thread-shaped, white, the length of the Germen. STIGMATA simple, *fig. 6.*

PERICARPIUM: CAPSULA ovata, obsoleta pentagona, ore quinqueedentata. *fig. 7.*


SEED-VESSEL: an oval CAPSULE, slightly pentagonal, the mouth quinqueedentate.

SEMINA rufa, subreniformia, scabra, 60 numeravi, *fig. 8. 9.*

SEEDS reddish brown, rough, about 60 in each capsule, *fig. 8. 9.*

SOME of our modern and most celebrated systematic Botanists seem very much divided with respect to the Genus to which this Plant should belong. HALLER makes it an *Alfine* or *Chickweed*; SCOPOLI a *Stellaria*, and LINNEUS a *Cerastium*. We shall not pretend to decide who is most in the right, but only observe that its general habit or appearance, and the form of its seeds, might easily induce HALLER to consider it as an *Alfine*; the shape of its petals, with the structure of its seeds, would justify SCOPOLI in calling it a *Stellaria*, while the number of its styles might lead LINNEUS with propriety to place it among the *Cerastium*. To us it appears to have the greatest natural affinity with the *Alfine media* or common *Chickweed*; it is true LINNEUS ranks that plant among those which have five Stamina, yet it is frequently observed to have more, and the structure of the flower evidently shows it to be formed for having ten, and those flowers which have not that number may be considered as imperfect. The seeds of these two plants are so similar as scarcely to be distinguished from each other, and their stalks are procumbent, tender, brittle, and stringy, indeed they frequently so much resemble one another, as to oblige the young Botanist to have recourse to the very different size of their flowers in order to discriminate them.

This Plant grows in moist places, on the banks of rivers and by streams of water, it flowers in July and August. SCOPOLI asserts that the plants of this kind afford excellent food for Kine.



EUPHORBIA PEPLUS. SMALL GARDEN SPURGE.

EUPHORBIA *Linnaei Gen. Pl. DODECANDRIA TRIGYNIA.*

Raii Syn. Gen. 22. HERBE VASCULIFERE FLORE TETRAPETALO ANOMALÆ.

EUPHORBIA (*Peplus*) umbella trifida, dichotoma, involuclis ovatis, foliis integerrimis obovatis petiolatis. *Lin. Syl. Vegetab. p. 375. Fl. Succ. p. 163.*

TITHYMALUS foliis rotundis, stipulis floralibus cordatis, obtusis, petalis argute corniculatis. *Haller. hist. vol. 2. p. 9. n. 1049.*

PEPLUS five Efula rotunda. *Bauhin pin. 292. Parkinson. Gerard. emac. 503.*

TITHYMALUS parvus annuus, foliis subrotundis non crenatis, Peplus dictus. *Raii Syn. p. 313. n. 9: Petty Spurge. Hudson Fl. Angl. p. 182.*

RADIX annua, lignosa, simplex, fibrosa, albida.

CAULIS, suberectus, dodrantalis, teres, glaber, ramosus, basi durior, tenuior, scaber, foliosus, laeviss.

RAMI pauci, sparsi, inferioribus longioribus oppositis.

UMBELLA trifida, dichotoma.

FOLIA obovata, petiolata, integerrima, sparsa, obtusiuscula, inferioribus subrotundis.

STIPULÆ *umbellæ* tres, ovato-acutæ, petiolis brevibus insidentes, *umbellulæ* alterne oppositæ, sessiles, cordato-ovate, inaequales, integerrimæ, basi quæ tendit germen quasi cavatæ.

CALYX ventricosus, persistens, fig. 1.

COROLLA nulla.

NECTARIA quatuor *bicorniculata*, fig. 2.

STAMINA plerumque duo, aut tria, visibilia, exserta:

ANTHÆ didymæ, subrotundæ, fig. 3.

PISTILLUM: GERMEN pedunculatum, nutans, triangulare, angulis longitudinaliter fuscatis, fig. 4, 6: STIGMATA tria, apice bifida, fig. 5.

PERICARPIUM: CAPSULA trilocca, trilocularis, trivalvis, valvulis levibus, et dum adhuc virides diffidentibus, fig. 6.

SEMEN unicum in singulo loculamento, ovatum, canum, *obtusiusculum*, appendiculatum, fig. 7.

N. B. Omnes partes fructificationis lente augentur.

ROOT annual, woody, simple, fibrous and whitish.

STALK generally upright, about nine inches high, round, smooth, and branched; at bottom harder, more slender, and of a reddish colour, leafy and milky.

BRANCHES few, not growing in any regular order, the lower ones longest and opposite.

UMBEL first trifid, then dichotomous.

LEAVES somewhat oval, but narrowest towards the base, having foot-stalks, entire at the edges, placed in no regular order, somewhat blunt, the lowermost leaves almost round.

STIPULÆ of the large *umbel* three in number, oval and pointed, placed on very short foot-stalks: of the small *umbel* alternately opposite, sessile, of an heart-shaped-oval form, unequal, and entire, at bottom on that side to which the Germen tends as if cut away.

CALYX bellying out and continuing, fig. 1.

COROLLA wanting.

NECTARIES four, each having two little horns, fig. 2.

STAMINA seldom more than two or three, which are visible, and placed without the Calyx: ANTHERÆ two on each filament joined together, of a roundish figure, fig. 3.

PISTILLUM: GERMEN placed on a foot-stalk, hanging down, triangular, the angles longitudinally grooved, fig. 4, 6: STIGMATA three, bifid at top, fig. 5.

SEED-VESSEL: a CAPSULE of three cavities, and three valves, the valves protuberant, smooth, and splitting with a kind of elasticity even while they are of a green colour, fig. 6.

SEED one in each cavity, oval, grey, with numerous depressions on its surface, and a little white button at one end, fig. 7.

N. B. All the parts of fructification are magnified.

MANY of the Spurges considerably resemble one another, and two of them that have this affinity, grow frequently together in Gardens, viz. the present Spurge, *Euphorbia Peplus*, and the Sun Spurge, *Euphorbia Helioscopia*; they may be distinguished however by the slightest attention. In the *Helioscopia* the leaves are *notched* or *serrated* at the edges, in the *Peplus* they are *entire*, in the *Helioscopia* the Petals or rather Nectaria are *round* and *entire*, in the *Peplus* each is furnished with *two little horns*, fig. 2; there are other marks of distinction but these are the most striking.

This species grows in Gardens and other cultivated ground, and flowers in Autumn.

The milky fluid which it abounds with, is by some applied to Warts, which it is said to destroy.

Most if not all the plants of this Genus contain in them this milky and gummy substance, which to the taste is exceedingly acid; and this lactitious property, joined to the peculiarity of its parts of fructification, point out almost at first sight this natural family of plants. But the botanic Student who would investigate this species according to the principles of the Linnaean System, not having these characteristics to assist him, finds a considerable difficulty in learning even the *Class* to which it belongs, nor is it possible for him to ascertain the *Class* by an examination of this or scarce any other English Spurge: the *Stamina* in the first place are very minute, it is seldom that more than two or three protrude beyond the Calyx, all the rest lie concealed within it, they seldom amount to twelve in number, and even if they did amount to that exact number, their minuteness and the milky juice which flows from the dissection, render the enumeration of them scarce practicable. The Student may however in a great degree surmount this difficulty, by an examination of some plant of this genus, which is larger in every respect, and the *Euphorbia Lathyris* improperly called the *Coper Tree*, (which is cultivated in many Gardens) will afford him a very good example, and tend to give him a clear idea of the flower and fruit of this singular genus of plants.


I would not be thought on account of this difficulty to inveigh against LINNÆUS's System, being sensible that difficulties occur, and must occur in all botanic arrangements, and instead of selecting faults inseparable from every mode of classification, (which seems to have been a favourite amusement of some Authors, and forms indeed the greatest part of their writings) I would use every endeavour to make it more perfect.

It is too much the fashion now, as well as formerly, for every Botanist as soon as he thinks he has some pretensions to eminence, to set about the arduous task of framing a new System; he may by this means give the public some idea of his self-consequence, and be enrolled in the Catalogue of System-makers, but not one jot will he advance the science of Botany. It is to be regretted that Botanists will not be contented with a System, a proof of whose superiority is the almost general reception it has met with throughout Europe, and unite in their endeavours to render that System more complete, by giving us an accurate account of the history of those plants not already given, their virtues and uses; this appears to me to be the true method of advancing this delightful Science, and making it useful to Mankind.

When one System of Botany is generally followed as is nearly the case at present, Botanists in different kingdoms perfectly understand each others language, but when each adopts a separate one, (which is frequently dictated by Pride or Caprice) it becomes Babel; and every one who wishes to acquire a knowledge of the plants treated of, must at considerable expence both of time and labour, acquire first the Authors new-created System-language, a tax which it is hoped every true Botanist will unite to oppose.



Euphorbia Peplus



EUPHORBIA HELIOSCOPIA. SUN SPURGE OR WART-WORT.

EUPHORBIA *Linnei Gen. Pl.* DODECANDRIA TRIGYNIA.

Cor. 4-l. 5-petala, calyci infidens. *Cal.* 1-phyllus, ventricosus.

Capsl. 3-cocca.

Raii Syn. Gen. 22. *HERBÆ VASCULIFERÆ, FLORE TETRAPETALO ANOMALÆ.*

EUPHORBIA umbella quinquefida : trifida : dichotoma, involucellis obovatis, foliis cuneiformibus ferratis. *Linn. Syst. Vegetab. p.* 377. *Sp. Plant.* 658. *Fl. Suecic. p.* 162.

TITHYMALUS foliis petiolatis, subrotundis, ferratis, stipulis rotundis, ferratis. *Haller bist. v.* 2. *p.* 10. n. 1050.

TITHYMALUS *helioscopius.* *Scopoli Fl. Carniol. p.* 337. n. 579:

TITHYMALUS *helioscopius.* *Bauhin Pin.* 291. *Gerard emac.* 458. *Parkinson.* 189.

TITHYMALUS *helioscopius* five folisequus. *I. B.* 3. 669. *Raii Syn.* 313. *Hudson Fl. Angl. p.* 183.

RADIX simplex, fibrosa, annua.

CAULIS erectus, teres, pilosus, inferne brachiatus, brachiis oppositis.

FOLIA sparsa, pauca, glabra, *ferrata*, cuneiformia, inferiora petiolata, superiora sessilia.

UMBELLA quinquefida, trifida, dichotoma, patens, fastigiata.

STIPULÆ minute ferratæ, glabræ, UMBELLÆ quinque, obovatæ, horizontales, æquales, Umbellulæ tres, ovatæ, inæquales, inferiore duplo minore, quæ sequuntur mucrone terminatæ.

CALYX subventricosus, flavescens, *fig.* 1.

COROLLA nulla.

NECTARIA quatuor, subrotunda, nuda, *fig.* 2.

STAMINA : FILAMENTA duo, tria, aut plura, visibilia, exserta; ANTHERÆ flavæ, biloculares, loculis subrotundis, *fig.* 3.

PISTILLUM : GERMEN pedunculatum, subrotundum, nutans; STIGMATA tria, apice bifida, *fig.* 4, 5.

PERICARPIUM : CAPSULA trilocca, trilocularis, trilvalvis, *fig.* 6.

SEMEN unicum in singulo loculamento, ovatum, rugosum ex purpureo fuscum, *fig.* 7.

ROOT simple, fibrous, annual.

STALK upright, round, slightly hairy, below branch-
ed, the branches opposite.

LEAVES growing in no regular order, few, smooth,
ferrated, and wedge-shaped, the lower ones
standing on foot-stalks, the upper ones sessile.

UMBELL dividing into five, next three, then two,
spreading, of an equal height at top.

STIPULÆ minutely ferrated and smooth, those of the
UMBELL five, somewhat oval, spreading hori-
zontally, and equal; those of the *smaller* UM-
BELL three, oval, unequal, the interior one
twice as small as the others; those which
follow terminating in a point.

CALYX somewhat swelled, of a yellowish colour,
fig. 1.

COROLLA wanting.

NECTARIA four, roundish and naked, *fig.* 2.

STAMINA : two, three, or more FILAMENTS, visible
beyond the Calyx; ANTHERÆ yellow, bilocu-
lar, the cavities containing the Pollen roundish,
fig. 3.

PISTILLUM : GERMEN placed on a foot-stalk, round-
ish, hanging down; STIGMATA three, bifid
at top, *fig.* 4, 5.

SEED-VESSEL a CAPSULE of three protuberating
valves, and three cavities, *fig.* 6.

SEEDS one in each cavity, oval, wrinkled, of a purp-
lish brown colour, *fig.* 7.


IN speaking of the *Euphorbia Peplus*, I had occasion to take notice of the difficulty which Students in Bota-
ny find in investigating the *Class* and *Order* of this Genus, and endeavoured to make it easier to them: in
this plant the parts of the fructification are somewhat larger; and it differs from the other Spurges in having
its leaves finely ferrated. In its acrimonious quality it is inferior to none; hence it has often been applied
to Warts for the purpose of destroying them; but even in this case, great care should be used in its applica-
tion. My friend Mr. WILLIAM WAVELL lately informed me of a case which fell under his notice in the
Isle of Wight, where from the application of the juice of this Spurge to some Warts near the eye of a little
girl, the whole face became inflamed to a very great degree.


It is very common in gardens and cultivated ground, flowering in Autumn.



Euphorbia Helioscopia.







POTENTILLA REPTANS. COMMON CINQUEFOIL OR FIVE LEAVED GRASS.

POTENTILLA *Linnaei Gen. Pl. ICOSANDRIA POLYGYNIA:*

Raii Gen. 15. HERBÆ SEMINE NUDO POLYSPERMÆ.

POTENTILLA *reptans* foliis quinatis, caule repente, pedunculis unifloris. *Lin: Syll. Vegetab. p. 398. Fl. Suecic. p. 178.*

FRAGARIA foliis quinatis serratis, petiolis unifloris, caule reptante. *Haller hist. v. 2. p. 47.*

QUINQUEFOLIUM majus repens. *Baubin pin. p. 325. Gerard emac. 987.*

PENTAPHYLLUM vulgatifolium *Parkinson 398. Raii Syn. p. 255.*

POTENTILLA *reptans. Hudfon. Fl. Angl. p. 197. Scopoli Fl. Carniol. p. 361*

RADIX perennis, fusiformis, paucis fibrillis instructa, intra terram profunde penetrans, crassitie digiti minimi aut pollicis etiam in annosis, externe fœdide castanea.

CAULES numerosi, teretes, glabri, repentes, purpurei.

FOLIA quinata, etiam septena occurrunt, serrata, venosa, inæqualia, parum hirsuta, petiolis longis indentia, per paria e geniculis caulium ad magna intervalla proceduntia.

STIPULÆ geminæ, trifoliatæ, foliolis ovatis.

PETIOLI uniflori, longi, suberecti.

CALYX: PERIANTHIUM monophyllum, planiusculum, decemfidum, laciniis alternis minoribus, sæpe reflexis, *fig. 3, 4, 5.*

COROLLA: PETALA quinque, subrotundo-cordata, flava, unguibus calyci inserta, *fig. 6.*

STAMINA: FILAMENTA viginti, subulata, Corolla breviora, margini interiori glandulosæ calycis inserta, in duas series distributa; ANTHERÆ oblongæ, compressæ, flavæ, biloculares, loculæ membranâ divisæ, insidentes, *fig. 7, 8.*

PISTILLUM: GERMINA numerosa, in capitulum collecta; STYLI filiformes filamentis breviores, lateri Germinis inserti, persistentes; STIGMATA minima, obtusa, *fig. 9, 10.*

SEMINA numerosa, parva, fusca, stylo persistente terminata, *fig. 11, 12.*

ROOT perennial, tapering, furnished with few fibres, penetrating deeply into the earth, the size of the little finger, or even of the thumb when old, externally of a dark chestnut colour.

STALKS numerous, round, smooth, and creeping.

LEAVES quinate, or growing five together, sometimes even seven, serrated, veiny, unequal in their size, slightly hairy, sitting on long footstalks, which proceed in pairs from the joints of the stalks at considerable distances.

STIPULÆ growing in pairs, composed of three oval-shaped leaves.

FOOT-STALKS of the flowers uniflorous, long, and nearly upright.

CALYX: a PERIANTHIUM of one leaf, flattish, divided into ten segments, the segments alternately smaller and frequently turned back, *fig. 3, 4, 5.*

COROLLA: five PETALS of a roundish heart-shaped figure, and yellow colour, inserted into the Calyx by their Ungues or claws, *fig. 6.*

STAMINA twenty FILAMENTS tapering: shorter than the Corolla, inserted into the inner edge of the Calyx, which puts on a glandular appearance, and placed in two rows; ANTHERÆ oblong, flat, bilocular, the bags or cavities divided by a membrane, sitting on the filaments, *fig. 7, 8.*

PISTILLUM: the GERMINA numerous, collected into a little head; the STYLES filiform, shorter than the filaments, inserted into the side of the Germin and continuing; the STIGMATA very small and blunt, *fig. 9, 10.*

SEEDS numerous, small, brown and terminated by the Style, *fig. 11, 12.*

The Roots of *Cinquefoil* and many other plants of the Class *Icosandria*, possess considerable virtues as astringent medicines, and may be used in the same Cases in which *Bistort* is recommended.

It has likewise been used in some places for the purpose of tanning Leather where better materials for that purpose are with difficulty acquired.

A Tea or infusion of the leaves is in use among the Country People as a drink in Fevers.

Most sorts of Cattle are fond of the leaves, but it does not appear to be a plant worth cultivating on that account.


The Larva or Caterpillar of the *Phalena Rubi*, *vid. Rosfel, Suppl. tab. 69, Albin tab. 81*, feeds on the leaves in Autumn, although a plant to which Insect is by no means confined.

It grows very common in meadows and on banks by the road sides, and flowers in July, August, and September.

It affords the botanic Student a very good example of the *Caulis repens* or *Creeping Stalk*.



Potentilla reptans





Ranunculus bulbosus.

RANUNCULUS BULBOSUS. ROUND-ROOTED OR BULBOUS CROWFOOT.

RANUNCULUS *Linnei Gen. Pl. POLYANDRIA POLYGYNIA.*

Raii Syn: Gen. 15. HERBÆ SEMINE NUDO POLYSPERMÆ.

RANUNCULUS *bulbosus*, calycibus retroflexis, pedunculis sulcatis, caule erecto multifloro, foliis compositis.

Linnei Syst. Vegetab. p. 430. Sp. Pl. 778. Fl. Suecic. 196.

RANUNCULUS radice subglobosa, foliis hirsutis, semitrilobis, lobis petiolatis acute serratis. *Haller. hist.*

v. 2. p. 74.

RANUNCULUS *Scopoli Fl. Carn. v. 1. p. 400. DIAGN. Radix globosa. Calyces reflexi. Squamula nectarifera obtuse trigona.*

RANUNCULUS *pratensis radice verticilli modo rotunda. Baubin. pin. 179. Fuschii Icon. 160. Gerard. emac. 953. Parkinson 329. Raii Synop. 247. Hudson Fl. Angl. 211. Fl. Dan. Icon. 551.*

RADIX perennis, *subrotunda*, albida, solida, superne et inferne depressior, hinc radicem Rapæ quodammodo referens.
CAULIS pedalis, teres, *erectus*, fistulosus, hirsutus, ramosus.
FOLIA *radicalia* petiolis longis, hirsutis, basi vaginantibus infidentia, subprocumbentia, hirsuta, venosa, trilobata, lobo medio majori et longius petiolato, semitrifido, segmentis acute incisis; lobis lateralibus trifidis, segmentis inferioribus profundius divisis; caulina subseffilia in lacinias plures tenuiores divisa.

PEDUNCULI *sulcati.*

CALYX: PERIANTHIUM pentaphyllum, foliolis ovatis, concavis, *reflexis*, pilosis, apice obtusiusculis, margine membranaceis, basi *subpellucidis*, *fig. 1.*

COROLLA PETALA quinque obcordata, flava, nitentia, *fig. 2.*

NECTARIUM: squamula flava subemarginata ad basin petali *fig. 3.*

STAMINA: FILAMENTA plurima, receptaculo inserta; ANTHERÆ oblongæ, flavæ, subincurvæ, *fig. 4.*

PISTILLUM: GERMINA numerosa in capitulum collecta; STYLI nulli; STIGMATA minima reflexa, *fig. 5.*

SEMINA plurima compressa, fusca, mucronata, lævia, arillata, *fig. 6.*

Fig. 7. ARILLUS, fig. 8. semen denudatum.

ROOT perennial, *roundish*, white and solid, flattened a little both at top and bottom, hence somewhat resembling a Turnep.

STALK a foot high, round, *upright*, hollow, hairy and branched.

LEAVES: the *radical* leaves placed on long hairy foot-stalks, which at bottom embrace the stalk, somewhat procumbent, hairy, veiny, and divided into three lobes; the mid-lobe largest and placed on a longer foot-stalk than the others, divided half way down into three segments which are sharply cut in; the side-lobes triid, the lower segments more deeply divided than the others; the leaves of the *stalk* nearly sessile, deeply divided into numerous and narrower segments.

FOOT-STALKS of the flowers *grooved.*

CALYX: a PERIANTHIUM of five leaves, the leaves oval, hollow, *turned back* and hairy, blunth at top, membranous at the edges, *thin and somewhat transparent* at bottom, *fig. 1.*

COROLLA: five PETALS, heart-shaped, yellow, and shining, *fig. 2.*

NECTARY: a small yellow scale at the bottom of the petal, with a slight indentation at top, *fig. 3.*

STAMINA: FILAMENTS numerous and inserted into the receptacle; ANTHERÆ oblong, yellow, and bending a little inwards, *fig. 4.*

PISTILLUM: GERMINA numerous, collected into a little head; STYLES none; STIGMATA very small and bending back, *fig. 5.*

SEEDS numerous, flat, brown, smooth, pointed, and covered with an Arillus, *fig. 6.*

Fig. 7. the ARILLUS, fig. 8. the seed taken out of it.

THIS *Crowfoot* has been considered by some Authors as the same Species with the *Ranunculus repens*, but certainly without any propriety, for there can be no doubt but they are as distinct as any two species of *Ranunculus* whatever. It is distinguished from the *repens* by several peculiarities, the principal of which are, 1st, its *reflexed calyx*, the turning back of which does not depend on any accidental circumstances, but solely on its particular structure; if it be plucked off, and held up to the light, the lower half of it will appear thin and almost transparent, hence not having a sufficient degree of solidity to support itself upright, it is reflected downwards;—2^{dy}, the root in this species is *round*, and *solid*; in the *repens* it is *fibrous*: and 3^{dy}, (which perhaps may be considered as the most essential difference) the stalk of the *bulbosus* is never known to *throw out any Stolones or Creepers*, which the *repens* always does in every soil and situation.

This Species blows earlier than either the upright or creeping *Crowfoot*, and is the second flower, which next to the Dandelion covers our meadows and pastures with that delightful yellow, which almost dazzles the eye of the beholder.

Like the rest of the *Crowfoots* it possesses the property of inflaming and blistering the skin, but more particularly the Root, which is said to raise blisters with less pain and more safety, than Spanish flies; and hence where Blisters have been thought necessary, these roots have been applied for that purpose, particularly to the Joints in cases of the Gout. On being kept they lose their stimulating quality, and are even eatable when boiled.

HOFFMAN informs us that Beggars make use of them to blister their skins in order to excite compassion.

The Juice of this herb is said to be more acrid than that of the *Ranunculus sceleratus*, and if applied to the nostrils it provokes sneezing.

Hogs are fond of the roots and will frequently dig them up.

It abounds in dry pastures, and flowers in May; it is cultivated when double as well as the upright meadow *Crowfoot*, which last occurs in almost every Garden, under the name of *Yellow Bachelors Buttons*.





Ranunculus acris

RANUNCULUS ACRIS. UPRIGHT MEADOW CROWFOOT.

RANUNCULUS *Linnei Gen. Pl.* POLYANDRIA POLYGYNIA.

Raii Gen. 15. HERBÆ SEMINE NUDO, POLYSPERMÆ.

RANUNCULUS *acris* calycibus patulis, pedunculis teretibus, foliis tripartito-multifidis, fannis linearibus.
Linnei Syll. Vegetab. p. 430. Fl. Suecic. p. 196.

RANUNCULUS foliis hirsutis, semitrilobatis, lobis lateralibus bipartitis, foliis caulinis semitrilobis. *Haller. Hist. n. 1169.*

RANUNCULUS pratensis erectus *acris.* *Baubin. pin. 178. Gerard. emac. 951. Parkinson 329. Raii Synopsi, p. 248. Hudson. Fl. Angl. p. 211. Scopoli. Fl. Carniol. p. 398.*

RADIX perennis, e pluribus radiculis albidis constans.

ROOT perennial, consisting of numerous white fibres.

CAULIS bipedalis, erectus, fistulosus, teres, subpilosus, apice ramosus.

STALK generally about two feet high, upright, hollow, round, somewhat hairy, much branched at top.

FOLIA *Radicalia* petiolis longis erectis infidentia, tripartita, lobo medio trifido, lateralibus bilobis, omnibus acute dentatis aut incis, subhirsutis, supernè ad basin præsertim sæpe purpureis, venis fubtus extantibus.

LEAVES: *Radical* leaves standing on long upright foot-stalks, tripartite, the middle lobe trifid, the side ones bilobous, and all of them sharply indented, slightly hirsute, the upper surface particularly at the base frequently of a purple colour, the veins underneath prominent

Caulina radicalibus similia, in lacinias tenuiores vero divisa et petiolis brevioribus infidentia, tandem linearia, sessilia.

Leaves of the *Stalk* like the radical leaves, but more finely divided, and standing on shorter foot-stalks, at top linear and sessile. The FOOT-STALKS with their sheaths hairy.

PETIOLI cum vaginis hirsuti.

PEDUNCULI teretes.

FOOT-STALKS of the Flowers round.

CALYX: PERIANTHIUM pentaphyllum, patens, flavescens, pilosum, foliolis ovatis, concavis, obtusis, margine membranaceis, fig. 1.

CALYX: a PERIANTHIUM of five leaves, spreading, of a yellow colour and hairy, the leaves oval, concave, and membranous at the edges, fig. 1.

COROLLA: PETALA quinque flava, nitentia, subcordata nunc emarginata, nunc integra, fig. 2.

COROLLA: five PETALS, yellow and shining, nearly heart-shaped, sometimes notched, sometimes entire, fig. 2.

STAMINA: FILAMENTA plurima, apice paululum dilatata, fig. 5. 4. ANTHERÆ flavae, subincurvatae, obtusæ, fig. 4.

STAMINA: FILAMENTS numerous, a little dilated at top, fig. 5. 4. ANTHERÆ yellow, obtuse, bending a little inward, fig. 4.

NECTARIUM: squamula emarginata, ad basin petalorum, fig. 3.

NECTARY: a small scale, slightly notched at top, at the base of each Petal, fig. 3.

PISTILLUM: GERMINA numerosa, in capitulum collecta, STYLI nulli; STIGMATA reflexa, fig. 6.

PISTILLUM: GERMINA numerous, forming a little head; STYLES none, STIGMATA reflex, fig. 6.

SEMINA: plurima, subrotunda, compressa, fusca, apice reflexa. fig. 7.

SEEDS numerous, roundish, flat, of a brown colour, bending back at the tip, fig. 7.

Most of the Ranunculi or Crowfoots are acrid and in some degree poisonous, and the species above described possess this property in a very considerable degree; hence Linnaeus has given it the name of *acris*; even pulling up the plant and carrying it to some little distance we have known sufficient to produce a considerable inflammation in the palm of the person's hand who held it. Cattle in general will not eat it, yet sometimes when they are turned hungry into a new field of Grass, or have but a small spot to range in they will feed on it, and hence their mouths, as we have been credibly informed, have become sore and blistered. When made into hay it loses its acrid property, but is too stinky and hard to afford good Nourishment. It should seem therefore to be the interest of the Farmer as much as possible to root out this species from his Meadows that its place may be supplied with good sweet grass.

It grows too frequently in most of our meadows, and flowers in June and July.

The common people about Town and in many parts of the country call this and the other yellow Crowfoots by the names of *Butter-cups* and *Butter-flowers*, and this name seems to have originated from a supposition that the yellow colour of butter was owing to these plants; that this should be the case seems scarce probable, certainly it receives no good taste from it.

CALTHA PALUSTRIS. MARSH-MARIGOLD.

CALTHA *Linnei Gen. Pl.* POLYANDRIA POLYGYNIA *Cal.* 6. Petala quinque. Nectaria 6. Capsulæ plures polyspermæ.

Raii Syn. HERBÆ MULTISILIQUÆ SEU CORNICULATÆ.

CALTHA palustris. *Linnei Syst. Vegetab. p.* 432. *Flor. Suecic.* 198.

CALTHA *Haller. bist. belo. p.* 32. n. 1188.

POPULAGO *palustris. Scopoli Fl. Carniol. p.* 404.

CALTHA palustris flore simplici. *Bauhin pin* 276.

POPULAGO. *Tournefort. Tabernamont.*

CALTHA palustris vulgaris simplex. *Parkinson* 1213.

CALTHA palustris major. *Gerard. emac.* 817.

Raii Syn. 272. Marsh Marigold. *Hudson Fl. Angl. p.* 214.

RADIX perennis, e plurimis fibris, teretibus, majusculis, albidis, constans.

CAULES ex eadem radice nascuntur plures, suberecti, pedales, fistulosi, pene teretes, glabri, ramosi, ad basin purpurei.

FOLIA radicalia petiolata, cordato-reniformia, glabra, crenata, caulina subsessilia, ad apicem acutiora, et acute crenata.

STIPULÆ fuscae, membranaceæ, marcescentes.

RAMI dichotomi.

PEDUNCULI uniflori, erecti, sulcati.

CALYX nullus.

COROLLA: PETALA plerumque quinque, flava, magna, subrotundo-ovata, plana, patentia, superne non splendentia, *fig.* 1.

STAMINA: FILAMENTA numerosa, filiformia, Corollâ breviora, ANTHERÆ oblongæ, compressæ, incurvatæ, flavæ, *fig.* 2.

PISTILLUM: GERMINA quinque ad decem, oblonga, compressa, erecta; STYLI nulli; STIGMATA simplicia, *fig.* 3.

PERICARPIUM: CAPSULÆ totidem, acuminatæ, patentēs, futura superiore dehiscentes, *fig.* 4.

SEMINA plurima, subovata, pulchra, inferne olivacea, superne rufa, *fig.* 5.

ROOT perennial, consisting of numerous, round, large, white fibres.

STALKS: several arise from the same root, almost upright, about a foot high, hollow, nearly round, smooth, branched, and purple at bottom.

LEAVES: the radical leaves placed on long foot stalks, betwixt an heart and kidney shape, smooth, shining, and notched or crenated; the leaves of the STALK nearly sessile, more pointed at top, and sharply crenated.

STIPULÆ brown, membranous and withered.

BRANCHES dichotomous.

PEDUNCLES supporting one flower, upright, and grooved.

CALYX wanting.

COROLLA generally consists of five large PETALS of a roundish oval shape and yellow colour, flat, spreading, and without any gloss on the upper side, *fig.* 1.

STAMINA: FILAMENTS numerous, filiform, shorter than the Corolla; ANTHERÆ oblong, flat, bending inward, and of a yellow colour, *fig.* 2.

PISTILLUM: GERMINA from five to ten, oblong, flattish, and upright; STYLES none; STIGMATA simple, *fig.* 3.

SEED-VESSEL: so many CAPSULES as Germina, pointed, and spreading, opening at the superior future, *fig.* 4.

SEEDS numerous, somewhat oval, beautiful, at bottom of an olive, and at top of a reddish colour.

LINNEUS informs us that the *Caliba* is the first flower which proclaims the Spring in *Lapland*, and that it begins to blow about the end of May, with us it usually flowers in March and April, and last Spring, 1775, this plant was found in Blossom in the month of February, so remarkably forward was the Spring of that year.

It grows in wet Meadows and by the sides of Rivers, where it makes a very noble appearance, and when double, is often cultivated in Gardens, where it will grow very readily if the soil be favourable.

In the Country, Children collect it to ornament their Garlands on May day.


I scarce ever observed the leaves to be eaten by any animals, but the flowers are often destroyed by a species of CHRYSOMELE.

HALLER says that it is acrid and caustic and yet that it is eaten by Cows.

The flower Buds are pickled and used as Capers.



Callithra pulchra





16. 2. p. 2.

Verbena officinalis.

VERBENA OFFICINALIS. VERVAIN.

VERBENA *Lin. Gen. Pl. DIDYNAMIA GYMnosPERMIA.*
Raii Gen. 14. SUFFRUTICES, ET HERBÆ VERTICILLATÆ.
 VERBENA *officinalis*, tetrandra, spicis filiformibus, paniculatis; foliis multifo-ido-lacinatis, caule solitario.
Lin. Syst. Vegetab. p. 62.
 VERBENA foliis tripartitis rugosis, spicis nudis gracilissimis *Haller. hist. v. 1. p. 96.*
 VERBENA communis cæruleo flore. *Bauhin, Pin. 260. mas, seu recta et vulgaris. Parkinson 674. communis Gerard 664. Raii Syn. 236. Hudson Fl. Angl. p. 505. Scopoli Fl. Carniol. p. 433.*

<p>RADIX perennis, lignosa, crassitie digiti minimi, raro major, in terram profunde penetrans, fibrosa, lutescens, sapore subamaro.</p> <p>CAULES plerumque plures ex eadem radice, erecti, pedales aut bipedales, quadrangulares, duo latera excavata, duo subconvexa, fuscata, idque alterne, aculeis brevibus armati, brachiati.</p> <p>FOLIA opposita, sessilia, venosa, profunde dentata, aut incisa, ad basin angustiora.</p> <p>FLORES in spicas longas, filiformes, erectas dispositi, BRACTEA ovato-lanceolata, acuminata, calyce breviori suffulti, fig. 11.</p> <p>CALYX: PERIANTHIUM monophyllum, angulatum, quinqueidentatum, <i>denticulo quinto minimo, persistens, fig. 1 2, 3.</i></p> <p>COROLLA monopetala, inæqualis, purpurascens, TUBUS cylindraceus, incurvatus; FAUX villosa, fig. 5; LIMBUS quinquefidus, lacinii rotundatis, subæqualibus, fig. 4.</p> <p>STAMINA: FILAMENTA quatuor brevissima, vix conspicua, ANTHERÆ quatuor, quarum duæ breviores reliquis, ejusdem formæ cum Didynamis fig. 6.</p> <p>PISTILLUM: GERMEN tetragonum, STYLUS filiformis apice paululum incrassatus; STIGMA obtusum fig. 7.</p> <p>PERICARPIUM nullum, Calyx continens Semina.</p> <p>SEMINA quatuor, oblonga, obtusa, interne planiuscula alba, externe fusca, convexa, <i>fulcato-reticulata fig. 8, 9, 10.</i></p>	<p>ROOT perennial, woody, about the thickness of the little finger, seldom larger, running deep into the earth, fibrous, of a yellowish colour, and slightly bitter taste.</p> <p>STALKS: in general several arise from the same root, upright, from one to two feet high, four square, two sides hollowed out, two roundish and grooved, and that alternately, armed with short prickles, the branches alternately opposite.</p> <p>LEAVES opposite, sessile, veiny, deeply indented or cut in, narrowest at bottom.</p> <p>FLOWERS disposed in long filiform erect spikes, supported by an oval pointed FLORAL-LEAF shorter than the Calyx, fig. 11.</p> <p>CALYX: a PERIANTHIUM of one leaf, quinqueidentate, the fifth tooth exceedingly minute, continuing, fig. 1, 2, 3.</p> <p>COROLLA monopetalous, unequal, purplish, the TUBE cylindrical and crooked, the MOUTH villous, fig. 5. the LIMB divided into five segments, which are round and nearly equal, fig. 4.</p> <p>STAMINA: four FILAMENTS very short and scarce conspicuous, four ANTHERÆ two of which are above the others, of the same form with those of the Class Didynamia in general, fig. 6.</p> <p>PISTILLUM: the GERMEN four square, the STYLE filiform, growing thicker towards the extremity, the STIGMA obtuse, fig. 7.</p> <p>PERICARPIUM wanting, the CALYX containing the Seeds.</p> <p>SEEDS four, oblong, obtuse, on the inside flatish and white, on the outside brown, convex, grooved and reticulated, fig. 8, 9, 10.</p>
---	---

The Vervain may be considered as a kind of domestic plant, not confined to any particular soil, but growing by the road sides, pretty universally at the entrance into Towns and Villages.

It produceth its blossoms in the months of August and September.

There is only one Species of this Genus which grows wild in this country, but in different parts of the world the species are numerous, and what is remarkable, some have four and others but two Stamina, hence LINNÆUS ranks them among his *Didynamia* plants, making a division of them into such as have *flores Diandri* and *flores Tetrandri*. As our species hath four Stamina, two of which are above the other two, as the Style proceeds from the center of the four united Germens, and as four naked seeds follow, which are contained within the Calyx, we have placed it with SCOPOLI among the *Didynamia Gynnospermia* plants, a Class to which the botanic Student, who had been instructed in the Linnæan principles of Botany, would readily have been induced to refer it.

The seed of this plant has something remarkably curious in its appearance, on the inside it is of a snowy white, externally brown, and beautifully reticulated.

The Plant which the Romans called *Verbena*, appears to have been used on particular occasions at a very early period, as a token of mutual confidence betwixt them and their Enemies. It was also constantly applied to the purposes of Superstition and Enchantment, in making wreaths and brooms for their Altars, and chaplets for their Priests. It is probable from *Pliny's* account, that the plant which we now describe was the same with that of the Antients, but in a larger sense, they called the Laurel and Myrtle or whatever was bound round the Altar *Verbena*. The dry harsh nature of this herb, agrees but ill with the *Pinguis Verbena* of Virgil, perhaps it acquired that title from being anointed with the fat of the sacrifice.

In later times Vervain has been accounted a sovereign remedy in a multitude of disorders; SCHRODER recommends it in upwards of thirty different complaints, on which Mr. Ray judiciously observes "*Mirum tot viribus pollere plantam nulla insigni qualitate sensibili dotatam*!" strange that a plant which inherits no remarkably sensible quality should possess so many virtues!


Mr. Morley a late writer on the Vervain, considers it as extremely useful in the cure of the Schrophula or Kings evil, and in his Essay on the nature and cure of Schrophulous diseases, has given us a figure of the plant with particular directions for its use, which consists in hanging the root (which is to be of a larger or smaller size according to the age of his Patients) tied with a yard of white fatten ribband round the neck, there to be worn till they recover.

Those who know any thing of the effects of Medicines on the human body, will not easily be persuaded that such a kind of application can produce any very wonderful effect in this case, even making the greatest allowance for the powers of the imagination; and Mr. Morley as if sensible of the inefficacy of his Vervain Amulet, calls to his assistance a number of powerful medicines, among others we find Mercury, Antimony, Hemlock, Jalap, &c; and by a repeated and oftimes a long continued application of Baths, Cataplasms, Ointments, Poulices, Plasters, &c. and the exhibition of gentle purges and alterative medicines, some have been relieved and others cured; but can any one hence infer with any degree of reason that the Vervain Root had any share in the cure? certainly no; out of all Mr. Morley's cases there is not one which proves it, and the virtues of this plant still remain to be ascertained by rational experiments.

It should be observed that the Schrophula is a disease which at certain periods of life and at certain seasons of the year, is liable to be much worse than at others, and frequently exceeding bad cases of this kind have been cured by the most simple applications.

Many people have no doubt applied to Mr. Morley from a supposition that his motives were perfectly disinterested, and it must be confessed that there are Empirics much more mercenary and infinitely more dangerous; yet it does not appear but Mr. Morley acts nearly on the same principle with other Practitioners in Physick, with this difference indeed, that they receive their fees in specie, he takes his in kind.

That we may not be thought to act dissingenuously by Mr. Morley we shall quote his own words—"Many many Guineas have been offered me but I never take any money. Sometimes indeed genteel People have sent me small acknowledgements of Tea, Wine, Venison, &c. Generous ones, small pieces of Plate or other little Presents. Even neighbouring Farmers a Goose or Turkey, &c. by way of Thanks.



LAMIUM PURPUREUM. RED LAMIUM OR DEAD NETTLE

LAMIUM *Linnei Gen. Pl.* DIDYNAMIA GYMNOSPERMIA. *Corollæ* labium superius integrum, fœnicatum, labium inferius bilobum; faux utrinque margine dentata. *Lin. Descrip. Gen. abbrev.*

Raii Syn. Gen. 14. SUFFRUTICES ET HERBÆ VERTICILLATÆ.

LAMIUM *purpureum* foliis cordatis obtusis petiolatis. *Linnei Syst. Vegetab.* p. 446. *Sp. Pl.* 809. *F. Succ.* 203.

LAMIUM foliis cordatis, obtusis, in summo ramo congestis. *Haller. infl.* v. 1. 1:8.

LAMIUM *purpureum*. *Scopoli Fl. Carniol.* p. 407. n. 701.

LAMIUM *purpureum* fetidum, folio subrotundo, sive Galeopsis Dioforidis. *Bauhin. p'n.* 230. *Lanium rubrum*. *Gerard emac.* 703. *Parkinson.* 604. *Raii. Synesis* Small Dead Nettle or red Archangel 240. *Hudson. Fl. Angl.* 225. *Oeder. Fl. Dan. icon.* 523.

RADIX annua, fibrosa.

CAULES plures, ad basin debiles, et ramosi, prope summitatem fore nudi, et sæpe colorati, semipedales, quadrangulares, fistulosi, scabriusculi.

FOLIA opposita, venosa; hirsutula, inferiora subrotundo-cordata, crenata, longe petiolata: superiora ovato-cordata, obtuse serrata, petiolis brevibus infidentia, alterne opposita, reflexa, dense et imbricatim congesta, et rubedine tincta.

FLORES purpurei, in summis caulibus verticillatim densius stipati. Verticilli multiflori.

CALYX: PERIANTHIUM monophyllum, tubulatum, superne patentius, quinque-dentatum, substriatum, hirsutulum, dentibus subæqualibus, acuminatis. *fig. 1.*

COROLLA monopetala, ringens, pallide purpurea, *fig. 2*; TUBUS brevis, cylindraceus, *fig. 6*; FAUX inflata, margine utroque bidentata, *fig. 4*; denticulo superiori spinæ simili, inferiore obtusiore, macula notata; labium superius, *fig. 3*, ovatum, concavum, villosulum, integrum, labium inferius bilobum, maculatum, lobis patentibus. *fig. 5.*

STAMINA: FILAMENTA quatuor, subulata, alba, sub labio superiori tecta, quorum duo longiora, *fig. 7*; ANTHERÆ oblongæ, barbatae, polline croceo repletæ. *fig. 8.*

PISTILLUM: GERMEN quadrifidum; STYLUS filiformis, longitudine et situ staminum; STIGMA bifidum, acutum, *fig. 9, 10, 11.*

SEMINA 4 in fundo calycis, pallida, triangularia, apice truncata, marginata, *fig. 12.*

ROOT annual and fibrous.

STALKS several, at bottom weak and branched, near the top almost naked, and frequently coloured, six inches or more in height, quadrangular, hollow, and slightly rough.

LEAVES opposite, veiny, slightly hairy, the lower ones of a roundish-heart shaped form, notched, and placed on footstalks, the uppermost ones oval-heart-shaped, obtusely serrated, with short footstalks, alternately opposite, growing thickly together, bent back and laying one over another, of a reddish colour.

FLOWERS purple, growing thickly together on the tops of the stalks in whorls; many flowers in each whirl.

CALYX: a PERIANTHIUM of one leaf, tubular, at top spreading, with five teeth, somewhat striated and hairy, the teeth nearly equal and long pointed. *fig. 1.*

COROLLA monopetalous, gaping, of a pale purple colour, *fig. 2*; the TUBE short and cylindrical, *fig. 6*; the ENTRANCE of the TUBE inflated, the margin on each side furnished with two teeth, *fig. 4*; the uppermost pointed like a thorn, the lowermost blunter with a spot on it; the upper lip *fig. 3*; oval, hollow, slightly villous, entire, the under lip divided into two lobes, spreading a little from one another, and spotted, *fig. 5.*

STAMINA: four FILAMENTS, tapering and white, hid under the upper lip, two of which are longer than the rest *fig. 7*; the ANTHERÆ oblong, bearded, and full of a yellow pollen *fig. 8.*

PISTILLUM: GERMEN quadrifid; STYLUS filiform, the length of the Stamina; STIGMA bifid and pointed *fig. 9, 10, 11.*

SEEDS 4 in the bottom of the Calyx, of a pale brown, triangular, cut off as it were at top, with a margin round them, *fig. 12.*

Although this plant may perhaps with propriety be considered as a Weed in Gardens, yet the bright colour of its tops and flowers, joined to its early appearance, contributes not a little to ornament our banks in the Spring, when few other plants appear in blossom.


The Flowers are most commonly of a bright red colour, sometimes white, and are much resorted to by Bees of various kinds.

The Leaves and Flowers are those parts of the plant, which are used in Medicine, although in the present practice they are scarce regarded.

According to LINNÆUS it is boiled in *Upland*, a Province of *Sweden*, as a pot herb. A Variety of this plant occurs not unfrequently about Town, which has its leaves more deeply indented. RAY calls it *Lanium rubrum minus, foliis profunde incis.* I have found it growing on a bank on the right hand side of the way between *Pimlico* and *Celsea*.



Lamium purpuraceum.





Thymus acinus.

THYMUS ACINOS. BASIL THYME.

THYMUS *Linnaei Gen. Pl. DIDYNAMIA GYMNOSPERMIA.*

Calyx bilabiati faux villis clausa.

Raii Synop. Gen. 14. SUFFRUTICES ET HERBÆ VERTICILLATÆ.

THYMUS *Acinos* caulibus adscendentibus, foliis dentato-ferratis, calycibus basi ventricosif.

THYMUS *Acinos* floribus verticillatis, pedunculis unifloris caulibus erectis subramosis, foliis acutis, ferratis. *Linn. Syst. Vegetab. p. 452. Flor. Suecic. p. 209.*

CLINOPODIUM foliis ovatis acutis ferratis, flore foliis brevior. *Haller. hist. helv. n. 237.*

THYMUS *Acinos. Scopoli Fl. Carniol. p. 426. n. 735.*

CLINOPODIUM arvense ocimi facie. *Baubin. pin. p. 225.*

CLINOPODIUM minus five vulgare. *Parkinson. 21.*

OCYUM sylvestre. *Gerard. emac. 675.*

ACINOS multis. *Baubin. hist. 32. 259. Raii Syn. p. 238. Wild Basil. Hudson Fl. Angl. p. 230.*

RADIX annua, simplex, fibrosa.

CAULES adscendentes, semipedales, tetragoni, ramofi, hirsuti, purpurascetes; RAMI cauli similes longi, patentes, imi oppositi.

FOLIA opposita, petiolata, ovato-acuta, medium interius petiolo proximum integrum, exterius mucroni proximum dentatum, margines paululum reflexi, ciliati, nervo medio venisque subtus hirsuti, superne vix hirsuta, impunctata, venis quam in terpyllo profundius exaratis.

FLORES pedunculati, verticillati, spicati, plerumque sex in singulo verticillo.

CALYX: PERIANTHIUM monophyllum, tubulatum, basi ventricosum, striatum, hirsutum, quinque-dentatum, dentibus tribus superioribus brevioribus, reflexis, inferioribus setaceis, fauce villis clauso, fig. 1.

COROLLA monopetala, tubulosa, purpurea, bilabiata, labium superius brevius, obtusum, reflexum, emarginatum, inferius trifidum, lacinii subrotundis, medio productiore subemarginato, macula alba, lunulata, prominente, notata, fig. 3, 4, 5.

STAMINA: FILAMENTA quatuor, quorum duo longiora, Corollæ breviora; ANTHERÆ parvæ, rubræ, fig. 6.

PISTILLUM: GERMEN quadripartitum; STYLUS filiformis longitudine Stamina; STIGMA bifidum, acutum, fig. 7.

PERICARPIUM nullum

SEMINA quatuor oblonga intra Calycem, fig. 8, 9.

ROOT annual, simple and fibrous.

STALKS adscending, about six inches high, square, branched, hirsute, purplish; BRANCHES like the stalk, long, spreading, the bottom ones opposite.

LEAVES opposite, standing on foot-stalks, of a pointed oval shape, the inner middle part of them next the foot-stalks entire, the outer middle part next the point indented, the edges turned a little back and ciliated, the midrib and veins on the under side of the leaf hirsute, the upper surface of the leaves scarcely hairy, without any dots, the veins deeper than in the common Wild Thyme.

FLOWERS growing on foot-stalks, in whirls, forming a spike, generally six in each whirl.

CALYX: a PERIANTHIUM of one leaf, tubular, bellying out at bottom, striated, hirsute, having five teeth, the three uppermost of which are shortest and turned back, the lower ones slender and tapering, the mouth closed up with short hairs, fig. 1.

COROLLA monopetalous, tubular, purple, having two lips, the uppermost of which is shortest, blunt, turned back, with a slight notch in it; the lowermost divided into three roundish segments, the middle one of which is longer than the others, very slightly notched in, and marked with a raised white semilunar spot, fig. 3, 4, 5.

STAMINA: four FILAMENTS, two long and two short, within the Corolla; ANTHERÆ small and red, fig. 6.

PISTILLUM: GERMEN divided into four parts; STYLE filiform, the length of the Stamina; STIGMA bifid and acute, fig. 7.

SEED-VESSEL none.


SEEDS. Four oblong seeds within the Calyx, fig. 8, 9.

As there are only two species of *Thyme* growing wild in this Kingdom, and those very different from each other, the young Botanist cannot be at a loss in distinguishing them; with the *Thymus alpinus*, (figured by that accurate Botanist JACQUIN, in his *Fl. Aylriac*, who has contributed much to the advancement of botanic knowledge,) this plant has a much greater affinity, but may be distinguished by attending to the size of the flowers and the shape of the Calyx: the flowers of the *alpinus* are nearly twice as large as those of the *acinos*, and the Calyx of the latter has a protuberance at its base which we do not find either in the *alpinus* or *serpyllum*; a white circular mark in the mouth of the flowers, makes the blossoms of this species strikingly different from those of Wild Thyme.

The most common place of growth for this plant is in uncultivated fields, particularly where the soil is chalky, about *Charlton* it is found in abundance, flowering in July and August.

A variety with a white flower sometimes occurs.

The same agreeable aromatic flavour predominates in this species as in the Wild Thyme, whence it is probable that their virtues are very similar.





Cynophrasia Odontites.

EUPHRASIA ODONTITES. RED EYE-BRIGHT.

EUPHRASIA *Linnæi* Gen. Pl. DIDYNAMIA ANGIOSPERMIA.

Raii Syn. Gen. HERBÆ FRUCTU SICCO SINGULARI FLORE MONOPETALO.

EUPHRASIA *Odontites* foliis linearibus: omnibus ferratis. *Linnæi* Syst. Vegetab. Sp. Pl. p. 841. Fl.

Suecic. p. 213. n. 544.

ODONTITES bractæis ferratis hirsutis. *Haller.* hift. v. 1. p. 134. n. 304.

EUPHRASIA *Odontites.* *Scopoli* Fl. Carniol. p. 435.

EUPHRASIA *pratensis* rubra. *Baubin* Pin. p. 234.

EUPHRASIA *pratensis* rubra major. *Parkinson* 1329.

CRATÆOGONON *Euphrofync.* *Ger. emac.* 91. *Raii* Syn. p.* 284. Eye-bright Cow-wheat. *Hudson* Fl. Angl. p. 234.

RADIX annua, simplex, fibrosa, lignea.

CAULIS erectus, ramosissimus, semipedalis, ad bipedalem, hirsutus, obtuse quadrangularis.

RAMI cauli similes, oppositi.

FOLIA alterne opposita, sessilia, lineari-lanceolata, reflexa, rariter dentata, hirsutula, venosa, venis parvis, subtus hirsutis.

BRACTEÆ lanceolatae, suberectae, purpurascetes.

FLORES spicati, secundi, spicis apice subnutantibus,

CALYX: PERIANTHIUM monophyllum, tubulosum, quadridentatum, hirsutum, dentibus aequalibus, acutis, fig. 1.

COROLLA monopetala, ringens, labium superius concavum, submarginatum, inferius tripartitum, lacinis obtusis, aequalibus, fig. 2.

STAMINA: FILAMENTA quatuor, quorum duo paulo breviora, alba; ANTHERÆ bilobae, biloculares, apice filamentosae, basi spinulis duabus terminatae, deorsum ubi filamentum inferitur appendiculis clavatis pluribus instructae, fig. 3, 4, 5.

PISTILLUM: GERMEN ovatum, hirsutulum; STYLUS filiformis, in flore nondum explicato sub labio superiore Corollae involutus, postea Corollam longior; STIGMA capitatum, fig. 6.

PERICARPIUM: CAPSULA ovato-oblonga, compressa, bilocularis, fig. 7.

SEMINA plurima, albidia, striata, fig. 8.

* ROOT annual, simple, fibrous, and woody.

STALK upright, very much branched, from six inches to two feet high, hirsute, and obtusely square.

BRANCHES like the stalk and opposite.

LEAVES alternately opposite, sessile, betwixt linear and lanceolate, turning back, thinly indented, slightly hirsute, veiny, veins few and hirsute underneath.

BRACTEÆ lanceolate, nearly upright, purplish.

FLOWERS growing in spikes of a red colour, inclined all one way, the spikes nodding a little at top.

CALYX: a PERIANTHIUM of one leaf, tubular, quadridentate, hirsute, the teeth equal and sharp, fig. 1.

COROLLA monopetalous, gaping, the upper lip concave and slightly notched in; the lower lip divided into three, obtuse, equal segments, fig. 2.

STAMINA: four FILAMENTS, two somewhat longest, white; ANTHERÆ composed of two lobes and two cavities, at top thready, at bottom terminated by two little spines, and on the back part where the filament is inserted, furnished with several small club-shaped threads or appendages, fig. 3, 4, 5.

PISTILLUM: GERMEN oval, hirsute; STYLE filiform, before the flower opens bent in underneath the upper lip of the Corolla; afterwards longer than the Corolla; STIGMATA forming a little head, fig. 6.

SEED-VESSEL an oval, oblong, flattened CAPSULE, of two cavities, fig. 7.

SEEDS several, whitish and striated, fig. 8.

This species of *Eyebright*, which is exceedingly different from the common sort, grows very common in Pastures, sometimes in Corn-fields, and flowers in July and August: it differs very much in size according to the place it grows in, and is now and then found with white flowers.

It is not remarked either for its beauty or utility.



ANTIRRHINUM CYMBALARIA. IVY-LEAV'D ANTIRRHINUM.

ANTIRRHINUM *Linnei Gen. Pl. DIDYNAMIA ANGIOSPERMIA.*

Raii Syn. HERBÆ FRUCTU SICCO SINGULARI FLORE MONOPETALO.

ANTIRRHINUM *Cymbalaria foliis cordatis quinquelobis alternis, caulibus procumbentibus. Linnei*

Syst. Vegetab. p. 464. Sp. Pl. p. 851.

ANTIRRHINUM *caule repente, foliis reniformibus, quinquelobatis. Haller hist. p. 146. n. 339.*

ANTIRRHINUM *Cymbalaria Scopoli Fl. Carniol. n. 770.*

CYMBALARIA *Bauhin pin. 306.*

LINARIA *hederaceo folio glabro, seu Cymbalaria vulgaris. Tourn. 169. Gariel. 287. Gouan. Fl. Monsp. p. 100. Gerard Fl. Galloprov. p. 292. Raii Syn. p. *282. Hudson Fl. Angl. p. 237.*

Tota Planta glabra, cum odore ingrato.

RADIX perennis, fibrosa, intra fissuras murorum penetans; eradicatione difficilis.

CAULES plures, confertim nascuntur, basi repentes, procumbentes, ramosi, teretes, glabri, purpurascens, nervo intus duriore et tenaciore sicut in Alfine.

FOLIA quinquelobata, glabra, subcarnosa, opposita, aut alterna, sæpe purpurascens, fig. 12.

PETIOLI longi, superne sulcati.

PEDUNCULI teretes, petiolis paulo longiores.

CALYX: PERIANTHIUM quinquepartitum, lacinii lanceolatis, persistentibus, fig. 1.

COROLLA monopetala, ringens; TUBUS brevis, fig. 6; LIMBUS bilabiatus, labium superius bifidum, reflexum, purpureum, venis duabus saturatioribus striatum, fig. 2. inferius trifidum, lacinii subrotundis, albidis, fig. 3; PALATUM prominens, bifidum, flavum, fig. 5, FAUX villosa, croceum.

NECTARIUM purpureum, conicum, longitudine calycis, fig. 5.

STAMINA: FILAMENTA quatuor, duo breviora; ANTHERÆ bilobæ, albæ, conniventes, fig. 7.

PISTILLUM: GERMEN subrotundum, purpureum; STYLUS filiformis; STIGMA obtusum, fig. 8.

PERICARPIUM CAPSULA subrotunda, rugosa, seminibus protuberantibus, bivalvis, valvis apice in plures lacinias dehiscens, fig. 9, 10.

SEMINA nigra, subrotunda, rugosa, fig. 10.

The whole plant smooth, with a disagreeable smell.

ROOT perennial, fibrous, penetrating between the crevices of the walls, and scarce to be eradicated.

STALKS numerous, growing in a kind of tuft, creeping at bottom, procumbent, branched, round, smooth, purplish, and stringy as in Chickweed.

LEAVES quinquelobate, smooth, somewhat fleshy, some of them opposite, others alternate, frequently purplish, fig. 12.

FOOT-STALKS of the leaves long, on the upper part grooved.

FOOT-STALKS of the flowers, round, a little longer than the foot-stalks of the leaves.

CALYX: a PERIANTHIUM divided into five segments, which are lanceolate and continuing, fig. 1.

COROLLA monopetalous, ringent; the TUBE short, fig. 6: the LIMB divided into two lips; the upper lip bifid, turning back, and purple, striped with two veins of a deeper colour, fig. 2; the lower lip trifid, the segments round and whitish, fig. 3; the PALATE prominent, bifid, and yellow, fig. 4; the MOUTH or entrance into the tube villous and saffron-coloured.

NECTARY purple, conical, the length of the Calyx, fig. 5.

STAMINA: four FILAMENTS, two short and two long; ANTHERÆ composed of two lobes, white and connivent, fig. 7.

PISTILLUM: GERMEN roundish and purple; STYLE filiform; STIGMA blunt, fig. 8.

SEED-VESSEL a roundish CAPSULE, surface uneven, from the seeds protuberating, of two valves, which open at top into several lacinie, fig. 9, 10.

SEEDS black, roundish and wrinkled, fig. 10.

This Species of *Antirrhinum* is so perfectly distinct from all the others which grow wild in this country, that there is no possibility of mistaking it. It is found in great plenty in all those parts near London, that lay within the reach of the Thames; the seeds are carried by the flux and reflux of the tide up and down the river, and left at high water mark in the crevices of old walls, where they take root and encrease very fast. It is supposed to have been introduced to us from Italy, whether for the purposes of ornament or medicine is uncertain.

The Walls of the *Physic Garden*, at *Chelsea*, from whence it has probably originated in this country, are plentifully covered with it; it may also be found on the *Temple Walls*, and at the sides of the stream running under *Vauxhall Turnpike*.

In some situations the leaves grow much larger than those of the annexed specimen.



Antirrhinum Cymbalaria.

ANTIRRHINUM ELATINE. SHARP-POINTED FLUELLIN.

ANTIRRHINUM *Lin. Gen. Pl. DIDYNAMIA ANGIOSPERMIA.*

Raii Syn. Gen. 18. HERBÆ FRUCTU SICCO SINGULARI, FLORE MONOPETALO IRREGULARI.

ANTIRRHINUM foliis hastatis alternis, caulibus procumbentibus. *Lin. Sp. Pl. 85.*

ANTIRRHINUM caule procumbente, foliis hastatis, imis conjugatis, superioribus alternis. *Haller hist. v. 1.*

p. 14. 6. n. 340.

ELATINE folio acuminato, in basi auriculato, flore luteo. *Bauhin Pin. 253.*

ELATINE folio acuminato. *Parkinson 553.*

ELATINE altera. *Gerard emac. 623.*

LINARIA *Elatine dicta*, folio acuminato. *Raii Syn. *282.*

ANTIRRHINUM *Elatine*. *Hudson Fl. Angl. p. 237. Scopoli Fl. Carniol. p. 444. Oeder. Fl. Dan. Ic. 426.*

TOTA PLANTA pilosa.

RADIX fibrosa, annua, albidâ,

CAULES numerosi, teretes, subramosi, in junioribus plantis suberecti, tandem procumbentes, ad duos pedes et ultra sæpe extensi.

FOLIA petiolata, ima subrotunda, opposita; proxima dentata, alterna; quæ sequuntur magna ex parte hastata.

PEDUNCULI axillares, alterni, penduli, longitudine foliorum.

CALYX: PERIANTHIUM quinquepartitum, persistens, segmentis ovato-lanceolatis acutis, fig. 1.

COROLLA monopetala, ringens, flava; TUBUS brevissimus; LIMBUS bilabiatus, labium superius bifidum, segmentis obtusis, inferne purpureis, inferius trifidum, segmentis obtusis, medio productiore, et paulo minore; PALATUM prominulum, flavum, fig. 2; NECTARIUM subulatum, flavum, longitudine segmentorum calycis, fig. 3.

STAMINA: FILAMENTA quatuor, quorum duo paulo longiora; ANTHERÆ purpureo-fuscæ, coalescentes, fig. 4.

PISTILLUM: GERMEN subrotundum, compressum, apice villosum; STYLUS filiformis, longitudine staminum, apice incrassatus, uncinatus; STIGMA simplex, fig. 5, 6, 7.

PERICARPIUM: CAPSULA rotunda, bilocularis, bivalvis, valvis deciduis, foramine magno in utroque latere capsulæ relicto, valvæ orbiculatæ, concavæ, fig. 8, 9, 10.

SEMINA nigra, rugosâ, 8-10 in singulo loculamento, fig. 12.

THE WHOLE PLANT hairy.

ROOT fibrous, annual, whitish.

STALKS numerous, round, a little branched, in the young plants nearly upright, in the old ones trailing on the ground, frequently to the distance of two feet or more.

LEAVES standing on foot-stalks, the bottom leaves roundish and opposite, the next to those are indented and alternate, and those which follow are for the most part hastate.

PEDUNCLES alternate, pendulous, the length of, and proceeding from the AxE of the leaves.

CALYX: a PERIANTHIUM divided into five segments persisting, the segments lanceolate, fig. 1.

COROLLA monopetalous, ringent, and yellow; the TUBE very short; the LIMB divided into two lips, the upper lip bifid, the segments obtuse, and purple underneath; the lower lip trifid, the segments obtuse, the middle one longest and least; the PALATE prominent and yellow, fig. 2; the NECTARIUM the length of the segments of the Calyx, small and tapering, fig. 3.

STAMINA four FILAMENTS, two of which are a little longer than the others; the ANTHERÆ purplish-brown, adhering together, fig. 4.

PISTILLUM: the GERMEN roundish, flattened, at top hairy; the STYLE filiform, the length of the Stamina, thickened at top and hooked; the STIGMA simple, fig. 5, 6, 7.


SEED-VESSEL: a round CAPSULE of two cavities and two valves, the valves round and concave, on falling off leaving a large hole in each side of the Capsule, fig. 8, 9, 10.

SEEDS black, and wrinkled, from 8 to 10 in each cavity, fig. 12.

THIS species of *Antirrhinum* grows generally in Corn-fields, and in some parts of England is much more common than it is with us; in the Corn-fields about Peckham I have generally found it in bloom in July, August and September, and even later; it very much resembles the *Antirrhinum spurium* in its general habit, but is readily distinguished by its pointed leaves. Some Writers have considered it as possessed of healing properties, and affirm that the expressed juice of the plant, or its distilled water taken inwardly and applied externally, has checked and cured spreading and cancerous Ulcers; and RAY relates a Story from LOBEL of a poor Barber, who by the above use of this plant, saved his Nose, which had been condemned to be cut off by several eminent Physicians and Surgeons.



Antirrhinum Elatine.



ANTIRRHINUM LINARIA. COMMON YELLOW TOAD FLAX.

ANTIRRHINUM *Linnaei Gen. Pl.* DIDYNAMIA ANGIOSPERMIA.

Raii Syn. Gen. 18. HERBÆ FRUCTU SICCO SINGULARI FLORE MONOPETALO.

ANTIRRHINUM *Linaria* foliis lanceolato-linearibus confertis, caule erecto, spicis terminalibus seffilibus, floribus imbricatis. *Linn. Syst. Vegetab. p.* 466. *Fl. Suecic.* 217.

ANTIRRHINUM foliis linearibus adscendentibus congestis, caule erecto spicato. *Haller. hist. V. 1. p.* 145.

LINARIA vulgaris lutea flore majore. *Baubin pin. p.* 212.

LINARIA lutea vulgaris. *Gerard emac.* 550. vulgaris nostras. *Parkinson* 458. *Raii Syn. p.* *281. *Hudson Fl. Angl. p.* 238. *Scopoli Fl. Carniol. p.* 442:

RADIX perennis, alba, dura, lignosa, per terram rep-
tando immensum se propagans.

CAULES plerumque plures ex eadem radice, erecti,
pedales aut cubitales, foliosissimi, teretes, laeves.

FOLIA linearia, acuta, conferta, sparsa, glauca.

FLORES lutei, palato croceo, in summis caulibus in
spicas densas imbricatum congesti.

CALYX: PERIANTHIUM quinquepartitum, breve, per-
sistens, laciniis ovato-lanceolatis, superiore cæte-
ris paulo longiore, duabus inferioribus magis
dehiscentibus, *fig.* 1.

COROLLA monopetala ringens, lutea, *fig.* 3. TUBUS
brevis; LIMBUS bilabiatus, *fig.* 4. LABIUM
superius bifidum, laciniis primum deflexis, postea
reflexis conniventibus, *fig.* 5; LABIUM inferius
trifidum, laciniis obtusis, intermedio brevior
minore, *fig.* 6; FAUX clausa PALATO pro-
minente, bifido, croceo, ad basin villosa, *fig.* 7.

STAMINA: FILAMENTA quatuor, alba, sub labio su-
periori inclusa, quorum duo breviora, ad basin
villosa, *fig.* 9; ANTHERÆ flavæ, conniventes,
fig. 10.

PISTILLUM: GERMEN subrotundum, STYLUS filiformis,
albus; STIGMA obtusum.

PERICARPIUM: CAPSULA ovato-cylindræa, bilocu-
laris, apice in plures lacinas dehiscens, *fig.* 14,
15, 16.

SEMINA numerosa, nigra, plana, medio extuberantia,
fig. 17.

ROOT perennial, white, hard and woody, creeping
under the earth, and propagating itself very
much.

STALKS: generally several arise from the same root,
upright, from one to two feet high, very full
of leaves, round and smooth.

LEAVES linear, pointed, growing very thick together
on the stalk without any regular order, smooth,
and of a blueish colour.

FLOWERS yellow, with the palate of an orange or
saffron colour, placed one over another in thick
spikes on the top of the Stalks.

CALYX: a PERIANTHIUM divided into five segments
short and continuing, the segments oval and
pointed, the upper one a little longer than the
rest, the two inferior ones gaping widest, *fig.* 1.

COROLLA monopetalous, ringent, and yellow, *fig.* 3;
the TUBE short; the LIMB composed of two
LIPS, *fig.* 4; the upper LIP bifid, the segments
first bending down, afterwards turned back and
closing together, *fig.* 5; the lower LIP trifid,
the segments obtuse, the middle one shortest
and least, *fig.* 6; the MOUTH closed by a PA-
LATE prominent, bifid, of a saffron colour,
and villous at bottom, *fig.* 7.

STAMINA: four white FILAMENTS, inclosed under
the upper lip of the Corolla, two of which are
shorter than the other two, at bottom villous,
fig. 9; ANTHERÆ yellow, slightly connected
together, *fig.* 10.

PISTILLUM: GERMEN roundish, STYLE filiform and
white; STIGMATA obtuse.

SEED-VESSEL: a CAPSULE of an oval and cylindrical
shape, having two cavities, and splitting at top
into several divisions, *fig.* 14, 15, 16.


SEEDS numerous, black, flat, protuberant in the mid-
dle, *fig.* 17.

Mr. Ray in his *Historia Plantarum* has collected the Authorities of several writers who speak highly of the medical virtues of this Plant. At the same time that we by no means believe in all the Virtues which are attributed to many plants by the old Authors, we would be careful of rejecting all their accounts, particularly when there is some reason to think they may be founded in Truth, the mention of them may at least serve to excite such of the Faculty as have proper opportunities to give them a fair trial, and either reject them entirely, or bring them more generally into practice.


According to some it operates both by Stool and Urine, and so much by the latter, as to acquire among the Germans the name of *Harnkraut*. A small Glass of the distilled Water mixed with a drachm of the bark of the *Ebulus* or *Water Elder* in powder, powerfully provokes Urine, and is recommended in Dropsical Cases. The distilled water or juice of the Plant put in the Eyes, takes away the redness and inflammation of them, as *TRAGUS* asserts from his own long observation and experience. Made into an Ointment with lard and mixed with the yolk of Egg, it takes away the violent pain arising from the Piles.

The flowers of this plant are frequently found double with two or more Spurs, and a singular variety of it which *LINNEUS* calls *Peloria*, is said by Mr. HUDSON to grow about *Clapham* in *Surry*, this rare monstrosity we shall not fail to figure.

In its common state, the Toad Flax grows very common on banks by the road sides, which it decorates not a little by its singular and beautiful Flowers. It may with the greatest ease be cultivated in Gardens, and raised either from Seeds or Roots; the Seed is ripe at the latter end of September.



Antirrhinum 'Linaria'.



DIGITALIS PURPUREA. FOX-GLOVE.

DIGITALIS *Linnaei Gen. Pl.* DIDYNAMIA ANGIOSPERMIA.

Cal. 5-partitus. Cor. campanulata 5-fida, ventricosa. Caps. ovata bilocularis.

Raii Syn. Gen. 18. HERBÆ FRUCTU SICCO SINGULARI FLORE MONOPETALO.

DIGITALIS *purpurea* calycinis foliolis ovatis acutis, corollis obtusis : labio superiore integro. *Linnaei Syst. Vegetab. p. 570. Sp. Pl. p. 866.*

DIGITALIS foliis calycinis ovatis, galea simplici. *Haller. hist. p. 143. n. 330.*

DIGITALIS *purpurea. Scopoli Fl. Carniol. p. 447. n. 780.*

DIGITALIS *purpurea* folio aspero. *p. 243.*

DIGITALIS *purpurea. Gerard. emac. 790.*

DIGITALIS *purpurea vulgaris. Parkinson 1653. Raii. Syn. p.* 283. Purple Fox-glove. Hudson Fl. Angl. p. 240. Oeder Fl. Dan Icon. 774.*

RADIX biennis, fibrosa.

CAULIS tripedalis ad orgyalem, simplex, erectus, foliosus, teres, pubescens.

FOLIA ovato-acuta, ferrata, venosa, subtus albidia, pubescentia ; PETIOLI breves, alati.

FLORES spicati, nutantes, imbricati, secundi.

PEDUNCULI uniflori, pubescentes, apice incrassati, peracta florescentia suberecti.

CALYX : PERIANTHIUM quinquepartitum, lacinias ovato-acuminatis, nervosis, supremâ angustiore, *fig. 1.*

COROLLA monopetala, subcampanulata, purpurea, interne ocellata ; TUBUS magnus, patens, deorsum ventricosus, basi cylindracea, arcuata ; LIMBUS parvus, quadrifidus, lacinia superiore integra, quasi truncata, inferiore majore, inflexa.

STAMINA : FILAMENTA quatuor basi Corollae inserta, alba, apice paululum latiora, basi infracta, quorum duo longiora ; ANTHERÆ primum magnae, turgidae, ovatae, basi coadunatae, lutescentes, et saepe maculatae ; demum et forma et situ mire mutantur, *fig. 2, 3, 4.*

PISTILLUM : GERMEN subconicum, luteo-virens ; STYLUS simplex ; STIGMA bifidum, *fig. 5, 6, 7.*

NECTARIUM GLANDULA basin Germinis cingens, *fig. 8.*

PERICARPIMUM : CAPSULA ovato-acuminata, bilocularis, bivalvis, valvula inferiore fidente, *fig. 9.*

SEMINA plurima, nigricantia, parva, utraque extremitate truncata, *fig. 10.*

ROOT biennial and fibrous.

STALK from three to six feet high, simple, upright, leafy, round, and pubescent or downy.

LEAVES of a painted oval shape, serrated, veiny, underneath whitish and pubescent ; the FOOT-STALKS short and winged.

FLOWERS growing in a spike, pendulous, laying one over another all one way.

PEDUNCLES sustaining one flower, pubescent, thick-set at top, after the flower drops off, becoming nearly upright.

CALYX : a PERIANTHIUM divided into five segments, which are of an oval pointed shape, and nervous, the uppermost narrower than the rest, *fig. 1.*

COROLLA monopetalous, somewhat bell-shaped, purple, and marked in the inside with little eyes ; the TUBE large, spreading, bulging out backwards ; the base cylindrical, and as if it had been tied with a ligature ; the LIMB small and quadrifid, the upper segment entire and as if cut off, the lower segment larger and bent in.

STAMINA : four FILAMENTS inserted into the bottom of the Corolla, white, a little broadest at top, crooked at bottom, two long and two short ; ANTHERÆ at first large, turgid, oval, touching at bottom, of a yellowish colour and often spotted ; lastly changing both their form and situation in a singular manner, *fig. 2, 3, 4.*

PISTILLUM : GERMEN rather conical, of a yellow green colour ; STYLE simple ; STIGMA bifid, *fig. 5, 6, 7.*

NECTARY a GLAND surrounding the bottom of the Germen, *fig. 8.*

SEED-VESSEL : a pointed oval CAPSULE, of two cavities and two valves, the lowermost valve splitting in two, *fig. 9.*

SEEDS numerous, blackish, small, as if cut off at each end, *fig. 10.*

Was it not that we are too apt to treat with neglect the beautiful plants of our own country, merely because they are common and easily obtained, the stately and elegant *Fox-glove* would much oftener be the pride of our gardens than it is at present ; for it is not only peculiarly striking at a distance, but its flowers and their several parts become beautiful in proportion to the nearness of our view : How singularly and how regularly do the blossoms hang one over another ! How delicate are the little spots which ornament the inside of the flower ! and like the wings of some of our small Butterflies smile at every attempt of the Painter to do them justice : how pleasing is it to behold the netting Bee hide itself in its pendulous blossoms ! while extracting its sweets which furnish our tables with honey, and our manufacturers with wax : nor are the more interior parts of the flower less worthy of our admiration, or less adapted to the improvement of the young Botanist : here all the parts of the fructification being large, he will readily obtain a distinct idea of them ; but more particularly of the form of the Antheræ ; and the alteration which takes place in them, previous to and after the discharge of the Pollen. *vid. fig. 3, 4.*


The flowers of this plant are in general of a fine purple colour, and like all other purple flowers are liable to variations ; sometimes we find the blossoms of a milk white or cream colour, and some other varieties of it are mentioned by RAY, but the white is the most common. Such as would wish to cultivate it, may raise it either from seed, which is very small for the size of the plant, or from young plants. It grows naturally in a dry and gravelly soil, and in such situations is common enough over most parts of England ; about *Charlton-Wood* it is very plentiful, and flowers in July and August.

According to the testimony of many writers, the juice or decoction of this plant taken inwardly, acts as an emetic and purgative, and that too with considerable violence ; hence Mr. RAY very properly advises it to be given to such only as have robust constitutions. PARKINSON affirms that it is very efficacious in the cure of the Epilepsy ; but he unites with it in his prescription *Polygody of the Oak*, so that there is no knowing to which of the plants the merit of curing this stubborn disease is due.

The flowers or herb either bruised or made into an ointment, are strongly recommended in Schrophulous tumours and ulcers ; and so great an opinion have the Italians of its virtues as a vulnerary, that they have the following proverb concerning it. "*Arada tutte le piaghe salda.*" Fox-glove cures all wounds. *Raii Hist. Plant.*



Digitalis purpurea.



DRABA VERNA. VERNAL DRABA OR WHITLOW GRASS.

DRABA *Linnaei Gen. Pl. TETRADYNAMIA SILICULOSA.*

Raii Synop. Gen. 21. HERBÆ TETRAPETALÆ SILIQUOSÆ ET SILICULOSÆ.

DRABA *verna* scapis nudis, foliis subferratis. *Linnaei Syll. Vegetab. p. 489. Flor. Suec. p. 223.*

DRABA cauliculis nudis, foliis subhirsutis, subdentatis. *Haller. hist. helv. 1. 215.*

BURSA PASTORIS minor loculo oblongo. *Baubin. pin. 108. 2.*

PARONYCHIA vulgaris. *Gerard emac. 624. Raii Syn. 292. Hudson Fl. Angl. 243. Scopoli Flor. Carniol. n. 792.*

RADIX fibrosa, annua.

CAULES nudi, palmares, 1 ad 5 aut plures in solo fertili ex eadem radice nascuntur.

FOLIA ovato-lanceolata, basi angustiora integra et subferrata, (ferra nisi unica aut duo, raro plures) super terram expansa, scabriuscula, hirsuta, pili bi-trifurci.

PEDUNCULI alterni, uniflori.

CALYX: PERIANTHIUM tetraphyllum, foliolis erectis, concavis, gibbis, obtusis, subhirsutis. *fig. 1.*

COROLLA tetrapetala, petala alba, calyce duplo longiora, bipartita. *fig. 2.*

STAMINA: FILAMENTA sex incurvata, quorum 4 longitudine Pistilli 2 breviora; ANTHERÆ flavæ. *fig. 3. 4.*

PISTILLUM: GERMEN ovatum, compressum; STYLUS vix ullus; STIGMA capitatum, planum. *fig. 5.*

PERICARPIUM: SILICULA ovata, compressa, brevi mucrone obtuso terminata, bilocularis, bivalvis, valvulis plano-concavis. *fig. 6.*

SEMINA plura, ovata, fusca, margini DISSEPIMENTI affixa. *fig. 8. 9.*

† ROOT fibrous and annual.

STALKS naked, about three inches high, one to five and frequently more, if the soil be rich, spring from the same root.

LEAVES of an oval pointed shape, narrower at bottom, some of them entire, and others a little serrated, or indented, (seldom more than one or two indentations in a leaf,) spreading on the ground, roughish, hirsute, some of the hairs bifurcate, others trifurcate.

PEDUNCLES alternate, uniflorous.

CALYX: a PERIANTHIUM of four leaves, which are upright, hollow, gibbous, obtuse, and somewhat hairy, *fig. 1.*

COROLLA tetrapetalous, the PETALS white, twice the length of the Calyx, and bipartite, *fig. 2.*

STAMINA: six FILAMENTS which bend inward, 4 long the height of the Pistillum, and 2 short; the ANTHERÆ yellow, *fig. 3. 4.*

PISTILLUM: The GERMEN oval and flat; STYLE scarce any; STIGMA a small head flat at top.

SEED-VESSEL a short oval pod, flat, and terminated by a short blunt point, having two Cavities and two Valves, the Valves slightly concave, *fig. 6:*


SEEDS several, oval, brown, fixed to the edge of the DISSEPIMENT or Partition, *fig. 8. 9.*

ON Walls, dry Banks, and in barren Fields, the white blossoms of this diminutive plant, are very conspicuous in the months of March and April, a season when any kind of blossom is viewed with pleasure, as it cannot fail to excite the pleasing reflection that the season is approaching when


“ All that is sweet to smell, all that can charm
Or eye or ear, bursts forth on every side
And crouds upon the senses.”

Linnaeus informs us that in Smoland a Province of Sweden, they sow their Rye when this plant is in blossom, and that in the night time and in wet weather its flowers droop.

Galen says that *Paronychia* or *Whitlow Grass* has its name from its properties, for it heals Whitlows; but Commentators are much in doubt concerning the plant itself. From the account of the Antients, it appears that it is a different plant from what we are now describing; some have fixed on Wall Rue, (*Asplenium Ruta muraria*,) others on a plant resembling Spurge, such is the confusion that arises from imperfect descriptions.



Arabis verna





Thlaspi Bursa Pastoris

THLASPI BURSA PASTORIS. SHEPHERD'S PURSE.

THLASPI *Linnaei Gen. Pl. TETRADYNAMIA SILICULOSA.*

Silicula emarginata, obcordata, polysperma: valvulis navicularibus, marginato-carinatis.

Raii Syn. Gen. 21. HERBÆ TETRAPETALÆ SILIQUOSÆ ET SILICULOSÆ.

THLASPI *Bursa pastoris* siliculis obcordatis, foliis radicalibus pinnatifidis. *Linnaei Syst. Vegetab. p. 491.*

Spec. Pl. 903. Fl. Succic. 227.

NASTURTIUM siliquis triangularibus, *Haller hist. v. 1. p. 221*

PASTORIA BURSA *Fuschii icon. 611.*

BURSA PASTORIS major folio sinuato. *Baubin Pin. 108. Gerard emac. 276. Parkinsoni Theat. 866.*

Raii Syn. 306. Hudson, Fl. Angl. 247. Scopoli. Fl. Carniol. v. 2. 17.

RADIX annua, fibrosa, albida.

ROOT annual, fibrous and whitish.

CAULIS pedalis, erectus, ramifusus, teres, subasper.

STALK about a foot high, upright, branched, round, a little prickly.

FOLIA *radicalia* hirsutula, pinnatifida, laciniiis quoad formam mire variantibus, *caulina* amplexicaulia, dentata.

LEAVES: *radical leaves* slightly hirsute, pinnatifid, the laciniae or jags varying exceedingly in their form; the upper leaves embracing the stalk, and indented at the edges.

PEDUNCULI uniflori, demum fere horizontales,

PEDUNCLES, supporting one flower on each, nearly horizontal when the flowers are gone off.

CALYX: PERIANTHIUM tetraphyllum, foliolis ovatis, concavis, subpilosis, margine membranaceis, *fig. 1.*

CALYX: a PERIANTHIUM of four leaves, the leaves oval, hollow, slightly hairy, and membranous at the edges, *fig. 1.*

COROLLA: PETALA quatuor alba, calyce paulo longiora, apice rotundata, *fig. 2.*

COROLLA: four white PETALS, a little longer than the Calyx, round at top, *fig. 2.*

STAMINA: FILAMENTA sex, alba, quorum quatuor longitudine Styli, duo breviora incurvata; ANTHERÆ flavæ, *fig. 3.*

STAMINA: six white FILAMENTS, four of which are of the same length as the Style; two are shorter and bent a little inwards: ANTHERÆ yellow, *fig. 3.*

PISTILLUM: GERMEN oblongo-cordatum; STYLUS brevissimus; STIGMA villosum, *fig. 4.*

PISTILLUM: GERMEN of an oblong heart-shape; STYLE very short; STIGMA villous, *fig. 4.*

PERICARPIUM: SILICULA lævis, obcordata, bivalvis, *fig. 5.*

SEED-VESSEL; a short smooth pod, triangular or heart-shaped, with two valves, *fig. 6.*

SEMINA plurima, pedicellata, flavescencia, margini Dissepimenti affixa, *fig. 6.*

SEEDS numerous, of a yellowish colour, standing on little foot-stalks, which connects them to the edge of the Dissepimentum or Partition, *fig. 6.*

DISSEPIMENTUM utrinque acutum Valvis contrarium.


PARTITION pointed at both ends, placed cross-ways to the Valves.

THE radical leaves of this plant differ so exceedingly in their appearance, that the most expert Botanist is often obliged to have recourse to its most striking character, the shape of its Seed-vessels, before he can with certainty distinguish it. When it grows on walls and in dry situations, the leaves are more deeply divided, and the Laciniae become much narrower; in cultivated ground they are broader and less jagged: It differs likewise no less with respect to its size, sometimes being not more than two or three inches high, and at other times as many feet.

March and *April* are the months in which it is found most generally in blossom, yet like the *Groundsel* and *Poa annua*, it may be found in this state at almost any time of the year,

It acquires its name of *Shepherd's Pouch* or *Purse*, from the particular shape of its pods, by which it is obviously distinguished from all our other Tetrastynamous plants.

The plant is collected and given to small birds, who appear to be very fond of the seeds, and this is the only use to which we at present know of its being applied.



GERANIUM CICUTARIUM. HEMLOCK-LEAV'D CRANE'S-BILL.

GERANIUM *Linnei Gen. Pl.* MONADELPHIA DECANDRIA.

Monogyna. *Stigmata* quinque. *Fructus* rostratus, pentacoccus.

Raii Synop. HEREE PENTAPETALÆ VASCULIFERÆ.

GERANIUM *cicutarium* pedunculis multifloris, floribus pentandris, foliis pinnatis incisii obtusis, caule ramoso. *Liame. Soff. Vegetab. p. 90. Fl. Succic. p. 243.*

GERANIUM petiolis multifloris, caule procumbente, foliis duplicato-pinnatis, pinnulis acute incisii. *Hall. h³. No. 944.*

GERANIUM cicutæ folio minus, et supinum. *Baubin p.n. 319.*

GERANIUM cicutæ folio inodorum album. *Gerard emac. 945. 946.*

GERANIUM moschatum inodorum. *Parkinson 1708. Raii Syn. 357. Field Crane's-bill without scent. Hudson Fl. Angl. 262.*

RADIX annua, albida, simplex, carne tenera, cum nervo intus duriore et tenaciore, paucis fibris instructa, crassiuscula, et in terram alte descendens.

CAULES ex eadem radice nascuntur plures, crassiusculi, teretes, hirsuti, procumbentes, ramosi, variae longitudinis pro ratione loci.

FOLIA pinnata, pinnis sessilibus pubescentibus, pinnulis acute incisii.

STIPULÆ ad exortum foliorum membranaceæ, albide, ovato-acutæ, superiore integra, fig. 1; inferiore in duas divisæ, fig. 2.

PEDUNCULI axillares, alterni, hirsuti, multiflori, longitudine foliorum.

FLORES umbellati, rosei, a tribus ad sex.

INVOLUCRUM membranaceum, multidentatum, fig. 3; PEDICELLI basi crassiores, deflexi et demum assurgentes.

CALYX: PERIANTHIUM pentaphyllum, foliolis ovatis, striatis, hirsutis, concavis, mucronatis, fig. 4.

COROLLA: PETALA quinque, subovata, plana, subæqualia, rosea, basi hirsuta, calyce longiora, fig. 5.

STAMINA: FILAMENTA decem, quorum quinque alterna Antheris carent fig. 7; ANTHERÆ saturate purpurascens, fig. 6.

NECTARIA: Glandulæ quinque fuscae circa basin staminum locantur, fig. 9.

PISTILLUM: GERMEN quinquangulare, villosum; STYLUS subulatus, fuscatus; STIGMATA quinque purpurascens, paululum reflexa, fig. 10, 11.

PERICARPIUM nullum; FRUCTUS pentacoccus, rostratus.

SEMEN oblongum, læve, fuscum, arillatum, fig. 14; ARILLA hirsuta; ARISTA prælonga pilosa instructa quæ demum spiralis evadit, fig. 12, 13.

ROOT annual, whitish, simple, tender, the string or nerve in the middle of it hard and tough, furnished with few fibres, large for the size of the plant, and penetrating deep into the earth.

STALKS: several usually spring from the same root, thickish, round, hirsute, procumbent and branched, of various lengths according to their place of growth.

LEAVES pinnated, the pinnæ sessile and slightly hairy, the pinnulæ sharply indented.

STIPULÆ at the base of the leaves membranous, whitish, acutely oval, the uppermost entire, fig. 1; the lowermost generally divided into two, fig. 2.

FOOT-STALKS of the flowers springing from the base of the leaves, alternate, hirsute, the length of the leaves, and supporting many flowers.

FLOWERS growing in an Umbell, from three to six, of a rose-colour.

INVOLUCRUM membranous, with many teeth, fig. 3; the small foot-stalks of the flowers thickest at bottom, turning down, and lastly turning upward.

CALYX: a PERIANTHIUM of five leaves, the folioli oval, striated, hirsute, concave, and terminating in a fine point, fig. 4.

COROLLA: five PETALS, somewhat oval, flat, nearly equal, of a rose colour, hairy at bottom, somewhat longer than the calyx, fig. 5.

STAMINA: ten FILAMENTS, five of which want the Antheræ, the ANTHERÆ of a deep purple colour, fig. 6.

NECTARIA: five brown Glands placed round the base of the Stamina, fig. 9.

PISTILLUM: GERMEN quinquangular and villous, STYLE tapering and grooved; STIGMATA five, of a purple colour, bending a little back, fig. 10, 11.

SEED-VESSEL none; FRUIT as yet unripe, formed of five protuberating seeds, and terminating in a long beak.

SEED oblong, smooth, brown, inclosed within an ARILLUS, fig. 14, which is hirsute, and furnished with a long hairy ARISTA, finally becoming spiral, fig. 12, 13.


We have often had occasion to remark the very great difference in the appearance of plants arising from soil and situation; of this the young Botanist cannot be too well apprised, nor too often informed: from a want of attention to this circumstance, the plant which we have now described, has been divided by different Authors into several species.


It seems worthy of notice, that the alterations which are produced in plants from growing in a richer soil, are chiefly those of encrease of size, and a multiplication of their parts; the minutiae of the fructification suffer but little change in their form by culture, hence they are often most to be depended on, even in ascertaining different species.

When the *Geranium Cicutarium* grows on a dry sandy bank, or wall, as it very frequently does, it is quite diminutive; when it occurs in a moister and more luxuriant soil, the branches extend often a foot or two in length, and the whole plant becomes so altered in its general appearance, as readily to deceive the inexperienced Tyro; but the long pointed fruit which occurs in both, and from whence this plant has obtained the name of *Crane's-bill*, readily points them out to be the same.

The seeds of the *Geraniums* are, in general, enclosed within a membranous *Arillus*, which terminates in an *Arista* or *Tail*, of different lengths in different species; in some of them, when the seeds are become ripe, they detach themselves from the receptacle, to which they are affixed, with considerable elasticity, and the seeds being loosely contained within the *Arillus*, are thrown out to a considerable distance. In the present species, the seeds are more closely invested by the *Arillus*, which does not separate itself with so much force, and as soon as detached, the *Arista* begins to be twisted up in a spiral form. This may be very distinctly observed if we separate a seed, with its *Arillus*, as soon as ripe, and place it in the palm of the hand, the tail of the *Arillus* immediately appears in motion, as if endued with some sensitive property, and continues uninterruptedly this motion, 'till it has assumed the form of a screw, *vid. fig. 13*. The seed thus furnished with its twisted *Arista*, is more liable to attach itself to anything which may come in contact with it, by which means this plant is more universally disseminated.

The *Geranium moschatum* has a great affinity with this species, that plant however has a strong smell of musk, which this entirely wants; and has also many other peculiarities, which we shall not fail to particularize when it comes to be treated of.





GERANIUM ROBERTIANUM. STRONG-SCENTED CRANES-BILL, OR HERB ROBERT.

GERANIUM *Linnaei Gen. Pl.* MONADELPHIA DECANDRIA.

Stigmata quinque. Fructus rostratus, pentacoccus.

Raii Syn. 335. HERBÆ PENTAPETALÆ VASCULIFERÆ.

GERANIUM *robertianum* pedunculis bifloris, calycibus pilosis decemangulatis. *Linnaei Syst. Vegetab. p. 515. Fl. Suecic. 241. n. 619.*

GERANIUM *foliis duplicato pinnatis, pinnis ultimis confluentibus, calycibus striatis, hirsutis. Haller hijst. n. 943.*

GERANIUM *robertianum. Scopoli Fl. Carniol. n. 845. Hudson Fl. Angl. p. 264.*

GERANIUM *robertianum primum. Baubin. Pin. 319.*

GERANIUM *robertianum. Gerard. emac. 939.*

GERANIUM *robertianum vulgare. Parkinson 710. Raii Syn. p. 338.*

RADIX annua, fusca, fibris ramosis prælongis instructa.	ROOT annual, brown, furnished with long branched fibres.
CAULES plures, diffusi, ramosi, sanguinei ut ut tota planta haud infrequenter, geniculis tumidis, pilosi, præsertim in junioribus plantis.	STALKS several, spreading, branched, of a blood-red colour, as is frequently the whole plant, (the joints tumid,) hairy, particularly in the young plants.
FOLIA opposita, pilosa, præcipue in umbrosis, unumquodque folium e tribus foliolis pinnatifidis basi confluentibus componitur, foliolo medio longius pedicellato, lacinii spinula rubra terminatis.	LEAVES opposite, hairy especially when growing in the shade, each composed of three pinnatifid leaves, uniting at the base, the middle leaf standing on the longest foot-stalk, the laciniae or jags of the leaf terminated by a small red spine.
STIPULÆ ad singulum geniculum quatuor, utrinque binæ.	STIPULÆ four at each joint, two on each side of it.
PEDUNCULI biflori.	PEDUNCLES biflorous.
CALYX: PERIANTHIUM decemangulatum, persistens, foliis ovato-lanceolatis, nervosis, hirsutis, mucronatis, fig. 1, 2.	CALYX: a PERIANTHIUM having ten angles, and continuing, the leaves ovato-lanceolate, nervous, hairy, terminating in a point, fig. 1, 2.
COROLLA: PETALA quinque rosea, patentia, æqualia, lamina subcordata, unguis linearis, medio prominulo sulcato in tres nervos albos divaricante. fig. 3.	COROLLA: five rose-coloured PETALS, spreading and equal, the lamina somewhat heart-shaped, the claw linear, the middle part of it prominent, grooved, and spreading into three whitish nerves.
STAMINA: FILAMENTA decem fertilia, subulata, plana, alba, basi coherentia; ANTHERÆ purpurascentes, polline flavo repletæ, fig. 4, auct 5.	STAMINA: ten fertile FILAMENTS, tapering, flat, white, connected at bottom; ANTHERÆ purplish, filled with a yellow Pollen, fig. 4, magnified, fig. 5.
PISTILLUM: GERMEN quinquangulare; STYLUS subulatus, villosus; STIGMATA quinque, rubra, paululum reflexa, fig. 6.	PISTILLUM: GERMEN having five angles; STYLE tapering, villous; STIGMATA five, red, a little turned back, fig. 6.
SEMINA quinque Arillata, lævia, ovata, fusca ad unum latus compressa, fig. 9; ARILLUS rugosus, fig. 7, 8.	SEEDS five, contained within an Arillus, smooth, oval, brown, flattened on one side, fig. 9; the ARILLUS wrinkled, fig. 7, 8.


Although our English *Geraniums* cannot boast that grandeur and variety of splendid colours so conspicuous in many of the foreign ones, yet several of them are sufficiently beautiful to be entitled to a place in the gardens of the curious, particularly the *Bloody Cranes-bill*, (*Geranium Sanguineum*;) the *Crowsfoot Cranes-bill*, (*Geranium Pratense*;) the *Perennial Doves-foot Cranes-bill*, (*Geranium Perenne* of *Hudson*;) and the *Herb Robert* which we have now described: the latter of these grows naturally in woods, but more particularly under the hedges which surround woods; it likewise is frequently found in old hollow trees, and not uncommonly on the roofs of houses not much exposed to the sun: it is an annual plant; the seeds sow themselves in Autumn, soon after the young plants come up; flower the ensuing spring, and continue to blossom the whole Summer long, if the plant grows in the shade: towards the latter end of the year, both stalks and leaves become of a deep red or blood colour.

The whole plant has a disagreeable smell when bruised, by which it will be distinguished from our other species. It appears to grow all over Europe, and as a proof of its being still more universal, LINNÆUS mentions its growing in *Arabia felix*.


A variety with a white flower now and then occurs.

If credit may be given to writers on the *Materia Medica*, it is a plant of considerable efficacy in medicine, particularly as an Astringent, hence it is recommended in all kinds of Hemorrhages; and those who have the management of cattle, are said to give them an infusion of this plant when they make bloody urine.—Has not this practice originated from the doctrine of signatures? It is also celebrated as a vulnerary in schrophulous, cancerous and putrid Ulcers, to which either the juice is applied, or the parts fomented with a decoction of the herb; as likewise in Contusions, dissolving the extravasated blood when applied in the form of a Cataplasm; and lastly it is said to be exhibited with good success in the Stone and Gravel.—How far it merits these encomiums future experiments must determine.

The herb bruised and applied to places infested with Bugs, is said by LINNÆUS to drive them away.



Geranium robertianum.



OROBUS TUBEROSUS. WOOD PEA.

OROBUS *Linnei Gen. Pl.* DIADELPHIA DECANDRIA.

Raii Synop. Gen. 23. HERBÆ FLORE PAPILIONACEO, SEU LEGUMINOSÆ.

OROBUS *tuberosus* foliis pinnatis, lanceolatis; stipulis semifagittatis integerrimis, caule simplici. *Lin. Syst. Vegetab. p.* 550. *Fl. Suecic. n.* 642.

OROBUS caule simplici; foliis senis ellipticis; radice tuberosa. *Haller. hist. n.* 417.

ASTRAGALUS sylvaticus, foliis oblongis glabris. *Baubin. pin.* 351. *Gerard. emac.* 1237.

LATHYRUS sylvestris lignosior. *Parkinson, 1072.* *Raii Synop. p.* 324. Wood-Peasc, or Heath-Peasc. *Hudson, Fl. Angl. p.* 274. *Scopoli. Fl. Carn. n.* 883.

RADIX perennis, tuberosa.

CAULIS simplex, erectus, pedalis, alatus, subternatus.

FOLIA pinnata, CIRRHUS brevi recto terminata, Pinnarum paria duo, tria, elliptica, mucronata, glabra subtus cærulefcentia.

STIPULÆ semifagittatæ, sæpe integræ, sæpius vero ad basin hamatæ, dente unico aut pluribus.

RAMI florigeri, 1, 2, 3, aut plures ex foliorum alis, primum nutantes, FLORES pulchelli, ex rubro purpurei, demum cærulefcentes.

CALYX PERIANTHIUM monophyllum, tubulatum, purpureum, basi obtusum; ore quinque-dentato, denticulis tribus inferioribus acutioribus, duobus superioribus brevioribus, obtuse divisus, subafurgentibus, fig. 1.

COROLLA *Papilionacea*: VEXILLUM obcordatum, reflexum, fig. 2. ALÆ conniventes, Carinâ connexæ, Unguis linearis, fig. 5. Lamina obtusa. CARINA, fig. 7, acuminata, affurgens, marginibus cavis ad Alas recipiendas, fig. 9.

STAMINA: FILAMENTA diadelpchia (simplex et novem fidum) adscendentia, fig. 11, 17. ANTHERÆ flavæ, fig. 12. ad basin filamenti simplicis et superioris, foramina duo observantur, fig. 16.

PISTILLUM: GERMEN cylindraceum, compressum, Stylus filiformis, erectus, lateri interiori prope apicem villulosus, fig. 13.

PERICARPIMUM LEGUMEN teres, longum, primum rubrum, demum nigrum, fig. 14.

SEMINA plura, subrotunda, e luteo-fusca, fig. 15.

ROOT perennial and tuberosus.

STALK simple, upright, about a foot high, winged and somewhat twisted.

LEAVES pinnated, terminated by a short straight CIRRHUS consisting of two or three pair of Pinnæ which are elliptical, and end in a small sharp point, smooth and underneath blueish.

STIPULÆ semifagittate, frequently entire but more often jagged at bottom, with one or several teeth.

BRANCHES which sustain the flowers 1, 2, 3, or more, springing from the bosom of the leaves, at first drooping. the FLOWERS beautiful, of a reddish purple colour, becoming blue as they go off.

CALYX: a PERIANTHIUM of one leaf, tubular, purple, blunt at bottom, the mouth quinque-dentate, the three lowermost teeth sharpest, the two uppermost shortest, bluntly divided, and turned a little upwards, fig. 1.

COROLLA *Papilionaceous*: the VEXILLUM heart-shaped, turning back, fig. 2. the WINGS connivent and connected with the Carina, the Claw linear, fig. 5. the Lamina obtuse, fig. 6. the CARINA or Keel acuminate, rising upward, the edges hollow for the reception of the Alæ or Wings, fig. 9.

STAMINA: ten FILAMENTS, nine united into one body below, and one separate at top, fig. 11, 17. rising upward, ANTHERÆ yellow, fig. 12. at the base of the simple and uppermost filament two small holes are conspicuous, fig. 16.

PISTILLUM: GERMEN cylindrical, and flattish, STYLE thread-shaped, interiorly near the tip villous, fig. 13.

SEED-VESSEL, a LEGUMEN round, and long, first red, when ripe black, fig. 14.

SEEDS several, roundish, of a yellowish brown colour, fig. 15.


This elegant species of Orobus grows very plentifully in all our Woods about Town; it seems to delight in a strong clayey soil. It produces its blossoms in May and June and the seed is ripe in July. The root is large and tuberosus, deeply situated in the Earth and taken up with difficulty; it is not made any particular use of with us, but is considerably esteemed in some parts of Great Britain:


My very worthy and ingenious Friend the Rev. Mr. Lightfoot, of Uxbridge, has favoured me with the following account of its use, which he observed in his late tour through Scotland:

“The *Orobus tuberosus* is very common in Scotland, both in the Lowlands, Highlands, and the Hebrides. It is called “in the Erse Language *Cor-meille*. The Highlanders dig up the Roots and dry them in their pockets, and chew “them like Tobacco or Liquorice Root, to relish their Liquor, and to repel Hunger and Thirst. In *Breadalbane* “and *Ross-shire* they sometimes steep them in Water, and make an agreeable fermented Liquor with them, which “they esteem to be good for Disorders of the Thorax. It has a sweetish Taste somewhat like Liquorice Roots. Fond “as the Highlanders were of this Root they frequently used to change it with me for some Pig-tail Tobacco, their “favourite Indulgence.”









ERVUM HIRSUTUM. ROUGH-PODDED TINE-TARE.

ERVUM *Linnei Gen. Pl.* DIADÉLPHIA DECANDRIA. Calyx quinquepartitus, longitudine corollæ.

Raii Gen. 23. HERBÆ FLORE PAPILIONACEO SEU LEGUMINOSÆ.

ERVUM *hirsutum*, pedunculis multifloris, seminibus globosis binis. *Linn. Syst. Vegetab. p. 554. Spec. Plant. 1039. Fl. Suecic. 255.*

VICIA foliis linearibus, siliquis racemosis, dispermis, hirsutis. *Haller bist. helv. n. 422.*

ERVUM *hirsutum*. *Scopoli Fl. Carniol. n. 901. Hudson Fl. Angl. p. 280.*

VICIA segetum cum siliquis plurimis hirsutis. *Baubin. Pin. p. 345.*

VICIA sylvestris feu Cracca minima. *Gerard. emac. 1028.*


ARACHUS five Cracca minor. *Parkinson 1070. Raii Syn. small wild Tare or Tine Tare. Muller. Flor.*

Dan. icon. 639.

RADIX annua, tenuis, prælonga, paucis fibrillis instructa.	ROOT annual, slender, long, and furnished with few fibres.
CAULES pedales, aut bipedales, debiles, ramosi, quadrangulares, tortuosi.	STALKS from one to two feet high, weak, branch- ed, quadrangular and twisted.
STIPULÆ in plures lacinias tenues divisæ, superiore majore.	STIPULÆ divided into many slender lacinia, of which the uppermost is the largest.
FOLIA pinnata, ad octo aut duodecem paria, opposita, aut subalterna, lævia, lanceolata, apice truncata, nervo medio in macronem educto, capreolo ramofo terminata.	LEAVES pinnated, from eight to twelve pair, opposite, or nearly alternate, smooth, lanceolate, with the top cut off, and the midrib running out to a short point, terminated by a branched tendril.
PEDUNCULI longitudine foliorum, multiflori.	PEDUNCLES the length of the leaves, and supporting many flowers.
FLORES a tribus ad octo, pallide purpurei, racemati, et imbricatum dispositi.	FLOWERS from three to eight, of a pale purple colour, disposed in racemi, and laying one over another.
CALYX: PERIANTHIUM quinquedentatum, persistens, longitudine fere Corollæ, dentibus linearibus, acuminatis, subæqualibus, duobus superioribus more Orobi obtuse divis, fig. 1.	CALYX: a PERIANTHIUM with five teeth, continuing, almost the length of the Corolla, the teeth linear, and pointed, nearly equal, the two upper ones obtusely divided in the manner of the Orobus, fig. 1.
COROLLA papilionacea; VEXILLUM subrotundum, vix emarginatum, parum reflexum, fig. 2; ALÆ Carinæ adherentes, ovata, obtuse, ad basin lineares, fig. 3; CARINA alis brevior, fig. 4; interne macula purpurea utrinque notata.	COROLLA papilionaceous; the VEXILLUM roundish, scarcely nicked in, bending a little back, fig. 2; the WINGS adhering to the Carina, oval, obtuse, at bottom linear, fig. 3; the CARINA shorter than the Wings, fig. 4, marked internally on each side with a purple spot.
STAMINA: FILAMENTA decem affurgentia, supremum brevior cæteris, nec liberum, fig. 5; ANTHERÆ simplices, flavæ.	STAMINA: ten FILAMENTS which rise upward, the uppermost connected with, and shorter than the others, fig. 5; the ANTHERÆ simple and yellow.
PISTILLUM: GERMEN oblongum, STYLUS simplex, affurgens, STIGMA obtusum, villosum, fig. 6.	PISTILLUM: GERMEN oblong, STYLE simple and rising upward, STIGMA blunt and villous, fig. 6.
PERICARPIUM: LEGUMEN breve, hirsutum, dispermum, fig. 7.	SEED - VESSEL a short hairy LEGUMEN with two seeds, fig. 7.
SEMINA duo, subrotunda.	SEEDS two, and roundish.

This species of Tine-Tare, which at first sight bears so great a resemblance to the *Ervum tetraspermum*, grows like that, too frequently among Corn, to which it is in general more destructive, as being a stronger and more prolific plant. I have in wet seasons seen whole fields of corn overpowered and wholly destroyed by this plant.

It is easily distinguished from the *Tetraspermum*; in the first place, the leaves are not pointed as in that species, but appear as if cut off at the end, which although a material circumstance is not noticed by MULLER in his figure of it, *vid. Fl. Dan. icon. 639*; secondly the Stipulæ are divided into many more lacinia; the flowers and consequently the Pods grow in a kind of Cluster, whereas there is seldom more than two grow together in the *Tetraspermum*; and lastly, which seems to be the best distinction, the Pods are rough and contain two Seeds in each, while in the *Tetraspermum*, they are smooth and contain four Seeds.



ERVUM TETRASPERMUM. SMOOTH-PODDED TINE TARE.

ERVUM *Linnei Gen. Pl.* DIADELPHIA DECANDRIA.

Raii Syn. Gen. 23. HERBÆ FLORE PAPILIONACEO SEU LEGUMINOSÆ.

ERVUM (*tetraspermum*) pedunculis subbifloris, seminibus globosis quaternis. *Linn. Syst. Vegetab. p.* 554.

VICIA foliis linearibus, filiquis gemellis glabris. *Haller hist. v. 1. p.* 184.

ERVUM *tetraspermum*. *Scopoli Fl. Carniol. DIAGN.* Pedunculi subbiflori. Siliqua glabra, obtusa, tetrasperma.

VICIA segetum singularibus filiquis glabris. *Baubin Pin. p.* 345.

VICLÆ five Craccae minimæ species cum filiquis glabris. *I. Baubin.*

CRACCA minor filiquis singularibus, flosculis cæruleiscentibus. *Hoff. C. H. Alt. Raii Syn. p.* 322. Tine-

Tare with smooth pods. *Hudson Fl. Angl. p.* 280. *Oeder Fl. Dan. Icon.* 95.

RADIX annua, fibrosa.

CAULES in apertis locis læves, tenues, debiles, inter segetes vero, (ubi sæpius invenitur) capreolis erecte sese sustentant, pedales et ultra.

STIPULÆ ad basin foliorum, duo, simplices, utrinque acuminatæ.

FOLIA pinnata, lævia, lanceolata-linearia, parium trium ad quinque usque, capreolo ramofo terminata.

PEDUNCULI longitudine foliorum, plerumque biflori.

CALYX PERIANTHIUM quinquedentatum, persistens, dentibus inæqualibus, acutis, duobus superioribus brevioribus, latioribus, sursum tendentibus, obtuse divisis, *fig. 1.*

COROLLA papilionacea, *fig. 2*; VEXILLUM subemarginatum, limbus reflexus, venis purpureis pictus, *fig. 4*; ALÆ albe, conniventes, *fig. 5*; CARINA alis brevior, obtusa, *fig. 6.*

STAMINA: FILAMENTA diadelphea (simplex et novemfidum) adurgentia, *fig. 7*, 8, supremum liberum, *fig. 8*; ANTHERÆ simplices.

PISTILLUM: GERMEN compressum; STYLUS assurgens; STIGMA capitatum, villosum, *fig. 9.*

PERICARPIUM: LEGUMEN læve, teretiusculum, *tetraspermum*, *fig. 10.*

SEMINA subrotunda, fuscescentia, nigro marmorata, *fig. 11.*

ROOT annual and fibrous.

STALKS in open places are slender and weak, but among the Corn, (where this plant is most commonly found,) they support themselves upright by means of their tendrils, and grow to a foot or more in height.

STIPULÆ at the bottom of the leaves, two, simple, and pointed at each end.

LEAVES pinnated, smooth, lanceolate and linear, from three to five pair, terminated by a branched tendril.

PEDUNCLES the length of the leaves, generally sustaining two flowers.

CALYX a PERIANTHIUM having five teeth and continuing, the teeth unequal and pointed, the two uppermost shortest, broadest, and turning a little upwards, at bottom obtusely divided, *fig. 1.*

COROLLA papilionaceous, *fig. 2*; the VEXILLUM slightly nicked in at top, the limb somewhat turned back and streaked with purple, *fig. 4*; the ALÆ white and closing together, *fig. 5*; the CARINA shorter than the Alæ and obtuse *fig. 6.*

STAMINA: Ten FILAMENTS uniting into two bodies, of which one forms the lowermost, *fig. 7*, and one the uppermost which is free, *fig. 8*; ANTHERÆ simple.

PISTILLUM: GERMEN flatten'd; STYLE rising upward; STIGMA forming a little head and villos, *fig. 9.*

SEED-VESSEL: a LEGUMEN, smooth, roundish, and containing four seeds, *fig. 10.*

SEEDS nearly round, brownish and mottled with black, *fig. 11.*


This species of *Ervm* or *Tine-Tare* is found in most Corn-fields, often to the Farmers sorrow, as it frequently proves very injurious to the Corn, laying hold of it by means of its tendrils, and if the season favours its growth quite overcoming it. Like most plants of this kind it is exceedingly fertile; on one plant which I casually pulled up, I counted 220 pods, and as each pod contains four seeds, there must have been from a single seed the amazing produce of 880.

At first sight this species has a considerable resemblance to the *Ervm hirsutum*, but the slightest attention will discover the difference; in the *Ervm hirsutum* the pods contain only two seeds and are hairy; in the *Tetraspermum* they contain four and are smooth; in the *hirsutum* the flowers grow in a kind of cluster, in this species there is seldom more than two grow together.

The figure which I have given is intended to represent the plant as it grows among the Corn; when it is found by itself and in a poor soil it is often not so large.



Ervum tetraspermum





HYPERICUM PULCHRUM. Small Upright ST. JOHN'S WORT.

HYPERICUM *Linnaei Gen. Pl.* POLYADELPHIA POLYANDRIA.

Raii Synop. Gen. 24. HERBÆ PENTAPETALÆ VASCULIFERÆ.

HYPERICUM floribus trigynis; calycibus ferrato-glandulosis, caule tereti, foliis perfoliatis glabris. *Lin. Sp. Pl.* 1106.

HYPERICUM pulchrum Tragi. *J. Baubin. Hb. III.* 183. *Raii Synop.* 342.

HYPERICUM minus, erectum. *Baubin. Pin.* 279.

HYPERICUM foliis amplexicaulibus, cordatis, calycibus ovatis, ferratis, glanduligeris. *Haller. Hb. n.* 1041.

Gerard. emac. 540. *Hudson. Fl. Angl.* 290. *Oeder. Flor. Dan. Icon.* 75.

RADIX perennis.	ROOT perennial.
CAULIS pedalis ad bipedalem, erectus, teretis, fig. 1. glaber, subramosus, geniculi distantes.	STALK from one to two feet high, upright, round, fig. 1. smooth, and thinly branched, the joints remote from each other.
RAMI oppositi, breves, tenues, cauli similes.	BRANCHES opposite, short, slender and like the stalk.
PEDUNCULI teretes, plerumque triflori.	PEDUNCLES round, generally sustaining three flowers.
FOLIA CAULIS cordato-triangularia, glaberrima, amplexicaulia, saturate viridia, patentia, quam in cæteris Hypericis solidiora, versus marginem perforata, inferiora frequenter coccinea, RAMORUM ovata, caulis triplo minora, PEDUNCULORUM ovato-lanceolata.	LEAVES of the STALK triangularly heart-shaped, smooth, shining, embracing the stalk, nearly horizontal, of a deep green colour, more solid to the touch than the other St. John's Worts, perforated near the edge, and frequently of a bright red colour towards the bottom, those of the BRANCHES oval, three times smaller than those on the Stalk, and those of the PEDUNCLES lancet-shaped.
CALYX. PERIANTHIUM quinquepartitum, LACINIIS ovatis, acutis, striatis, margine ferratis, dentibus glanduliferis, glandulis nigro rufis, fig. 2.	CALYX. A PERIANTHIUM divided into five Segments, the SEGMENTS oval, pointed, striated, ferrated and, edged with little glands of a blackish red colour, fig. 2.
COROLLA. PETALA quinque, oblongo-ovata, flava, contorta, leviter striata, subtus aurantiaco lineata, margine subferrata et glandulis cincta, fig. 3.	COROLLA. Five PETALS, oblong, oval, yellow, slightly striated; on the under side tinged with a bright orange, slightly ferrated, and edged with glands, fig. 3.
STAMINA. FILAMENTA triginta sex, filiformia, in tres fasciculos ad basin coacta, in singulo fasciculo duodecim; ANTHERÆ biloculares, subrotundæ. POLLEN minuscule, fig. 4.	STAMINA. The FILAMENTS numerous, to thirty-six, filiform, uniting at bottom in three Fasciculi or Bundles, in each Fasciculus 12; the ANTHERÆ roundish and bilocular, fig. 4. The POLLEN bright scarlet.
PISTILLUM. GERMEN ovatum, STYLI tres, longitudine germinis, divaricantes, STIGMATA parva, subrotunda, fig. 5.	PISTILLUM. The GERMEN oval, three STYLES the length of the Germen, spreading, the STIGMATA small and roundish, fig. 5.
PERICARPIUM. CAPSULA subcônica, trilocularis, fusca, fig. 6, 7.	SEED-VESSEL a CAPSULE somewhat conical, of a brown colour, with three cavities, fig. 6, 7.
SEMINA plurima, oblonga, fusca, fig. 8.	SEEDS numerous, oblong and brown, fig. 8.

THE antient Botanists gave this Plant the name of PULCHRUM, from its beauty; and LINNÆUS has very properly continued it;—many will, no doubt, think it deserving of a place in their gardens, it is fond of a clayey soil, and woody situation, and is found in all the woods about town, as HORNSEY WOOD beyond *Uxington*, OAK OF HONOUR WOOD, (as it is generally called) a little beyond *Peckham*; CHARLTON-WOOD by *Greenwich*, likewise on *HOUNSLOW-HEATH*; it flowers in the month of *July*, and continues but a short time in blossom.

Its virtues as a medicine are probably the same with the common *St. John's Wort*.

THE UNIVERSITY OF CHICAGO


THE UNIVERSITY OF CHICAGO
THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO





Hypericum Perforatum

HYPERICUM PERFORATUM. COMMON St. JOHN'S WORT.

HYPERICUM *Linnaei Gen. Pl. POLYADELPHIA POLYANDRIA.*

Raii Synopsi Gen. 24. HERBE PENTAPETALE VASCULIFERE.

HYPERICUM *perforatum*, floribus trigynis, caule ancipiti, foliis obtusis pellucido-punctatis. *Linnaei Syst. Vegetab. p. 584. Fl. Suecic. n. 680.*

HYPERICUM caule terete, alato, ramosissimo; foliis ovatis, perforatis. *Haller. hist. vol. 2. p. 4.*

HYPERICUM vulgare *Bauhin. Pin. p. 279. Gerard. emac. 540. Parkinson 572. Raii Synop. 342. Hudson Fl. Angl. 290. Scopoli. Fl. Carniol. n. 944.*

Tota planta glandulis nigris adspersa.

The whole plant is sprinkled over with small black glands.

RADIX perennis, lignosa, fusca.

ROOT perennial, woody, of a brown colour.

CAULES plerumque plures ex eadem radice, bipedales, erecti, sublignosi, laeves, teretes, alternè ancipites fig. 1, ramosi.

STALKS several for the most part, springing from the same root, about two feet high, upright, woody, smooth, round, alternately two edged, fig. 1, much branched.

RAMI oppositi, suberecti, ancipites.

BRANCHES opposite, nearly upright, two edged.

FOLIA opposita, sessilia, ovato-oblonga, obtusa, perforata five pellucido-punctata, heptanervia ex luteo-viridia. fig. 2.

LEAVES opposite, sessile, of an oblong oval shape, obtuse, having the appearance of being all over perforated, of a yellowish green colour, with seven nerves or ribs, fig. 2.

PEDUNCULI ancipites, multiflori.

PEDUNCLES two edged, supporting many flowers.

PANICULA densa.

PANICLE bushy.

CALYX: PERIANTHIUM quinquepartitum, striatum, lacinis lanceolatis, acuminatis, nudis. fig. 3.

CALYX A PERIANTHIUM divided into five segments, and striated, the segments narrow and pointed, without any glands on them. fig. 3.

COROLLA: petala quinque, flava, ad unum latus crenulata, glandulis nigris adspersa. fig. 4.

COROLLA: five PETALS of a yellow colour, notched irregularly on one side, and sprinkled over with little black glands. fig. 4.

STAMINA: FILAMENTA plurima, in tria corpora vix coacta. fig. 5. ANTHERÆ flavæ, biloculares, loculis subrotundis, inter quos glandula nigra ponitur. fig. 6.

STAMINA: FILAMENTS numerous, uniting at bottom in three scarcely distinct bodies or fasciculi. fig. 5. ANTHERÆ yellow and bilocular, each of the Cavities of a roundish figure, and between them is situated a small black gland. fig. 6.

PISTILLUM: GERMEN subovatum, STYLI tres divaricantes: STIGMATA simplicia. fig. 7.

PISTILLUM: GERMEN somewhat oval, three STYLES which divaricate; the STIGMATA simple, fig. 7.

PERICARPIUM: CAPSULA subtrigona fig. 8. trilocularis fig. 9. pallide fusca.

SEED-VESSEL: a CAPSULE somewhat triangular, fig. 8, of a pale brown colour, with three Cavities, fig. 9.

RECEPTACULUM seu Thalamus seminum foramine triquetro gaudet, quod in pericarpio immaturo sectione transversa clare distingui potest, ut observavit Cl. Scopoli.

RECEPTACLE: the Receptacle which is continued through the Capsule, and connects the Cavities together, has a triangular hole in it, which is very obvious in a transverse section of it before it is ripe,—as the celebrated Scopoli has justly observed.

SEMINA plurima, oblonga, fusca. fig. 10. 11.

SEEDS numerous, oblong, and brown, fig. 10. 11.

It very often happens, that some of the minute parts of the Flower, and Seed, afford a more obvious, certain, and constant mark of specific difference, than any part of the plant besides, and we have a remarkable instance of the truth of this observation in the plant before us. A little gland, of a black colour, placed on the summit of the Anthera, at one view distinguishes this species, without any farther investigation: did such obvious distinctions prevail in all plants, a knowledge of them might with much ease be acquired; and fortunately we shall find, on examination, such marks more frequently occur than is generally imagined; whenever they do, we shall not fail to remark them.

The apparent perforation of the leaves, from whence this species is named, is not peculiar to it alone.

Although in the present practice this officinal plant does not seem to be much regarded, yet its sensible qualities, and the repeated testimonies of its virtues, entitle it as Dr. Cullen* observes to farther trials. To the taste it is astringent and bitter, and its effects seem to be chiefly diuretic. From possessing properties which have generally been called balsamic, it has been used as a vulnerary in external wounds, and internal hemorrhages, for the former purpose, the tops of the plant with the flowers are infused in oil, and for the latter, an infusion of the plant is made in the manner of Tea. It has likewise been given in ulcerations of the kidneys, and has even been supposed to possess virtues as a febrifuge.

It has had the ill fate to be abused by the superstition of the common people in France and Germany, who gather it with great ceremony on St. John's Day, and hang it in their Windows, as a certain charm and defence against Storms, Thunder, and evil Spirits; mistaking the meaning of some medical writers, who have fancifully given this plant the name of *Fuga Dæmonum* because they supposed, if given internally, it was a good medicine for maniacal and hypochondriacal Disorders.

The dried plant boiled with Allum dyes Wool of a yellow colour. It grows very common in hedges and fields that are but seldom tilled, and flowers in August and September.

*Vid. Dr. Cullen's Materia Medica p. 206.





Lactodon Tauriacum.

LEONTODON TARAXACUM. DANDELION.

LEONTODON *Linnaei Gen. Pl. SYNGENESIA. POLYGAMIA ÆQUALIS.*

Raii Synops. ed. 3. Gen. 6. HERBÆ FLORE COMPOSITO, NATURÆ PLENO LACTESCENTES.

LEONTODON *Taraxacum* calycis squamis inferne reflexis, foliis runcinatis denticulatis lævibus. *Linnaei Syll. Vegetab. p. 596. Sp. Plant 1122. Fl. Suec. 270.*

TARAXACUM calycibus glabris, squamis imis reflexis. *Haller hist. v. 1. p. 56.*

HEDYPSOIS *Taraxacum Scopoli Flor. Carn. n. 957.*

HEDYPSOIS major *Fuschii.*

DENS LEONIS latiore folio *Baubin. Pin. p. 126. Gerard. emac. 290. Parkinson 780. Raii Syn. ed. 3. p. 176. Hudson Fl. Angl. p. 297. Oeder. Fl. Dan. Icon. 574.*

RADIX perennis, subfusiformis, lactescens, externe pallide fusca.

FOLIA laciniato-pinnatifida, plus aut minus profunde incisa, lacinis acutis et acute dentatis, plerumque lævia, nonnunquam vero subsæpera.

SCAPI nudi, fistulosi, lacteicentes, versus apicem subtomentosi, uniflori.

CALYX communis lævis, glaucus, squamis inferioribus reflexis, fig. 1.

COROLLA composita, flava, corollulis hermaphroditis, numerosis, æqualibus. *Propria* monopetala, ligulata, truncata quinqueidentata, fig. 2.

STAMINA: FILAMENTA quinque capillaria, brevissima, fig. 3. ANTHERÆ flavae, in tubum cylindraceum coactæ, fig. 4.

PISTILLUM: GERMEN oblongum, fig. 5. STYLUS longitudine corollæ, fig. 6. STIGMATA duo revoluta, fig. 7.

SEMEN subincurvatum, subcompressum, subtrigonum, striatum, apice echinatum, pallide olivaceum, fig. 8, 9. PAPPIUS stipitatus, simplex, stipite brevior, fig. 10.

RECEPTACULUM nudum, alveolatum. fig. 11.

ROOT perennial, tapering, milky, externally of a pale brown colour.

LEAVES more or less deeply jagged, each jag or lacinia pointed, and sharply indented, generally smooth, but sometimes a little rough.

STALKS naked, hollow, milky, towards the top covered with a kind of down, supporting one flower on each.

CALYX: the common or general Calyx smooth, glaucous, the lowermost leaves or squamæ turning back, fig. 1.

COROLLA: the flower compounded of a great number of COROLLULÆ or lesser flowers, which are yellow, hermaphrodite and equal; each Corollula monopetalous, tubular at bottom, and flat towards the extremity, the apex truncated and quinqueidentate, fig. 2.

STAMINA: five FILAMENTS small and very short, fig. 3. the ANTHERÆ yellow, uniting and forming a cylindrical tube, fig. 4.

PISTILLUM: GERMEN oblong, fig. 5. STYLE the length of the COROLLA, fig. 6. STIGMATA two, rolling back, fig. 7.

SEED a little crooked, flattish, and somewhat four square, striated or grooved, at top prickly, of a pale olive colour, fig. 8, 9. the Down or pappus standing on a footstalk, simple, not feathery, shorter than the footstalk, fig. 10.

RECEPTACLE naked, and full of little holes, fig. 11.

As a medicinal plant the Dandelion is thought to possess considerable virtues, and has been frequently made use of in obstructions of the Viscera, particularly the Jaundice. Some recommend the juice, others a decoction of the whole plant. It appears to operate chiefly by urine, and from possessing this property in a considerable degree it has acquired its vulgar name of *Piss-a-bed*. Its other, and more common name, seems to be a corruption of the French term *Dent de Lion*.

As a kind of salad, this plant is by many preferred to any other, particularly by the inhabitants of Spitalfields, many of whom being descended from French families, that forsook their native country for one more favourable to religious liberty, still retain the peculiar customs of that people in their diet, &c. They blanch, or whiten it as the gardeners do endive, and the inferior class generally use the simple process of laying a tile on it, for whatever excludes the light from this or any other plant will make it become white, all plants deriving their colours from the fountain of light, the sun. And it is remarkable, that many plants containing bitter and acrid juices are rendered by this process mild, sweet, and agreeable: who, for instance, could eat endive, celery, or even lettuce, in their wild uncultivated states?

The Dandelion grows in the greatest plenty in rich meadows, although it is very common on walls, and in courts and areas; when growing in a barren soil or dry situation the leaves become more narrow and jagged.

It flowers in May, and is the first plant which covers our meadows with a beautiful yellow coat, a few weeks afterwards, when it produceth its seed, it changes this for a white one.

Children frequently amuse themselves with blowing off the seeds, which stand naked on the receptacle or top of the stalk, and the round white heads, formed by the expansion of their pappus or down, they call *clocks*.

The young botanist generally finds some difficulty in acquiring a clear idea of the structure of these compound flowers, occasioned by the minuteness of the parts of fructification, which however are much larger and more conspicuous in this than in many others of the class SYNGENESIA, and therefore a proper flower for him to begin with.

On examining the flower of the Dandelion he will find that it is not a double flower, properly so called, as he might be led to think from its fullness, but that it is composed of a great number of *Flosculi*, or lesser flowers placed close together on one common receptacle or bottom, and enclosed by one common or general calyx. On dissecting each of these *Flosculi*, he will find them to consist of a COROLLA, or PETAL fig. 2, which at bottom is tubular, but towards the extremity flat, that from the bottom or tubular part of the corolla, five FILAMENTS spring, which are small and short, yet loose and unconnected fig. 3, that these filaments are furnished with ANTHERÆ, which unite together and form a long slender tube fig. 4, beneath the corolla is placed the GERMEN, or future seed fig. 5, from whence the STYLE, or middle part of the Pistillum proceeds and passes up through the middle of the flower, betwixt the Filaments and through the tube formed by the union of the Antheræ, fig. 6, and is furnished at top with two STIGMATA which roll back, fig. 7, at a little distance from the Germen the lower part of the Stylus is surrounded by numerous upright hairs which are the future PAPPIUS or Down, fig. 10.

This, then, he will find to be the appearance of the parts of fructification in a full blown flower.


Those parts of the flower which were more immediately or more remotely necessary to the impregnation of the Seed having now performed their office decay, the Corolla with the Stamina and upper part of the Pistillum drops off, the Seed becomes larger, the lower part of the Pistillum remains, is elongated and becomes the footstalk of the Pappus, and the Seed as yet immature with the Pappus as yet moist are all enclosed and pressed by the Calyx into a conical form. This is its appearance in its second state.

The fructification still going forward the seed becomes ripe and brown, the PAPPIUS now deprived of its moisture expands itself every way, fig. 10, pushes back the Calyx, and assumes a spherical form. The seeds fitted for vegetation and thus exposed are carried away by the first strong wind, and "a new race planted far from their native soil."

Such then is the curious process which nature makes use of in the perfecting and dissemination of this plant.

1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900

1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900



LAPSANA COMMUNIS. NIPPLEWORT.

LAPSANA *Linnei Gen. Pl.* SYNGENESIA POLYGAMIA ÆQUALIS.

Receptaculum nudum. Cal. calyculatus, squamis fingulis interioribus calyculatis.

Raii Syn. Gen. 6. HERBÆ FLORE COMPOSITO NATURA PLENO LASTESCENTES.

LAPSANA *communis* calycibus fructus angulatis pedunculis tenuibus ramofissimis. *Linnei Syst. Vegetab. p. 602. Sp. pl. 1141. Fl. Suecic. p. 277.*

LAMPSANA caule brachiato; foliis ovatis longe petiolatis; petiolis pinnatis. *Haller hist. n. 6.*

LAMPSANA *communis. Scopoli Fl. Carniol. n. 988.*

SONCHO affinis Lampsana domestica. *C. Baubini pin. p. 124.*

LAMPSANA *Gerard. emac. 255.*

LAMPSANA vulgaris. *Parkinson 810. Raii Syn. 173. Hudson Fl. Angl. p. 303.*

RADIX annua, simplex, fibrosa.

CAULIS erectus, rigidus, bicubitalis, striatus, ramosus, hirsutus.

FOLIA opposita, hirsutula, ad radicem et in ima parte caulis uno vel altero pinnularum pari donata, segmento terminali magno, ovato, dentato, superiora oblonga, dentata.

CALYX: communis calyculatus, angulatus, lævis, squamæ ad basin minimæ erectæ, *fig. 1.*

COROLLA composita, imbricata, Corollulis hermaphroditis æqualibus; *propria* monopetala, ligulata, truncata quinque dentata, *fig. 2.*

STAMINA: FILAMENTA quinque capillaria, brevissima; ANTHERÆ cylindracea tubulosa, *fig. 2.*

PISTILLUM: GERMEN oblongiusculum; STYLUS filiformis, longitudine Staminum; STIGMA bifidum, reflexum, *fig. 2.*

SEMINA circiter octodecem, oblonga paululum incurvata, pappo destituta, intra calycem, *fig. 3, 4.*

ROOT annual, simple, and fibrous

STALK upright, rigid, about two cubits high, striated, branched, hairy.

LEAVES opposite, somewhat hairy, at the root, and on the lower part of the stalk furnished with one or two pair of pinnulæ; the segment which terminates the leaf large, oval, and indented; the upper leaves oblong and indented.

CALYX: the common Calyx smooth, and furnished at bottom with a few minute, upright squamulæ, *fig. 1.*

COROLLA compound, imbricated, the floscules hermaphrodite and equal; each of them monopetalous, ligulate, truncated, and having five teeth, *fig. 2.*

STAMINA: five small and very short FILAMENTS; ANTHERÆ uniting into a tube, *fig. 2.*

PISTILLUM: GERMEN oblong; STYLES filiform, the length of the Stamina: STIGMA bifid and turning back, *fig. 2.*


SEEDS about eighteen, oblong, a little bent in, without any down, contained within the Calyx, *fig. 3, 4.*

In gardens as a weed, this plant answers very well to the name of *communis*, being in general too common. Nature seems amply to have supplied the want of pappus or down in the seeds, by the great number of them produced in each plant. It also occurs on the sides of banks, and in all cultivated ground; flowering during most of the Summer months.

According to RAY it receives its name of *Nipplewort* from its efficacy in curing sore nipples: no other virtues or uses seem attributed to it.



Lapsana communis.





Erigeron 'aero'.

ERIGERON ACRE. PURPLE ERIGERON.

ERIGERON *Linnaei, Gen. Pl. SYNGENESIA POLYGAMIA SUPERFLUA.*

Raii Synopsi. HERBÆ FLORE COMPOSITO, SEMINE PAPPOSO NON LACTESCENTES, FLORE DISCOIDE.

ERIGERON *Acre pedunculis alternis unifloris. Lin. Sp. Pl. 1211.*

ERIGERON *polymorphum Scopoli. Fl. Carniol. DIAGN. folia lanceolata, basi et apice attenuata. Germina villosa. Pappus rufus.*

ERIGERON caule alterne ramoso, petiolis unifloris, semisfoculis pappum æquantibus, et semisfoculis pappum superantibus *Haller. hist. n. 85. 86.*

CONYZA cærulea acris *Baubin Pin. 265. Gerard emac. 484.*

ASTER arvensis cæruleus acris. *Raii Syn. 175. Blue-flowered sweet Fleabane.*

CONYZA odorata cærulea *Parkinson 126.*

SENECIO five Erigeron cæruleus *L. B. II. 1043 Hudfon Fl. Angl. 314. Oeder Fl. Dan. Tab. 292.*

RADIX perennis, fibrosa, fibris pallide fuscis.

* ROOT perennial and fibrous, the fibres of a pale brown colour.

CAULIS erectus, rigidus, pedalis, pupureus, striatus, foliosus, hirsutus, in quibusdam vix ramifusus in aliis ramifissimus.

* STALK upright, rigid, about a foot high, purple, striated, leafy, and hirsute, in some scarce branched at all, in others very much so.

FOLIA alterna, sessilia, hirsuta, inferiora obtuse ovata basi angustiora, superiora angusta, reflexa, tortuosa, ramorum linearia, subrecta.

* LEAVES alternate, sessile, hirsute, the bottom ones of a blunt oval shape, and narrow at bottom, the upper ones narrow, turning back and twisted, those of the branches linear and nearly upright.

FLORES erecti, nunquam sese explicantes sicut plerique flores Classis Syngenesiæ, externi purpurei, interni flavescentes, cum cavitate in medio.

* FLOWERS upright, never expanding themselves like most of the flowers of the Class Syngenesia, externally purple, internally yellow, with a cavity in the middle.

CALYX communis imbricatus, squamis subulatis, erectis, purpureis, hirsutis, laxis, fig. 1.

* CALYX: the common Calyx composed of a number of scales, which are narrow and pointed, upright, purplish, hirsute, and loosely connected fig. 1.

COROLLA composita, radiata; Corollulæ hermaphrodite tubulose, numerosæ in disco, fig. 2. femineæ ligulate, pauciores in radio, fig. 3. Propria hermaphroditi infundibuliformis, flava, limbo quinquefido, fig. 2: Femineæ ligulata, linearis, erecta, purpurea, hermaphroditi longior, fig. 3.

* COROLLA compound and radiated; the hermaphrodite flowers tubular and numerous in the middle, fig. 2. the female flowers ligulate, and fewer in the circumference, fig. 3: each hermaphrodite floscule, funnel-shaped, yellow, with the limb divided into five segments, fig. 2: each female floscule, linear, upright, purple, longer than the hermaphrodite flower, fig. 3.

STAMINA hermaphroditi: FILAMENTA quinque, capillaria, brevissima: ANTHERÆ in tubum coalite.

* STAMINA in the hermaphrodite flowers: five FILAMENTS, very small and short; the ANTHERÆ united into a tube.

PISTILLUM Hermaphroditi: GERMEN coronatum Pappo corolla paulo longior, fig. 4. STYLUS filiformis longitudine Pappi fig. 5; STIGMA bifidum fig. 6: Femineis: GERMEN tenue, Pappo longitudine fere Corollæ, fig. 7; STIGMATA duo, tenuissima, fig. 8.

* PISTILLUM of the hermaphrodite flowers; the GERMEN crowned with a Pappus or Down a little longer than the Corolla, fig. 4; the STYLE filiform, the length of the Pappus, fig. 5; STIGMA bifid, fig. 6: of the Female flowers; the GERMEN slender, the Pappus nearly the length of the Corolla, fig. 7; two STIGMATA very slender, fig. 8.

SEMINA oblonga, pallide fusca, hirsuta, lente aut: fig. 9: PAPPUS sessilis, lutescens, simplex, fig. 10.


* SEEDS oblong, of a pale brown colour, hirsute, magnified fig. 9; PAPPUS sessile, yellowish and simple fig. 10.

The Erigeron Acre is by no means a common plant in our neighbourhood, yet occurs very frequently on the hilly and chalky ground about *Charlton Wood*, particularly in the chalk pits on the left hand side of the lane behind the Church.

It flowers in the months of August and September, and is considered as a pretty sure indication of a barren soil. It has a taste somewhat warm and biting, and hence has received its name of *Acriis*.

We have rather chosen to retain LINNÆUS's name of *Erigeron* than adopt RAY's name of *Fleabane*, which tends to confound it with the Genus *Conyza*.

It frequently grows much taller, and is often found much smaller than the specimen we have figured.



SENECIO VULGARIS. GROUNDSEL.

SENECIO *Linnaei Gen. Pl.* SYNGENESIA POLYGAMIA SUPERFLUA. *Receptaculum nudum. Pappus simplex.*

Calyx cylindricus, calyculatus; squamis apice sphacelatis.

Raii Syn. HERBÆ FLORE COMPOSITO, SEMINE PAPPOSO NON LACTESCENTES, FLORE DISCOIDE.

SENECIO *vulgaris* corollis nudis, foliis pinnato-finuatis amplexicaulibus, floribus sparsis. *Linn. Syst. Vegetab.*

p. 630. Sp. Pl. 1216. Fl. Suecic. p. 290.

SENECIO corollis nudis, foliis pinnato-finuatis amplexicaulibus, floribus sparsis. *Haller bist. n. 58.*

SENECIO *vulgaris. Scopoli Fl. Carniol. p. 162. n. 1063. Hudson Fl. Angl. p. 315.*

SENECIO minor vulgaris. *Baubin Pin. 181.*

SENECIO *vulgaris. Parkinson 671.*

ERIGERON *Gerard. emac. 278. Raii Syn. p. 178. Common Groundsel or Simfon.*

RADIX annua, e plurimis fibrillis albidis constans.
CAULIS simplex, erectus, pedalis, ramosus, sæpe purpureus, subangulosus, in junioribus plantis versus apicem subtomentosus.

FOLIA obscure virentia, glabra, amplexicaulia, pinnato-finuata, pinnis acute dentatis.

PEDUNCULI striati, uniflori, primum erecti, perfecta florescentia penduli, demum erecti.

CALYX: communis primum cylindraceus, demum conicus; Squamis subulatis, plurimis, in cylindrum superne contractis parallelis, contiguis, æqualibus, paucioribus basin imbricatim tegentibus, apicibus omnium nigricantibus, fig. 1.

COROLLA Composita, longitudine calycis; Corollulæ hermaphrodite, tubulosæ, numerosæ in disco, infundibuliformes; limbo reflexo, quinquefido: Radio nullo, fig. 2, 3.

STAMINA: FILAMENTA quinque, capillaria, minima; ANTHERA cylindracea, tubulosa.

PISTILLUM: GERMEN ovatum; STYLUS filiformis, longitudine flaminum; STIGMATA duo oblonga, revoluta.

SEMEN oblongum, striatum, fuscum; PAPPUS simplex, albus, semine triplo fere longior, fig. 4; RECEPTACULUM nudum, scabrum.

ROOT annual, consisting of numerous white fibres. STALK single, upright, about a foot high, branched, often purple, slightly angular, in the young plants, towards the top, thinly covered with down.

LEAVES of a deep and dull green colour, smooth, embracing the stalk, pinnato-finuata, the pinnæ sharply indented.

PEDUNCLES striated, supporting one flower on each, at first upright, when the flowering is over they become pendulous, and lastly upright.

CALYX; the common Calyx first cylindrical and lastly conical; the Squamæ subulate, numerous, contracted above into a Cylinder, parallel, contiguous and equal; those at the base of the calyx fewer, lying one over another, the tips of all of them blackish, fig. 1.

COROLLA Compound, the length of the Calyx; the Florets hermaphrodite, tubular and numerous in the disk or middle, funnel-shaped, the limb reflex and divided into five segments: the Radius wanting, fig. 2, 3.

STAMINA: FILAMENTS five, capillary, and very minute: ANTHERÆ united into a tube.

PISTILLUM: GERMEN oval; STYLE filiform the length of the Stamina; STIGMATA two, oblong, and bent back

SEED oblong, striated and brown; the PAPPUS simple, white, almost three times the length of the seed, fig. 4; RECEPTACLE naked, and rough.

The Groundsel is a Plant which is well known to grow exceedingly common in Gardens, cultivated Ground, and on Walls, flowering all the year if the weather be mild.

Although it is scarcely used at present as a medicine, yet according to some Authors it is not without considerable virtues: the juice, or decoction of it taken internally, operates gently by vomit; and the plant externally applied, is said to be useful in inflamed Breasts, the Scrophula, and other Inflammations.


Mr. RAY suspects that it might be given with advantage in Worms, as Farriers and Horse-dealers give the juice of it to Horses that are troubled with those kind of Worms called *Bottes*, and to which it is presently fatal.

Birds of various kinds are fond of the seeds and tops of this plant; and a great variety of Caterpillars particularly those of the *Phalæna Jacobææ* eat it readily.



Senecio vulgaris.





BELLIS PERENNIS. COMMON DAISY.

- BELLIS. *Linnaei Gen. Pl.* SYNGENESIA POLYGAMIA SUPERFLUA.
Raii Synopsis Gen. 8. HERBE FLORE COMPOSITO DISCOIDE, SEMINIBUS PAPPO DESTITUTIS,
 CORYMBIFERE DICTÆ.
 BELLIS *perennis*, scapo nudo. *Linnaei System. Vegetab. p. 640. Fl. Suecic. p. 296. Haller hist. p. 39. Scopoli.*
Fl. Carniol. v. 2. 146.
 BELLIS *sylvestris minor* *Bauhin pin. 261. Gerard emac. 635. Parkinson 530. Raii Syn. p. 184. Hudson Fl.*
Angl. 320. Oeder. Fl. Dan. icon. 503.

- | | |
|--|---|
| <p>RADIX <i>perennis</i>, fibrosa.</p> <p>FOLIA ovata, dentata, hirsutula, in petiolo longos decurrentia; disrupsa fila trahentia.</p> <p>SCAPI teretes, hirsuti, triunciales, uniflori, ad apicem fistulosi.</p> <p>CALYX <i>communis simplex</i>, foliolis æqualibus <i>fig. 1.</i> apice membranaceis, hirsutis, obtusis <i>fig. 2. lente auct.</i></p> <p>COROLLA <i>composita</i>, radiata: <i>Corollulæ</i> hermaphrodite, tubulosæ, numerosæ in disco. <i>Feminae</i> ligulatae, calycis foliis plures in radio. Flosculi <i>Hermaphroditi</i> infundibuliformes quinquefidii flavi, <i>fig. 3, 4. lente auct.</i> <i>Feminae</i> ligulati, lanceolati, albi, <i>fig. 10.</i></p> <p>STAMINA <i>Hermaphroditis</i>: FILAMENTA quinque brevissima, <i>fig. 5.</i> ANTHERA cylindracea, tubulosa, <i>fig. 6.</i></p> <p>PISTILLUM <i>Hermaphroditis</i>: GERMEN ovatum, <i>fig. 9.</i> STYLUS filiformis, <i>fig. 8.</i> STIGMA crassiusculum, bifidum, <i>fig. 7.</i> <i>Feminae</i>: GERMEN ovatum, <i>fig. 13.</i> STYLUS filiformis. STIGMATA duo patula, linearia, <i>fig. 11.</i></p> <p>SEMINA ovata, compressa, marginata, pappo destituta, <i>fig. 14.</i></p> <p>RECEPTACULUM nudum, conicum, <i>fig. 15.</i></p> | <p>ROOT perennial, and fibrous.</p> <p>LEAVES oval, indented, slightly hirsute, running down the footstalks, which are long and if broke across appear stringy.</p> <p>STALKS round, hirsute, about three inches high, supporting one flower, at top hollow.</p> <p>CALYX the common calyx simple, the leaves equal, <i>fig. 1.</i> at the top membranous, hairy and obtuse, <i>fig. 2.</i> one of the tips magnified.</p> <p>COROLLA Compound and radiated: the <i>Corollulæ</i> or flosculi in the disk or middle numerous, tubular, and <i>hermaphrodite</i>, those in the radius or circumference flat, more numerous than the leaves of the calyx, and <i>female</i>. The <i>Hermaphrodite</i> Flosculi funnel shaped, divided into five segments and yellow, <i>fig. 3, 4.</i> magnified. The <i>Female</i> flosculi tubular at bottom, flat towards the extremity, lanceolate, and white, <i>fig. 13.</i></p> <p>STAMINA in the <i>Hermaphrodite</i> flower: five FILAMENTS very short, <i>fig. 5.</i> ANTHERÆ united into a tube, <i>fig. 6.</i></p> <p>PISTILLUM of the <i>Hermaphrodite</i> flower: GERMEN oval, <i>fig. 9.</i> STYLE thread-shaped, <i>fig. 8.</i> STIGMA thickish and bifid, <i>fig. 7.</i> of the <i>Female</i> flower: GERMEN oval, <i>fig. 9.</i> STYLE thread-shaped, two STIGMATA narrow and spreading, <i>fig. 11.</i></p> <p>SEEDS oval, flat, margin'd without any pappus or down, <i>fig. 14.</i></p> <p>RECEPTACLE naked and of a conical figure, <i>fig. 15:</i></p> |
|--|---|

The Daisy has been recommended by some writers to be given in hectic fevers, caused by drinking cold water when the blood has been heated by exercise, either infused in water or milk.

In some parts of Germany, it is said to be boiled and eaten with meat as a pot-herb; but it does not seem to promise much either as physic or food for man. Sheep and horses refuse it, and it is very probable that none of our cattle eat it willingly; if so the owners of lands pay dear for their enamelled meads, and daisied carpets, but this part of husbandry seems as yet little understood or attended to. As rural economists we have ventured to say thus much in dispraise of this flower, notwithstanding the lavish encomiums the father of our English poets has bestowed on it:

— In special one called Se of the daie
 The Daisie, a floure white and rede,
 And in french called *La bel Margarete*
 O commendable floure, &c. —
 — Above all flours in the mede
 Than love I most those flours white and rede.
 Such that men call in Daisies in our Town.

Chaucer is perhaps the first that takes notice of the Horologium Floræ or opening and shutting of flowers at a particular time of the day.

— She that is of all flours the floure,
 Fullfilled of all virtue and honour;
 And ever alike fair and fresh of hewe,
 As well in winter as in summer newe,
 As soon as ever the Sunne ginneth West
 To sene this floure, how it will go to rest,
 For fear of night so hateth the darknesse
 Her chere is plainly spread in the brightnesse
 of the Sunne. —
 Well by reason men it calle maie
 The Daisie, or else the Eye of the Daie
 And at the last there tho began anon
 A Lady for to sing right womanly
 A Bargonet in praising the Daisie
 For as methought among her notis swete
 She said *Si douce est la Margarete*

Retuned by Dryden in his own numbers:
 And then the Band of Flutes began to play,
 To which a Lady sung a Virelay;
 And still at every clofe she would repeat
 The Burden of the Song the *Daisy is so sweet*
The Daisy is so sweet when she begun
 The troops of Knights and Dames continued on
 The Comfort and the voice so charm'd my Ear
 And sooth'd my Soul that it was Heaven to hear.

Etymologists agree with the Old Bard in his derivation of the Daisy, viz. Days Eye. Under the French name *Margarette* it is probable a compliment was intended to some lady, but Critics are not agreed who this lady was.

Like many other flowers the Daisy becomes double by culture, and frequently *proliferous*, in this state it is called the *Hen and Chicken* Daisy.







Viola odorata.

VIOLA ODORATA. SWEET VIOLET.

VIOLA *Linnaei Gen. Pl.* SYNGENESIA MONOGAMIA.

Calyx pentaphyllus. *Corolla* pentapetala, irregularis, postice cornuta: *Capsula* fupera, trivalvis, unilocularis.

Raii Syn. Gen. 24. HERBÆ PENTAPETALÆ VASCULIFERÆ.

VIOLA *odorata* acaulis, foliis cordatis, stolonibus reptantibus, bracteis supra medium peduncul.

VIOLA *odoratâ*, acaulis, foliis cordatis, stolonibus reptantibus. *Linn. Syst. Vegetab.* 1: 668.

VIOLA acaulis stolonifera, foliis cordatis, *Haller hist. helv.* n. 558

VIOLA *odorata*, *Scopoli Fl. Carniol.* n. 1097.

VIOLA *maria* purpurea flore simplici odore. *Bauhin Pin.* p. 199: *maria* alba. p. 199.

VIOLA *nigra* five *purpurea*. *Ger. emac.* 550:

VIOLA *simplex* *maria*: *Parkinson 735*: *Raii Syn.* p. 364: Purple Sweet Violet, and White Sweet-scented Violet. *Oeder Fl. Dan. icon.* 309:

RADIX perennis, fibrosa, albida, in senescente plantâ basi petulorum quotannis relicta pars superior radicis tuberculosa evadit, et supra terram eminet; e sinu horum nodorum nascuntur stolones, qui humi repant, et foliis instruuntur stipulisque ejusdem formæ ac illæ quæ ad basin plantæ inveniuntur.

FOLIA subrotundo-cordata, crenata, superne glabra, inferne hirsutula, junioribus involutis.

STIPULÆ radicales, ovato-lanceolatae, membranaceæ, serratæ, dentibus glanduliferis.

PEDUNCULI radicales, infra Bractæas quadrangulares, supra Bractæas dorso canaliculati, apice incurvati, uniflori.

BRACTEÆ duæ, lanceolatae, plerumque oppositæ, appressæ, supra medium peduncul.

CALYX: PERIANTHIUM pentaphyllum, persistens, foliolis oblongo-ovatis, obtusis, e viridi purpurascens, fig. 1.

COROLLA pentapetala, irregularis, violacea, odorata, petalum infimum Nectario corniculato, obtusiusculo, apice compresso instructum, Petala lateralialia prope basin barbata, fig. 2.

STAMINA: FILAMENTA quinque brevissima ægre distinguenda: ANTHERÆ flavescens, biloculares, vix connexæ, membranâ ovato-acuta aurantiaca terminatæ; e parte posteriori duarum Antherarum exit Nectariumque intrat appendiculâ viridis, linearis, compressa, fig. 5, 4, 3.

PISTILLUM: GERMEN subrotundum; STYLUS basi tenuior et paululum tortuosus; STIGMA uncinatum, Antheris paulo longius, fig. 6, 7.

PERICARPIUM priusquam dehiscit, subrotundo-triangulari, purpurascens, villosum; trivalve valvulis subrotundis concavis, fig. 8.

SEMINA plurima, rotunda, nitida, straminea, appendiculata, fig. 9.

ROOT perennial, fibrous and whitish; in old plants the upper part of the root becomes knobby, and appears above ground, the knots or knobs being formed from the bottoms of the foot-stalks of the leaves which are yearly left; from the bottoms of these knobs spring the stolones or shoots which creep on the ground, and are furnished with leaves and the same kind of Stipulæ which are observable at the bottom of the plant.

LEAVES heart-shaped and somewhat round at the tip, crenated, on the upper side smooth and shining, underneath slightly hairy, when young rolled in at the edges.

STIPULÆ springing from the root, ovato-lanceolate, membranous, serrated at the edges, each serrature or tooth terminating in a minute gland.

PEDUNCLES springing from the root, below the Bractæ quadrangular, above the Bractæ grooved on the upper side, at top incurved, supporting one flower.

BRACTEÆ two, lanceolate, generally opposite to each other, pressed to the stalk, and placed above the middle of the Peduncle.

CALYX: a PERIANTHIUM of five leaves, continuing, each leaf of an oblong oval shape, obtuse at the tip, and of a greenish purple colour, fig. 1.

COROLLA: of five PETALS, irregular, of a bluish purple colour and sweet smell, the lowermost terminating in a blunt horned NECTARIUM, a little flattened at the extremity, the two side Petals bearded near the base, fig. 2.

STAMINA: five FILAMENTS so short as hardly to be distinguished; ANTHERÆ yellowish, bilocular, scarcely connected together, terminated by an oval-pointed, orange-coloured membrane; from the back of two of the Antheræ, springs a slender, flat, greenish appendage, which enters the Nectarium, fig. 5, 4, 3.

PISTILLUM: GERMEN roundish; STYLE slenderest at bottom and a little twisted; STIGMA hooked, and a little longer than the Antheræ, fig. 6, 7.

SEED-VESSEL, before it bursts, roundish, rather approaching to triangular, of a purplish colour, and villous appearance, splitting into three roundish hollow valves, fig. 8.

SEEDS several, round, shining, of a straw colour, terminated by a little appendage, fig. 9.

The *Viola odorata* delights to grow under warm hedges, particularly near Woods; if the Spring be favourable, it is generally in full bloom in the month of March, and towards the latter end of Summer ripens its seeds. A variety of this plant frequently occurs with a white flower, not inferior in its agreeable scent to the blue one; and sometimes this plant is found double, in which state it is often introduced into Gardens, and being furnished with abundance of creeping shoots, it is by means of these propagated with the utmost facility.

This species of Violet bears a considerable resemblance to the *Viola hirta*, the mode of distinguishing them we shall point out when we describe the latter.

A syrup made from the flowers is usually kept in the shop, and frequently given to children where a gentle laxative is required: it is likewise in use as a test to try acid and alkaline substances. The

The seeds are said by Authors to possess a diuretic quality, and hence the powder of them has been recommended in the stone and gravel.

The great BACON, who frequently descended from his sublimer studies, and amused himself with enquiries into the qualities and properties of plants, has left us a curious method of preserving the scent of this flower

"Take Violets, and infuse a good pugil in a quart of Vinegar, let them stand three quarters of an hour, and take them forth, and refresh the infusion with like quantity of Violets seven times; and it will make a Vinegar so fresh

"of the flower, as if a twelve month after it be brought you in a saucer, you shall smell it before it come at you."

"Note. It smelleth more perfectly of the flower a good while after than at the first."

The illustrious prescriber has given no directions concerning the use of this preparation, but it appears to us to be one of the most grateful preservatives against infection, especially if the strongest distilled vinegar which has been drawn over in glass, be made use of.

The Violet has been much complimented by the antient Poets; and our SHAKESPEARE gives it a conspicuous place in his catalogue of flowers.

"Violets dim,
"But sweeter than the lids of JUNO's eyes,
"Or CYTHEREA's breath."

The Commentators have not been successful in informing us how the "lids of JUNO's eyes" bear any resemblance to "Violets dim," not recollecting that *ιοβλαφαρος* (*having violet eyelids*) was a complimentary title with the Greek poets. This epithet alludes to a well known custom which still prevails in Greece, of colouring the eye lids blue. "A Grecian girl is painted blue round the eyes; and the insides of the sockets, with the edges on which the lashes grow, are tinged with black: For colouring the lashes and socket of the eye, they throw incense of Gum of Labdanum on some coals of fire, intercept the smoke which ascends, with a plate, and collect the foot: This I saw applied; a girl sitting cross-legged, as usual, on a sofa, and closing one of her eyes, took the two lashes between the fore finger and thumb of her left hand, pulling them forward, and then thrusting in, at the external corner, a bodkin which has been immersed in the foot, and extracting it again, the particles before adhering to it remained within, and were presently ranged round the organ, serving as a foil to its lustre, besides contributing, as they say, to its health, and increasing its apparent magnitude." CHANDLER *Travels into Greece*.


Altho' the poet of nature has been rather obscure on this subject, where he copies the ancients; he makes ample amends when he gives us the genuine effusions of his own imagination. With what precision and delicacy does he describe the soft enchantment of plaintive music, as resembling the sweetness of this flower illustrating in a beautiful simile the object of one sense by that of another!

"That strain again;—it had a dying fall;
"Oh! it came o'er my ear, like the sweet south,
"That breathes upon a bank of violets,
"Stealing and giving odour!"

*A Greek poet supposed to be a Christian, from the severity of his manners and purity of his instructions, forbids this custom of painting the eye lids, in the rules of conduct which he addresses to young women,

"Μὲν ἄλλοις τέτοιον οὐκ ἐπιφαίνεται σπουδῆς." NAUMACHIUS.

It is probable that the Greeks borrowed this fashion from their Asiatic neighbours; JEREMIEL, a native of Zidon, put her eye in painting, as translators tell us in the margin of our bible; the Prophets also allude to and censure this custom, see *Jeremiah* iv. 30. *Ezekiel* xxiii. 40.





Viola hirta.

VIOLA HIRTA. HAIRY VIOLET.

VIOLA *Linnei Gen. Pl.* SYNGENESIA MONOGAMIA.

Calyx pentaphyllus. *Corolla* pentapetala, irregularis, postice cornuta. *Capsula* supra, trivalvis, unilocularis.

Raii Synop. Gen. 24. HERBÆ PENTAPETALÆ VASCULIFERÆ.

VIOLA *hirta* acaulis, foliis petiolisque hirsutis, bractæis infra medium pedunculi.

VIOLA *hirta* acaulis, foliis cordatis piloso hirsutis. *Linn. Syst. Vegetab. p. 668.*

VIOLA acaulis, foliis cordatis hirsutis. *Haller hist. helv. n. 559.*

VIOLA *hirta* *Hudson Fl. Angl. p. 330.*

VIOLA *maria major* hirsuta inodora. *Hist. ox. II. 475.*

VIOLA *trachelii* folio vulgo. *Raii hist. 1051. Syn. p. 365.* Violet with Throat-wort leaves.

So great is the similarity betwixt this Species and the *Viola odorata*, that to describe it in the same manner as I have that plant, would be to repeat nearly the same words. To avoid this sameness of expression, I shall adopt a description in the way of contrast, which will enable me to point out the differences of each in a manner more striking, and I hope equally satisfactory to my botanic readers.

I would first premise, that as it is my greatest wish to clear up every difficulty respecting the species and varieties of those plants which come properly before me, so I have with that view, not only examined this plant with the greatest attention, where it has grown wild, but also cultivated it in my garden along with the *odorata*, and hence, seeing and noticing its mode of growth throughout the year, have perhaps been able to obtain a clearer idea of its history, than those who may have viewed it at one particular season only.

The *Viola odorata* throws out from the upper part of its root a number of stolones or shoots, which trail on the ground, and quickly take root at the joints, whereby it propagates itself very fast: the *hirta* likewise encreases itself by throwing out young stalks; but then they are not procumbent, nor do they ever strike root as those of the *odorata* do; hence the *hirta* does not encrease so fast, nor spread so wide. Although LINNÆUS makes a considerable difference in the form of the roots of these plants, yet from what I have observed, this difference proceeds chiefly from the age of the roots; for in both species, the older they are, the more full are they of tubercles or cicatrices, formed by the annual shedding of the leaves.

The radical *Stipule* are lanceolate and serrated in both species.

The *footstalks* of the leaves form perhaps the most obvious difference; in the *odorata* they are nearly smooth; in the *hirta* they are very hirsute, and this hairiness puts on a kind of silvery appearance in the young plants of this species, where it is remarkably conspicuous.


In the leaves themselves the difference is for the most part, not very remarkable, for in both species they are somewhat hirsute underneath; those of the *hirta* however, are sometimes remarkably so, from growing in particular soils or situations: the leaves of the *odorata* have a more glossy appearance on their upper surface, but this scarce discriminates them unless they are contrasted. With respect to shape and size likewise, the difference is not very obvious; both species when in bloom are small, compared with the size to which they afterwards grow. In make they are somewhat longer, and not so perfectly heart-shaped.

In the specimens of this plant which I have examined, I could not perceive that sensible difference which LINNÆUS notices (*vid. Maniff. Plant. ult. p. 483.*) in the shape of the *Peduncle* above the *Bractææ*; in both species they certainly are channelled at the back: in the situation of the *Bractææ*, however, there is a very considerable difference, which does not appear to have been taken notice of, and this seemed to me to be so obvious a character, that I trust it will apologize for my altering its specific description: in the *odorata*, the *Bractææ* are placed above the middle of the *Scapus*, or *Peduncle*; in the *hirta*, they are situate below it: but there is one caution necessary to be observed respecting this character, viz. that the *Bractææ* of each be observed, just when the flowers are fully expanded, for as that part of the *Scapus*, which is situated above the *Bractææ*, grows considerably longer by the time that the flowers of the *odorata* are faded, so they should both be examined when of an equal age, otherwise this distinction will not appear so remarkable.

The flowers of the *hirta*, in general, appear about a week later than those of the *odorata*, are of a paler blue colour, and entirely want that sweet fragrance which renders the *odorata* so grateful a harbinger of the Spring. In the other parts of the fructification, these plants are very similar to each other; but there is one circumstance respecting the manner in which they produce and disperse their seeds, which may not be generally known.

LINNÆUS in his *Flora Suecica*, n. 789, observes that the flowers which the *Viola mirabilis* first produces from the root, are furnished with Petals, yet that these for the most part are barren, while those which blow later the same Spring, and rise from the stalk, although destitute of Petals, produce perfect seed: and JACQUIN, in his excellent work the *Flora Austriaca*, where this plant is figured, (*Vol. 1. pl. 19.*) confirms the truth of LINNÆUS's observations, and says that the barrenness of those flowers appeared to arise from a deficiency of the *Stylus*. LINNÆUS in his valuable treatise above quoted, observes likewise, that the flowers of the *Viola montana*, which appear first, are furnished with Petals, but that those which are afterwards produced have no Petals, yet nevertheless are fertile; and this I find, on repeated examination, to be the case with the *Viola odorata* and *hirta*, but more particularly the latter: they differ from the *Viola mirabilis* in this respect, that all the flowers which are formed, both with and without Petals, produce perfect seed. I was led to this discovery from observing a single plant of the *Viola hirta*, to produce about the middle of Summer, ten or twelve capsules of ripe seeds, on which I was certain in the Spring no more than two or three blossoms had appeared: the next Spring I discovered, that besides those perfect blossoms which first spring up, this plant continues for a month or more to throw out new flowers, which are entirely destitute of Petals, or have only the rudiments of them which never appear beyond the *Calyx*; but all the other parts of the fructification are perfect. The capsules in both these species, when they become nearly ripe, lay close to the ground, so that when they burst, the seeds have an easy access into the earth.

There is some difference with respect to the soil and situation in which these two plants delight; the *odorata* grows very generally under warm hedges, and in woods; the other appears to be pretty much confined to a chalky soil, and often occurs in more exposed situations, in the fields and on the banks about *Chalfen*, it may be found in tolerable abundance.





Viola tricolor

VIOLA TRICOLOR. WILD PANSIE.

VIOLA *Linnaei* Gen. Pl. SYNGENESIA MONOGAMIA.

Raii Synop. Gen. 20. HERBÆ PENTAPETALÆ VASCULIFERÆ.

VIOLA tricolor, caule triquetro diffuso, foliis oblongis incisif, stipulis pinnatifidis. *Lim. Syst. Vegetab. p. 668. Fl. Suecic. 397.*

VIOLA caule diffuso, ramoso, foliis ovatis dentatis, flore calyce paulo majori. *Haller. hist. tom. 1. n. 569.*

VIOLA bicolor arvensis. *C. Bauhin. pin. 200.*

VIOLA tricolor sylvestris. *Parkinson. 755.*

JACEA bicolor frugum et hortorum vitium. *I. Bauhin. III. 548. Raii Syn. p. 336. 11. Hudfon. Fl. Angl. p. 331. Scopoli. Fl. Carniol. p. 183.*

RADIX simplex, fibrosa.

CAULIS palmaris et ultra, plerumque diffusus, ramofus, angulofus, ad basin fordidè purpureus; rami alterni.

FOLIA longe petiolata, elliptica, crenata, inferioribus sæpe minoribus, subrotundis, superioribus angustis, subdentatis.

STIPULÆ ad basin laciniato-pinnatifidæ, lacinii linearibus, extrema oblonga, dentata.

PEDUNCULI subquadrangulares, alterni, apice incurvati, dorso canaliculati, stipulis duobus parvis, membranaceis, prope florem, instructi.

CALYX : PERIANTHIUM pentaphyllum, persistens, foliolis acutis, tria *superiora* minora, ad basin æqualia, *suprema* erecta, petalis supremis longiora, duo *inferiora* apice et basi cæteris longiora, basique latiora, petalis infimis breviora, fig. 2.

COROLLA pentapetala, irregularis, duo superiora subrotunda, integerrima, albidæ, deorsum spectantia; lateraliū *lamina* ovata, obtusa, ad basin barbata, lineaque brevi purpurea notata; infimum latum emarginatum, ad basin flavum, lineis quinque purpureis pictum, CALCARE SEU NECTARIO NECTARIUM terminatum, longitudine calycis, apice violaceo, obtuso, fig. 3, 4, 5, 6.

STAMINA: FILAMENTA quinque, brevissima; ANTHERÆ albidæ, vix coadunatæ, biloculares, membranâ crocæ terminatæ, edubus inferioribus exeunt, nectariumque intrant, appendiculæ duæ lineares, fig. 7, 8, 9, 10.

PISTILLUM: GERMEN subconicum, fig. 11; STYLUS ad basin tortuosus, staminibus longior, fig. 12; STIGMA capitatum, oblique perforatum, persistens, fig. 13.

PERICARPIMUM: CAPSULA ovata, glabra, unilocularis, trivalvis, fig. 14, 15.

SEMINA plurima, ovata, fusca, nitida, appendiculata, valvis seriatiim affixa, fig. 15.

ROOT simple and fibrous.

STALK about four or six inches high, generally spreading, branched, angular, at bottom of a dull purple colour; the branches alternate.

LEAVES placed on long foot-stalks, elliptical, crenated, the lowermost often smaller and roundish, the uppermost narrow and slightly indented.

STIPULÆ at bottom jagged and pinnatifid, the lacinie or jags linear, that which terminates the Stipula oblong and indented.

FOOT-STALKS of the flowers, nearly quadrangular, alternate, bent in at top, channelled on the back, and furnished with two small membranous Stipulæ near the flower.

CALYX : a PERIANTHIUM of five leaves and continuing, the leaves sharply pointed, the three *upper ones* smallest, and equal at bottom, the uppermost upright and longer than the uppermost petals, the two *under leaves* longer both at bottom and top than the rest, and at bottom likewise broader, shorter than the lowermost petals, fig. 2.

COROLLA pentapetalous and irregular, the two uppermost petals roundish, entire, and reflected; the lamina or broad part of the five petals oval, obtuse, bearded at bottom, and marked with a short purple line; the lowermost petal broad, emarginate, yellow at bottom, and streaked with five purple lines, and terminated by a NECTARY SPUR of NECTARY the length of the Calyx, with a blueish and blunt point, fig. 3, 4, 5, 6.

STAMINA: five FILAMENTS very short; ANTHERÆ whitish, scarcely united, bilocular, terminated by a saffron coloured membrane; from the two lowermost two linear appendages go off and enter the Nectary, fig. 7, 8, 9, 10.

PISTILLUM: GERMEN somewhat conical, fig. 11; STYLE twisted at bottom and longer than the Stamina, fig. 12; STIGMA forming a little head, obliquely perforated and continuing, fig. 13.

SEED-VESSEL: an oval smooth CAPSULE of one cavity and three valves, fig. 14, 15.

SEEDS numerous, oval, brown and shining, with a button to each, affixed in rows to the inside of the valves, fig. 15.

Few plants have acquired a greater variety of names than the *Viola Tricolor*; in different Authors and different countries we find the following, viz. *Wild Pansie*, *Heb Tinity*, *Hea is ease*, *Three faces under a hood*, *Cull me to you*, *Love in Idleness*, &c. what has occasioned some of these is the different appearance it puts on from cultivation and change of soil; in a garden there are few flowers that can boast a greater variety or richness of colour, few that continue longer in blossom, or are cultivated with more ease; it is probable that the large yellow Violet, *Viola lutea*, is no more than a variety of this species.

The Pansie in its wild state occurs very frequently in cultivated fields, and blossoms through most of the summer months. It is so hardy as to appear in Lapland amongst the few other plants which ornament the wastes of that Country during its short summer. It is eaten by Kine and Goats.

The difference in the form of the Stigma seems to divide the plants of this Genus into two families, viz. *Pansies* and *Violets*, in the former the Stigma is round, with a remarkable hole on one side of it, in the latter it is hooked.

Linnaeus remarks the black lines which sometimes appear on the Petals, MILTON has observed the same, "*Pansies speak with 'fate'*" In a poor soil the purple and yellow in the bloom of this flower frequently become very faint, and sometimes fade into a perfect white, this variation in colour gives a propriety to the Metamorphosis of this flower in which SHAKESPEAR pays an elegant compliment to his royal mistress.

That very time I saw, (but thou could'st not)
Flying between the cold Moon and the Earth,
Cupid all-arm'd: a certain aim he took,
At a fair Vestal, thronged by the west,
And loo'd his love-shaft sinfully from his bow,
As it should pierce a hundred thousand hearts:
But I might see young Cupid's fiery shaft
Quench'd in the chaste beams of the watery moon,
And the imperial votress pass'd on,
In maiden meditation fancey-free.
Yet mark'd I where the bolt of Cupid fell,
It fell upon a little western flower;
Before, milk-white; now purple with Love's wound,
And Maidens call it Love in Idleness.

210111

1. The first of the three main parts of the book is a history of the development of the theory of the structure of the atom.

2. The second part is a description of the experimental methods used to determine the structure of the atom.

3. The third part is a discussion of the results of the experiments and the conclusions drawn from them.

4. The fourth part is a summary of the results of the experiments and the conclusions drawn from them.

5. The fifth part is a summary of the results of the experiments and the conclusions drawn from them.

6. The sixth part is a summary of the results of the experiments and the conclusions drawn from them.

7. The seventh part is a summary of the results of the experiments and the conclusions drawn from them.

8. The eighth part is a summary of the results of the experiments and the conclusions drawn from them.

OPHRYS APIFERA. BEE ORCHIS.

OPHRYS. *Lin. Gen. Plant.* ed. 3. GYNANDRIA DIANDRIA.

ORCHIS. *Rail Synops.* ed. 3. 379. HERBÆ BULBOSIS AFFINES.

OPHRYS. bulbis subrotundis, scapo folioso, nectarii labio quinquelobo; lobis subtus inflexis. *Hudson. Flor. Angl.* 340.

ORCHIS. radicibus subrotundis, labello holosericeo, emarginato, appendiculato. *Haller. Hist. Vol. 2. 1266. tab. 24.*
Duas species *apiferam* et *musciferam* HUDSONIS et HALLERI sub uno nomine *Insectiferæ* conjungit
CL. LINNÆUS.

Fuschii icon 360. *Baubin pin.* 83. *Gerard, emac.* 212.

RADIX. Bulbi duo, subrotundi, inæquales, radiculis longis vix fibrosis supra instructi.

CAULIS. Semipedalis aut pedalis, teres, *fig. 1.* foliosus.

FOLIA. Vaginantia, ovato lanceolata, subtus subargentea, fibris lineata, sæpe mutilata et fusca.

BRACTEÆ. *magnæ, vaginantes, virides, longitudine floris.*

FLORES. a tribus ad sex, spicati.

COROLLA. PETALA quinque, tria exteriora reliquis majora, ovata, concava, reflexa, purpurascens, ferioribus pallidioribus, subcarinata, carina viridi, *fig. 2;* duo interiora exterioribus quadruplo minora, angusta, *bisfuta, postice canaliculata, ad basin latiora, anteriorum extantia.*

NECTARIUM. Labellum amplum, leniter convexum, suborbiculatum, fusco sericeum, maculis flavis frequenter variegatum, quinquelobum, lobis inflexis, *fig. 3. laterales subtriangularibus, bisfatis; fig. 4. medio anteriorum productiore, apice recurvato flavo, fig. 5.*

Machina staminum five Stylus longa, subrecta, apice incurvata et sursum recurvata, *fig. 11.* antice bilocularis, loculis apertis, *fig. 12,* angustis, marginibus albis, membranaceis. *fig. 13.*

STAMINA. FILAMENTA duo, *fig. 6.* e squamula nectarifera ad basin Styli exentia, nutantia, Stigmati frequenter adherentia, *fig. 8.* basi glandula five globulo albo pellucido instructa, *fig. 7.* ANTHERÆ subrotundæ flavæ, *fig. 9.*

PISTILLUM. GERMEN oblongum, hexangulare, angulis obtusis rectis, STIGMA, *fig. 10.* melleo liquore obductum, cui particulæ antherarum frequenter adherent.

PERICARPIUM. CAPSULA oblonga, fusca, uncialis, *fig. 14,* unilocularis, *fig. 16,* trivalvis, valvis carinatis. *fig. 15.*

SEMINA plurima, minuta, oblonga, utraque extremitate membranacea, pellucida, reticulata, *fig. 18,* lente aucta, interiori parti carinæ longitudinaliter affixa, *fig. 17.*

ROOT. Two roundish, unequal bulbs, furnished at top with a few small longish fibres, but little branched.

STALK from half a foot to a foot high, round, *fig. 1,* leafy.

LEAVES embracing the stalk, of an oval pointed shape, underneath silvery, with linear fibres, frequently imperfect, and of a brown colour.

FLORAL LEAVES large, in the form of a sheath, green, and of equal length with the flowers.

FLOWERS from three to six growing in a spike.

COROLLA. five PETALS, the three exterior larger than the rest, oval, concave, turning back, purplish, somewhat keel shaped, the keel green, *fig. 2.* the latter flowering palest; the two interior four times smaller than the others, narrow, hairy, below behind, broadest at bottom, and projecting forward.

NECTARY. The lip of the Nectary, large, somewhat convex, roundish, of a silky brown colour, frequently variegated with yellow spots; having five lobes, the lobes bending underneath, *fig. 3.* The two side lobes, somewhat triangular and hairy, *fig. 4.* the middle of the anterior running out to a point, which turns back, and is of a yellow colour, *fig. 5.* the Style which in this plant supports the Stamina, long, upright, at the tip bending downwards and again upwards, *fig. 11.* anteriorly having two cavities which are open and narrow, *fig. 11.* the edges white and membranous, *fig. 13.*

STAMINA. two FILAMENTS, *fig. 6,* arising from the bottom of the style out a nectariferous scale, hanging down, frequently adhering to the Stigma, *fig. 8.* furnished at bottom with a small transparent gland or globule, *fig. 7.*—The ANTHERÆ roundish and yellow, *fig. 9.*

PISTILLUM. the GERMEN oblong, having six angles, the angles obtuse, not twisted, the STIGMA, *fig. 10,* covered with a viscid substance like honey, to which small particles of the Antheræ frequently adhere.

SEED VESSEL. a CAPSULE about an inch in length, oblong, brown, *fig. 14.* of one cavity, *fig. 16,* and three valves, the valves keel shaped. *fig. 15.*

SEEDS numerous, small, oblong; at each end membranous, transparent, and reticulated, *fig. 18.* magnified, affixed lengthwise to the inside of the keel of each valve, *fig. 17.*

Flowers in the Months of JUNE and JULY, the Seed is ripe the latter End of AUGUST.

Grows generally on chalky ground near woods, and sometimes in meadows; is become so rare about London, as scarcely to be found with any certainty.—Mr. Alborne informs me he has frequently gathered it in the pits behind Charlton church, and in the woods near Chiffelbury in Kent.—But it is often met with in plenty at a greater distance from town.

The root appears to possess the same virtues with those of the Orchis from which Salop is made, but being much smaller, is not worth cultivating on that account. The great resemblance which the flower bears to a Bee, makes it much sought after by Florists, whose curiosity indeed, often prompts them to exceed the bounds of moderation, rooting up all they find, without leaving a single specimen, to clear the heart of the Student in his botanic excursions. The best time of transplanting them is when they are in flower. This, with most of the other Orchids, was cultivated with the greatest success by the late PETER COLLINSON, Esq; (whose memory will always be revered by every Botanist) in his garden at Mill-hill.—His method was to place them in a soil and situation as natural to them as possible, and to suffer the grass and herbage to grow round them.

I have not yet heard of their being propagated by seed; it is to be wished that some intelligent Gardiner would exert himself in making some experiments to raise them in this way.

Botanists have often been at a loss in classing many plants, to find some resemblance by which they might distinguish that particular species; but in this plant the case is otherwise, the flower is so like the insect that gives it its name, that it strikes every beholder with admiration; what clerical purpose is intended by it, we do not at present know: Some future Observer may perhaps discover, for they who will examine Nature herself, “have much to see.”



Orchis pinnatifida





Asplenium scolopendrium.

ASPLENIUM SCOLOPENDRIUM. HARTSTONGUE.

ASPLENIUM *Lin. Gen. CRYPTOGRAMIA FILICES.*

Raii Synops. HERBÆ CAPILLARES ET AFFINES.

ASPLENIUM frondibus simplicibus cordato-lingulatis integerrimis, stipitibus hirsutis. *Lin. Sp. Pl. 1537.*

ASPLENIUM Frondes lanceolatæ, acuminatæ, basi cordatæ, integerrimæ, medio latiores. *Scopoli Fl. Can. n.*

ASPLENIUM petiolis hirsutis, folio longe lineari-lanceolato, integerrimo, circa petiolum exscisso. *Haller, 1. fl. n. 1695.*

HEMONITIS Fuschii *Icon. 294*

PHYLLITIS vulgaris *Clus. hist.*

SCOLOPENDRIA vulgaris *Tragi.*

LINGUA CERVINA officinarum *Baubin. pin. 350. Gerard. emac. 1138. Parkinson. 1046. Raii Synop. 116. Hudson. Fl. Angl. 384.*

RADIX perennis, fibrosissima, fusca, fibris fibrillis tenuissimis instructis.

ROOT perennial, exceedingly fibrous; the fibres brown, and furnished with other fibres, which are very minute.

STIPITES plures, pilosi.

STALKS numerous and mossy, or hairy.

FRONDES cordato-lingulatæ, longitudine pedales, latitudine fere bipollicares, glaberrimæ, margine undulato, nervo medio inferne piloso.

LEAVES tongue-shaped, at bottom cordate, about a foot in length, and one inch and a half in breadth, of a bright yellowish green colour and shining, the margin a little waved, and the mid-rib on the under side mossy.

FRUCTIFICATIO. Glomera linearia, obliqua, in pagina inferiore frondis nervo medio utrinque seriati disposita, *fig. 1. 2. 3.*

FRUCTIFICATION placed in oblique lines on the under side of the leaf, on each side of the mid-rib, *fig. 1. 2. 3.*

INVOLUCRUM. Squama linearis, bivalvis, longitudinaliter dehiscens, *fig. 2.*

INVOLUCRUM a linear membrane or case, of two valves, opening longitudinally, *fig. 2.*

CAPSULÆ numerosæ, subglobosæ, uniloculares, pedicellatæ, annulo elastico cinctæ, *fig. 5. 7. lente auctæ.*

CAPSULES numerous, standing on foot-stalks, nearly globular, surrounded by an elastic ring, and having one cavity, as they appear magnified, *fig. 5. 7.*

SEMINA numerosa, subrotunda, minutissima, *fig. 7. lente valde auctæ, fig. 8.*

SEEDS roundish, very numerous and minute, *fig. 7. as they appear through a greater magnifier, fig. 8.*

THIS is one of those plants which some Botanic Writers have called *Epiphyllisperma* from producing their seeds on the back of the leaves; LINNÆUS, includes it in his class, *Cryptogamia*, as neither Stamina nor Pistilla have yet been discovered on it. The first appearance of fructification that we observe are some little bags or cases of a yellowish or whitish green colour placed in rows on the under side of the leaves; *fig. 1.* on opening of which almost as soon as they become visible, we find the capsules or seed-vessels, *fig. 2.* very numerous, standing upright, and close together; at this time they appear of a green colour, as they approach towards maturity they change this for a deep brown; the cases then open lengthways in the middle, the two sides, by the protrusion of the capsules are turned quite back, and wholly disappear, *fig. 3.* This membranous substance or case, may be considered as similar to the *Calyptra* in mosses, or *Calyx* in other plants, and serves to secure and defend the tender seed and capsules, which being now become ripe, exhibit a most striking proof of that wisdom which the benevolent Author of Nature manifests in all the works of his creation.

Each capsule or seed-vessel consists of three parts, first the foot-stalk, *fig. 4.* which supports and connects them to the leaf; secondly, the jointed spring, *fig. 5.* which nearly surrounds the third part, or cavity containing the seeds, *fig. 6. 7.* The seeds being ripe, the cavity containing them is forced open by the elasticity of the jointed spring, and the seeds scattered and thrown to a considerable distance; one half of the cavity remains connected to one end of the spring, and the other half to the other end, *fig. 7.*

Some of the capsules being sooner ripe than others, discharge their seed sooner, so that it is a considerable time before they all become empty. On applying an entire row before the microscope for the first time, I was immediately struck with the motion that appeared in them, and afterwards found that the warmth of my breath occasioned a great number of the capsules to keep continually discharging their seeds, so as almost to give them the appearance of something alive. The closeness of the capsules to one another affording me but a confused idea of their structure, I separated them with the point of a penknife, from their connection to the leaf, and again placed them before the microscope, which then gave me a very different, and after a little examination, a very clear idea of their structure; many appeared with the seeds discharged, several in the act of discharging them, and some as yet entire; it frequently happened that while I was intently looking at one which I expected would open, at the instant of discharging it would be carried out of my sight by the strength and elasticity of the spring, and it was not 'till after repeated trials that I was able clearly to observe the manner of their opening. The seeds are very numerous, and scarcely visible to the naked eye; when magnified, they appear of a roundish figure, and full of little projecting points.

Both GRAY and SWAMMERDAM have given figures on this subject, but those of SWAMMERDAM are by much the most natural. As a great deal of the satisfaction in viewing objects of this kind, depends on the kind, as well as goodness of the microscope, that none of my Readers may be disappointed in the experiments they may make with this entertaining instrument, I may inform them, that the microscope I make use of is that which is sold in the shops, by the name of FLETCHER'S *Aquatic Microscope*, and which is made for this purpose with particular care and accuracy by GEORGE ADAMS of Fleet-Street, Mathematical Instrument Maker to his Majesty.

This plant may be found in seed from September to November in shady lanes and on walls, and is frequently found growing within-side of old wells. It is met with but rarely about town, though cultivated in most of our botanic gardens. The leaves are subject from a richness of soil to be much divided at their extremities, and very much curled at the edges.

It is an official plant, and is recommended by RAY from his own experience as a good medicine against convulsive disorders.

ALBION W. V. N. C. (1891)

1891

ALBION

INDUSTRIAL

1891

1891

1891


1891


1891

1891

1891

1891





Polypodium vulgare

POLYPODIUM VULGARE. COMMON POLYPODY.

POLYPODIUM *Linnei Gen. Pl.* CRYPTOGAMIA FILICES.

Frustrific. in punctis subrotundis sparsis per discum frondis;

Raii Syn. HERBÆ CAPILLARES ET AFFINES.

POLYPODIUM *vulgare* frondibus pinnatifidis : pinnis oblongis subferratis obtusis. *Linn. Syst. Vegetab.*
p. 786. *Fl. Suecic.* p. 373.

POLYPODIUM foliis pinnatis, lanceolatis, radice squamata. *Haller hist. n.* 1696.

POLYPODIUM *vulgare.* *Scopoli Fl. Carniol. n.* 1266.

POLYPODIUM *vulgare.* *Baubin. pin.* 359.

POLYPODIUM *vulgare.* *Parkinson* 1039.

POLYPODIUM *Gerard emac.* 1138. *Raii Syn. p.* 117, Polypody. *Hudson Fl. Angl. p.* 387.

RADIX oblique sub terræ superficie reptat, fibras suas ex tuberculis quibus plurimis scatet demittens, ad crassitudinem fere minimi digiti accedens, squamis fulcibus testæ, colore foris buxæ, intus fere herbaceæ, sapore dulci, tandem acerbo et adstringente.

STIPITES læves, interne sulcati.

FRONDES semipedales aut pedales, pinnatifidæ, pinnæ oblongæ, subferratæ, obtusæ, inferne pallidiores.

CAPSULÆ in acervulis, magnis, flavis, rotundis, nervo utrinque seriatim locatæ, pedicellatæ, subrotundæ, superficie granulata a feminibus protuberantibus, annulo elastico brevi instructæ, in valvulas duas dehiscentes, *fig.* 2, 3, 3, 5, 6.

SEMINA plurima, ovata, aut subreniformia flava, *fig.* 7, 8.

ROOT creeps obliquely under the surface of the earth, sending forth a number of fibres from little tubercles, which are plentifully distributed over its surface, about the thickness of the little finger, sometimes slenderer, covered with brown mossy scales, externally of a pale yellow colour, internally greenish, of a taste at first sweet, but finally sewerish and astringent.

STALKS smooth, grooved on the inner side.

LEAVES from half a foot to a foot in length, pinnatifid; the pinnæ oblong, slightly serrated, obtuse, paleish underneath.

CAPSULES placed in a row on each side the midrib of the leaf, in large, yellow, round dots, standing on foot-stalks, of a roundish shape, with the surface granulated from the seeds protuberating, furnished with a short elastic spring, and opening into two valves, *fig.* 2, 3, 4, 5, 6.

SEEDS numerous, oval or somewhat kidney-shaped, of a yellow colour, *fig.* 7, 8.

IN all those plants of the *Fern Tribe* which I have hitherto had an opportunity of examining, there appears to be much the same mechanism in their parts of fructification; one of the most striking and useful of which is the elastic ring which surrounds the Capsules, by means of which they are forced open and the seeds discharged. So necessary a part one should not conceive would be wanting in any of these plants, nor will it, I believe, be found to be so: yet many Botanists, and those too of eminence, not only deny its existence, but make the want of it a character to distinguish this Genus. GLEDITCH gives us the following as part of the generic character of the *Polypodium* "*Capsule annulo destituta.*" ADANSON also gives it the same character, "*sans anneau.*" It will perhaps not be difficult to account for this mistake; and at the same time it will shew us how injurious it is to science, for Authors to take things for granted without examining for themselves. In *TOURNEFORT's* elegant figures of the Genera, the Capsules of the *Polypodium* are represented without any ring: on the truth of these figures it is highly probable that those Authors have relied; for had they made use of their own eyes, assisted by a small magnifier, they could not have avoided seeing what *MALPIGHI* long before their time delineated, though rudely, and *GLEICHEN* since more elegantly figured.

There is one circumstance attending this species of *Polypodium*, which however does not run through the whole of this Genus, *viz.* the want of an Involucrum or Membrane; the little dots or assemblage of Capsules are not covered with any membrane; or if there be a membrane, it is very early deciduous, and not visible when the Capsules have arrived at a tolerable degree of maturity.

This species of *Polypodium* grows very common in woods and shady lanes on the old stumps of various trees; it differs much in size: sometimes it occurs on the Oak, in which case its virtue as a medicine has been more celebrated.

Its effects when taken inwardly are slightly purgative: it has been recommended in various disorders of the Viscera, in the Cachexy, swelling of the Spleen, Jaundice, obstructions of the Mesenteric Glands, Hypochondriac Disease, Cough, Asthma, &c. but it has generally been given with some other medicines.

In the present practice it is but little regarded.

1. The first part of the paper

is devoted to a general

discussion of the

principles of the

method of the

present paper


The second part of the paper is devoted to a detailed description of the method of the present paper. The method is based on the use of the principle of the least squares. The method is described in detail in the following sections.

3.

The third part of the paper

is devoted to a detailed

description of the





Bryum Leptocarpum

BRYUM SCOPARIUM. BROOM BRYUM.

BRYUM *Linnaei Gen. Pl. CRYPTOGAMIA MUSCI.*

Raii Syn. Gen. 3: MUSCI.

BRYUM *Scoparium*, Antheris erectiusculis, pedunculis aggregatis, foliis secundis recurvatis, caule declinato.

Linnaei Syst. Vegetab. p. 797:

HYPNUM foliis falcatis, heteromallis; vaginis multifloris. *Haller hist. n. 1777.*

HYPNUM *Scoparium. Scopoli Fl. Carn. p. 334. DIAGN. Florescentia Hyemalis. Folia arcuata, secunda, tenuia. Setae saepe plures.*

BRYUM *Scoparium: furculo declinato, ramoso, foliis secundis, recurvatis, primordialibus plumulosis. Necker. method. musc. p. 224.*

HYPNUM *Scoparium. Weis. Cryptogam. p. 213.*

BRYUM reclinatum, foliis falcatis, scoparum effigie: *The sickle-leaf'd bending Beasom Bryum. Dillen. musc. p. 357. tab. 46. fig. 16*

BRYUM erectis capitulis angustifolium, caule reclinato. *Cat. Gifs. 222. Raii Syn. 95. Hudson Fl. Angl. p. 406.*

CAULES unciales aut biunciales et ultra, suberecti, ramosi, in denso caespite collecti, fordide rufi, infra multo tomento fusco obfiti.

FOLIA caulem inequaliter circumstant, hinc in quibusdam locis nudiusculus relinquitur, in aliis foliis crebrioribus vestitur, praecipue ad apicem, longa, linearia, acuminata, canaliculata, *fig. 1*, recurvata, secunda.

PEDUNCULI unciales aut biunciales, ad basin rubicundi, erecti, ex uno latere caulium plerumque oriuntur, aliquando vero ex apice, ut plurimum foliarii, subinde vero duo ex eodem perichætio proveniunt, basi bulbillo cylindraceo terminati, *fig. 7*, qui foliis pluribus latiusculis, pilo terminatis, acù facile separabilibus includitur, *fig. 8, 9.*

CAPSULÆ oblongæ et fere cylindraceæ, nunc erectæ, nunc paululum incurvatae, *fig. 3*; OPERCULUM rostratum, tenue, longitudine capsulae et concolor, *fig. 4*; ORA ciliata five denticulata, *fig. 5*; CALYPTRA straminea, longitudine Capsulae, postquam medio disrumpitur, basi suo capfulam arcte cingit, *fig. 2*; POLLEN viride, *fig. 6.*

STALKS from one to two inches high and more, nearly upright, branched, growing thickly together, of a dirty red colour, and covered at bottom with a dark brown woolly substance.

LEAVES: the leaves cover the stalk unequally, hence in some places it is left rather naked, in others more thickly covered with leaves, particularly towards the top, are long, linear, pointed, grooved, *fig. 1*, bent back, and turning all one way.

FOOT-STALKS an inch or two inches high, towards the bottom reddish, upright, arising generally from the side of the stalks, but sometimes from the top, most commonly single, but now and then two proceed from the same perichætium, furnished at bottom with a cylindrical bulb, *fig. 7*, which is inclosed by many broadish leaves, terminating in a hair, and easily separated by a needle, *fig. 8, 9.*

CAPSULES oblong and almost cylindrical, sometimes upright, sometimes a little incurvated, *fig. 3*; the OPERCULUM the length of the Capsule, and of the same colour, terminating in a long slender point, *fig. 4*; the MOUTH ciliated or furnished with little teeth, *fig. 5*; the CALYPTRA straw-coloured, the length of the Capsule, after bursting in the middle closely embracing the Capsule by its base, *fig. 2*; the POLLEN green, *fig. 6.*

DILLENIUS very justly remarks, that this Moss seems to partake of the nature of both *Bryum* and *Hypnum*, but in his opinion, it comes nearest to the *Bryum*, and of the same sentiment appear to be LINNÆUS and NECKER, while HALLER, SCOPOLI, and WEIS, rank it among the *Hypnums*, and this they are led to, chiefly from the Peduncles being furnished at bottom with a kind of *Perichætium*; but DILLENIUS very properly observes, that although the peduncle is surrounded at bottom by many *squamæ* or *folioli*, yet these are not similar to what occur in the generality of *Hypnums*, as they may with the point of a pin be easily separated from one another, and then the base of the peduncle appears to be furnished with a *bulbillus* as in most of the *Bryums*: this circumstance added to its general habit, appears fully to justify this most excellent Botanist in placing it with the *Bryums*, from whence it ought not to have been separated without more weighty reasons than have been advanced.

This Moss distinguishes itself from most others by its beautifull and lively verdure; when young it puts on a very different appearance from what it has when farther advanced, being much shorter and its leaves upright; and DILLENIUS has sometimes remarked in this species *Stellula feminea*.

It grows in very large Clumps or Patches forming a soft and delightfull Carpet, on the banks which surround woods, at the bottom of trees, and on heaths.

It is found on some parts of Hamptstead heath producing its fructifications in February and March.

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION

CHICAGO, ILL., U.S.A.

1914

Published by the American Medical Association, 535 North Dearborn Street, Chicago, Ill., U.S.A.
Subscription price, \$5.00 per annum in advance. Single copies, 15 cents.
Entered as Second-Class Matter, May 2, 1912, Post Office at Chicago, Ill., under No. 102,363.
Acceptance for mailing at special rate of postage provided for in Act of October 3, 1917, authorized on July 1, 1918.
Postage paid at Chicago, Ill.

Subscription price, \$5.00 per annum in advance.

Single copies, 15 cents.

Entered as Second-Class Matter, May 2, 1912, Post Office at Chicago, Ill., under No. 102,363.
Acceptance for mailing at special rate of postage provided for in Act of October 3, 1917, authorized on July 1, 1918.

Postage paid at Chicago, Ill.

Postage paid at Chicago, Ill.

1914

Published by the American Medical Association, 535 North Dearborn Street, Chicago, Ill., U.S.A.
Subscription price, \$5.00 per annum in advance. Single copies, 15 cents.
Entered as Second-Class Matter, May 2, 1912, Post Office at Chicago, Ill., under No. 102,363.
Acceptance for mailing at special rate of postage provided for in Act of October 3, 1917, authorized on July 1, 1918.
Postage paid at Chicago, Ill.

It is the policy of the Association to publish only original researches, and to exclude all material that is purely editorial, or that is of a general nature, or that is of a purely local interest. The Association is not responsible for the opinions or views expressed by its members, or for the results of their experiments, or for the accuracy of their statements. The Association is not responsible for the results of the experiments of its members, or for the accuracy of their statements. The Association is not responsible for the results of the experiments of its members, or for the accuracy of their statements.


Published by the American Medical Association, 535 North Dearborn Street, Chicago, Ill., U.S.A.

Subscription price, \$5.00 per annum in advance. Single copies, 15 cents.

Entered as Second-Class Matter, May 2, 1912, Post Office at Chicago, Ill., under No. 102,363.

Acceptance for mailing at special rate of postage provided for in Act of October 3, 1917, authorized on July 1, 1918.

Postage paid at Chicago, Ill.



Elodea canadensis



BRYUM UNDULATUM. CURLED BRYUM.

BRYUM *Linnei. Gen. Pl. CRYPTOGAMIA Musci.*

Raii Syn. Gen. 3. Musci.

BRYUM (*undulatum*) antheris erectiusculis, pedunculis subfoliariis, foliis lanceolatis carinatis undulatis patentibus ferratis. *Linne. Syst. Vegetab. p. 797.*

BRYUM foliis lanceolatis ferratis, capsulis cylindricis inclinatis aristatis. *Haller. hist. tom. 2. 1823.*

BRYUM *phylitidisfolium*: furculo simplici, foliis undato-ferratis, primordiis plumulosi. *Necker's method. muscor. p. 203.* cur nomen triviale a Cl. Neckero mutaretur non video, cum analogia unde nomen ejus sumitur obscura fit, observante Cl. Scopoli.

BRYUM Phylitidis folio rugoso acuto, capsulis incurvis. *Dillen musc. 360. tab. 46. fig. 18.*

BRYUM undulatum. *Scopoli. Fl. Carniol. n. 1301. Raii Syn. p. 95. 16. Hudson Fl. Angl. 466. Weiss Cryptogam. 196. Oeder Fl. Dan. tab. 497. nostris duplo saltem minor, cum operculo nimis recto et acuto.*

SURCULI uncialis, aut biunciales, plerumque simplices, erecti, foliosi.

FOLIA lanceolata, undulata, carinata, *ferrato-aculeata*, patentia, arefactione involuta, *fig. 1.*

PEDUNCULI simplices, (duo ex eodem furculo nonnunquam proveniunt) furculis plerumque longiores, erecti, rubri, *fig. 2.*

CAPSULA sive ANTHERA cylindracea, incurvata, lente visa substriata, primum viridis, dein ex livido-fusca, demum rufa, *fig. 3.* Basis OPERCULI hemisphærica, rubra, apex pallida, setacea, obtusifuscula, *fig. 5.* Capsulæ ORA ciliata CILIIS inflexis, *fig. 7.* ANNULUS ruber, *fig. 6.* POLLEN seu SEMEN viride, *fig. 8.*

CALYPTRA pallide fusca, acuminata, primum erecta, flexurâ capsulæ dirumpitur, et recta manet, basique suâ a Capsula secedit, *fig. 4.*

STALKS from one to two inches high, generally simple, upright and leafy.

LEAVES lanceolate, waved, keel-shaped, *minutely and sharply ferrated at the edges*, spreading, when dry curling in, *fig. 1.*


FOOT-STALKS of the fructification simple, (sometimes two proceed from the same stalk) generally longer than the stalks, upright, and of a reddish colour, *fig. 2.*

CAPSULE or ANTHERA cylindrical, incurvated, if magnified appearing somewhat striated; first green, then livid-brown, and lastly of a reddish brown colour; *fig. 3.* the bottom of the OPERCULUM hemispherical and red, the top paler, very slender and rather blunt; *fig. 5.* the MOUTH of the Capsule furnished with CILIÆ which bend inward, *fig. 7.* the ANNULUS or RING red, *fig. 6.* the POLLEN or SEED green, *fig. 8.*

CALYPTRA of a pale brown colour, and terminating in a long point, first upright, afterwards by the bending of the Capsule it becomes burst at bottom, and remains strait, with its base at some little distance from the Capsule.

THIS species of BRYUM is one of the largest we have in this Country, it produces its fructification from November to February and may be found in most of the woods near Town, as well as on the heaths, but more particularly in *Charlton Wood*, where it abounds.

As all its parts of fructification are large and distinct, the botanic Student who would investigate this difficult class of plants, cannot with this view, select any moister more proper for his purpose.





Bryum hernandezii.

BRYUM HORNUM. SWANS-NECK BRYUM.

MNIUM *Linnaei Gen. Pl. CRYPTOGAMIA MUSCI.* Masculus flos pedunculatus. Feminus flos in distincto sæpius individuus.

Raii Synopsis Gen. 3. MUSCI.

MNIUM *hornum* antheris pendulis, pedunculo curvato, furculo simplici, foliolis margine scabris. *Linnaei Syst. Vegetab. 796.*

BRYUM *hornum* furculo capitulifero ramosiusculo: stellifero simplici, primordialibus plumulosis. *Necker. Method. Musc. p. 215.*

MNIUM foliis lanceolatis, imbricatis, capulis pendulis cylindricis obtusis. *Haller. hist. helv. 3. p. 54.*

MNIUM *hornum* ferratifolium. *Weis Cryptogam. 149.*

BRYUM antheris oblongis nutantibus, pedunculo curvato, foliolis ovatis, margine scabris. *Hudson. Fl. Angl. p. 415.*

BRYUM stellare *hornum* sylvarum, Capulis magnis nutantibus. *Dillen. musc. 402.*

BRYUM nitidum capitulis majoribus reflexis, calyptra imum vergente, pediculis oblongis e cauliculis novis egredientibus. *Raii Syn. p. 102. 51.*

Ad majores accedit hæc species.

CAULES unciales aut biunciales, radiculis ferrugineis, valde tomentosis instructi, erecti, plerumque ramosi, pedunculiferi et stelliferi, ad basin rubicundi, STELLULÆ et PEDUNCULI, nunc seorsim, nunc ex eadem radice proveniunt, unusque aut plures Surculi e basi caulis semper fere nascuntur.

FOLIA faturate viridia, ovato-lanceolata, suberecta, pellucida, ad lentem minute serrata, fig. 1; nervo medio distincto et in mucrone brevem educto, in furculis femineis dictis apice stellatim expansa, et paulo latiora, in junioribus angustiora et cauli magis adpressa.

PEDUNCULI terminales, biunciales, rubræ, versus apicem ut recte observavit DILLENIIUS instar colli coloris incurvati.

CAPSULÆ oblongæ, tumidæ, virides, nutantes, lente aucta, fig. 7; per longitudinem secta ut RECEPTACULUM conspicitur, fig. 9; CALYPTRA longa, acuminata, caduca, fig. 6; OPERCULUM breve, flavescens, fig. 8; ORA ciliata.

This species comes near to the largest size.

STALKS from one to two inches in height, furnished with roots which are of a ferruginous colour, and covered with a kind of woolly substance, upright and generally branched, reddish at bottom, producing both PEDUNCULI and STELLULÆ, which proceed sometimes from the same, sometimes from different roots, and one or several Surculi usually spring from the bottom of the stalk.

LEAVES of a deep green colour, of an oval pointed shape, nearly upright, pellucid, when viewed with a glass finely serrated at the edges, fig. 1; the midrib distinct, and terminating in a short point, on the tops of those stalk, which are considered as female, they are expanded like a little star and somewhat broader, in the young shoots they are narrower and pressed closer to the stalk.

PEDUNCLES springing from the summit of the stalks, about two inches in height, bent near the top like a Swans Neck as DILLENIIUS has properly observed.

CAPSULES oblong, tumid, of a green colour and drooping, magnified, fig. 7; cut longitudinally through the middle that the RECEPTACULUM may be seen, fig. 9; the CALYPTRA long, pointed, and soon falling off, fig. 6; the OPERCULUM short, of a yellowish colour, fig. 8; the MOUTH of the Capsule ciliated.

On examining with a Microscope the tops of those Stalks which are called *Stellulæ femineæ*, fig. 2, and which are considered by many as the female parts of the fructification in this Moss, there appeared in the center of the *Stellula*, a great number of small upright bodies, or Corpuscles, of two kinds, fig. 3, the one white, pellucid, and jointed; the other of a greener colour, shorter, and of an oblong oval shape, vid. fig. 4, 5. They do not appear to me to have any thing in their Structure, in the least similar to any of the parts of fructification in plants, what their real structure and uses are, may perhaps be discovered by future observations.

This species occurs not unfrequently on moist banks in Woods, as in *Charlton Wood*, and the Woods about *Hamstead*, producing its Fructifications in February and March.

As the *Capitula pulverulenta* of DILLENIIUS, or *Sphaerophylli* as they are called by NECKER, are entirely wanting in this Moss, and as the existence of those singular little heads seems very obviously to distinguish the Genus *Mnium*, I have chosen rather to arrange it with DILLENIIUS and HUDSON among the *Bryums*, than with LINNÆUS among the *Mniums*; for if we make *Mniums* of all the Mosses which have *Stellulæ*, we shall involve ourselves in considerable difficulties: many of those *Stellulæ* are indeed very obvious, as in the present one, but in others they are very obscure, so that it is difficult to say whether they exist in them or not; but if they were obviously to be distinguished, there is not the least likeness between a *Stellula* and *Sphaerophyllum*, why then unite in one Genus plants which have such very different appearances? Would it not be better to consider the Mosses which produce *Sphaerophylli* or little balls as *Mniums*, according to DILLENIIUS, and divide the *Bryums*, if necessary, into two families, viz. such as have obvious *Stellulæ*, and such as have none?

The name of rough *Bryum*, which Mr. HUDSON seems to have given to this Moss for brevity's sake, conveys an idea with which this *Bryum* does not seem perfectly to correspond, it having no roughness except at the edges of the leaves, which are minutely serrated: I have therefore adopted DILLENIIUS's name of *Swans Neck Bryum*, as being justifiable from the singular shape of the Peduncles, and being more likely to be remembered from its striking analogy.

THE HISTORY OF THE

REIGN OF

CHARLES THE FIRST

BY

JOHN BURNET

OF

THE UNIVERSITY OF OXFORD

IN TWO VOLUMES

LONDON

Printed by J. Streater, at the Sign of the Gun, in St. Dunstons Church-yard

1679

THE SECOND VOLUME

OF


THE

SAME

WORK

BY

JOHN BURNET



HYPNUM PROLIFERUM. PROLIFEROUS HYPNUM.

HYPNUM *Linnaei* Gen. Pl. CRYPTOGRAMIA MUSCI.

Raii Syn. Gen. 3. Musci.

HYPNUM *proliferum* furculis proliferis, plano-pinnatis, pedunculis aggregatis. *Linnaei* Syst. Vegetab. p. 802.

HYPNUM ramis teretibus pinnatis, pinnulis pinnatis, foliis adpressis. *Haller. hist.* 3. p. 33.

HYPNUM filicinum, Tamarisci foliis minoribus, non splendentibus. *Dillen.* p. 276. icon. 35. fig. 14.

HYPNUM repens filicinum minus, luteo virens. *Catal. Gifs.* 217. *Raii Synop.* p. 86. n. 36. *Hudson, Fl.*

Angl. p. 422. *Wetis Cryptogam.* p. 230.

CAULES palmares ad dodrantales, repentes, hinc inde radículas fufcas exferentes, fepe vero adeo intricate connexi ut humi íerpere nequeant, foliis ovato-acuminatis, carinatis, mucronatis, fparfe teñtis, fig. 1. horum foliorum fuperficies, microfcopio valde aucta granulofa apparet, fig. 2.

RAMI pulchre pinnati, deflexi, virefcetes, ad luteum colorem plus minufve accedentes pro ratione fitus aut anni temporis, omni fplendore deftituti, rachis concolor, ad extremitatem plerumque ineraffatus. **RAMULI** et **PINNULÆ** foliis exiftimis, confertis, nudo oculo vix confpiciuis imbricatim teñti; e difco rami, aut frondis, novus caulis aut furculus plerumque exurgit, unde plantula mire extenditur ac propagatur, et hinc *Prolifer* vocatur.

PEDUNCULI feíquiunciales, rubri, plerumque quatuor aut quinque, aliquando plures e caule aggregatim affurgunt, et in quibufdam caulibus, *Perichætia* plura aut potius eorum rudimenta occurrunt, e quibus *Pedunculi* fequente anno probabiliter nafcuntur.

PERICHÆTIUM fig. 3. aut bafis pedunculi, ovatum, foliolis tenuibus pilo longo flexuofe terminatis veftitum. **CAPSULÆ** five **ANTHÆRÆ** fig. 4. quæ Semen aut Pollinem continent, incurvatæ, ex fulco aurantiacæ. **OPERCULUM** fig. 6. (quod collo capfulæ infigitur, et femine matrefcente decidit) breve, et acuminatum. **Orbículum** Capfulæ duplici ferie **Ciliarum** inítruitur fig. 8. 9. **CILLÆ** exteriores fig. 8. aurantiacæ, divergentes, apicibus aliquando paululum inflexis, et cum aridæ íint fragiles; interiores fig. 9. convergentes, membrana reticulata connexæ, ad quam videndam microfcopio opus eft. **POLLEN** five **SEMEN** viride. **CALYPTRA** fig. 5. quæ anthera cum fuo Operculo partim tegitur et quæ primum decidit albida eft.

STALKS from three to nine inches in length, creeping on the ground, and here and there fending forth fmall brown fibres, but very often fo intricately connected together as to be hindered from creeping, thinly covered with leaves of an oval pointed fhape, having a ftrong midrib, which runs out to a fine point fig. 1. when greatly magnified the furface of thefe leaves exhibits a granulated appearance fig. 2.

BRANCHES beautifully pinnated, and bending downward, of a green colour, more or lefs inclined to yellow, according to its place of growth, and the feafon of the year, without any glifs; the midrib of the fame colour with the leaves and generally thicker at its extremity; the fmall leaves, laying one over another, and fcarce difcernible to the naked eye. From the middle of the branch or Fronds moft commonly arifes a new ftalk, or furculus, by which means this plant is fingularly extended and propagated, and from this circumftance it acquires the name of *Proliferous*.

PEDUNCLES about an inch and a half in length, of a bright red colour, generally about four or five, fometimes more, fpring from the ftalk nearly together, in fome of the ftalks there is the appearance of feveral *Perichætia* without peduncles, which probably arife from them the next year. The **PERICHÆTIUM** fig. 3. which is the bafe of the peduncle, is of an oval fhape, and covered with fmall leaves which terminate in a long flexible point. The **CAPSULES** or **ANTHÆRÆ** containing the pollen or feed fig. 4. are incurvated, and of a brown or orange colour. The **OPERCULUM** fig. 6. (which fits on to the top of the Capfulæ, and when the feed contained within it, is ripe, falls off) is fhort, and pointed; the mouth of the Capfulæ has two rows of **CILLÆ** fig. 8. 9; the exterior row fig. 8. orange coloured, and diverging, the tops of them fometimes bending a little inward, and brittle when dry, the interior row fig. 9. converging, of a membranous texture, and when very much magnified, appearing reticulated. The **POLLEN** or **SEED** contained within the Capfulæ is green. The **CALYPTRA** fig. 5. which partly covers the anthera and operculum, and firts drops off is of a white colour.

There is fcarce a Wood in the environs of this City, on the borders of which this elegant fpecies of Mofs doth not occur.

It produceth its fruítifications from December to February; in this ftate however it is but feldom met with, yet may be found by diligent fearching. *Linnaeus* in one of his journeys through Sweden, obferved this Mofs growing in the thickeft Woods, obfcured with perpetual fhade, and where all other plants perifhed.

Moft of the writers who have made this clafs of plants more particularly the object of their enquiries, have generally made two diftinct Genera of the Hypnum and Bryum, yet fo great is the affinity betwixt them, and fo much do they run into one another, that what fome of thefe Authors call a Bryum, others denominate a Hypnum; indeed this divifion feems adopted more to facilitate the investigation of the plants of this numerous family, than from any real natural divifion which takes place between them. The difference between fome of the Hypnums and fome of the Bryums is obvious to almoft every one, but to afcertain the limits where the one begins and the other terminates, feems a talk too difficult for the moft accurate Botanift.

The principal Charactersiftics of a Bryum according to *Linnaeus*, are, that the peduncle which fufains the Anthera or Capfulæ, grows out of the top of the furculus or ftalk, and is furnifhed at its bafe with a little naked tubercle or bulb; in the Hypnum on the contrary, the peduncle grows out of the fide of the ftalk and the tubercle at its bafe is covered with leaves and called a *Perichætium*.

Hypnum Poliferum.



